

ANALYSIS I EXAMPLES 1

G.P. Paternain Lent 2011

Comments on and/or corrections to the questions on this sheet are always welcome, and may be e-mailed to me at g.p.paternain@dpms.cam.ac.uk.

1. Prove that if $a_n \rightarrow a$ and $b_n \rightarrow b$ then $a_n + b_n \rightarrow a + b$.
2. Sketch the graphs of $y = x$ and $y = (x^4 + 1)/3$, and thereby illustrate the behaviour of the real sequence a_n where $a_{n+1} = (a_n^4 + 1)/3$. For which of the three starting cases $a_1 = 0$, $a_1 = 1$ and $a_1 = 2$ does the sequence converge? Now prove your assertion.
3. Let $a_1 > b_1 > 0$ and let $a_{n+1} = (a_n + b_n)/2$, $b_{n+1} = 2a_n b_n / (a_n + b_n)$ for $n \geq 1$. Show that $a_n > a_{n+1} > b_{n+1} > b_n$ and deduce that the two sequences converge to a common limit. What limit?
4. The real sequence a_n is bounded but does not converge. Prove that it has two convergent subsequences with different limits.
5. Investigate the convergence of the following series. For those expressions containing the complex number z , find those z for which convergence occurs.

$$\sum_n \frac{\sin n}{n^2} \qquad \sum_n \frac{n^2 z^n}{5^n} \qquad \sum_n \frac{(-1)^n}{4 + \sqrt{n}} \qquad \sum_n \frac{z^n(1-z)}{n}$$

6. Show that $\sum 1/(n \log^\alpha n)$ converges if $\alpha > 1$ and diverges otherwise. Does $\sum 1/(n \log n \log \log n)$ converge?
7. Consider the two series $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \dots$ and $1 + \frac{1}{3} - \frac{1}{2} + \frac{1}{5} + \frac{1}{7} - \frac{1}{4} + \dots$, having the same terms but taken in a different order. Let s_n and t_n be the corresponding partial sums to n terms. Show that $s_{2n} = H_{2n} - H_n$ and $t_{3n} = H_{4n} - \frac{1}{2}H_{2n} - \frac{1}{2}H_n$, where $H_n = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{n}$. Show that s_n converges to a limit s and that t_n converges to $3s/2$.
8. Suppose that $\sum a_n$ diverges and $a_n > 0$. Show that there exist b_n with $b_n/a_n \rightarrow 0$ and $\sum b_n$ divergent.
9. (Abel's test.) Let a_n and b_n be two sequences and let $S_n = \sum_{j=1}^n a_j$ and $S_0 = 0$. Show that for any $1 \leq m \leq n$ we have:

$$\sum_{j=m}^n a_j b_j = S_n b_n - S_{m-1} b_m + \sum_{j=m}^{n-1} S_j (b_j - b_{j+1}).$$

Suppose now that b_n is a decreasing sequence of positive terms tending to zero. Moreover, suppose that S_n is a bounded sequence. Prove that $\sum_{j=1}^{\infty} a_j b_j$ converges. Deduce the alternating series test.

Does the series $\sum_{n=1}^{\infty} \frac{\cos(n)}{n}$ converge or diverge?

10. For $n \geq 1$, let

$$a_n = \frac{1}{\sqrt{n}} + \frac{(-1)^{n-1}}{n}.$$

Show that each a_n is positive and that $\lim a_n = 0$. Show also that $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$ diverges. [This shows that, in the alternating series test, it is essential that the moduli of the terms decrease as n increases.]

11. Let $z \in \mathbb{C}$. Show that the series

$$\frac{z}{1-z^2} + \frac{z^2}{1-z^4} + \frac{z^4}{1-z^8} + \frac{z^8}{1-z^{16}} + \cdots$$

converges to $z/(1-z)$ if $|z| < 1$, converges to $1/(1-z)$ if $|z| > 1$, and diverges if $|z| = 1$.

12. Prove that every real sequence has a monotonic subsequence. Deduce the Bolzano-Weierstrass theorem.

13. Can we write the open interval $(0,1)$ as a disjoint union of closed intervals of positive length?

14. In lectures (or problem 12) you learned a proof of the Bolzano-Weierstrass theorem based on the fundamental axiom (every increasing sequence bounded above converges). Show that the Bolzano-Weierstrass theorem is in fact equivalent to the fundamental axiom, that is, give a proof of the fundamental axiom assuming Bolzano-Weierstrass.