

Stochastic Financial Models

J.R. Norris

August 26, 2026

Contents

1	Martingales	3
1.1	Conditional probabilities and expectations	3
1.2	Definitions	6
1.3	Examples	7
1.4	Optional stopping	8
2	Pricing contingent claims	11
2.1	Multi-period asset price model	11
2.2	Contingent claims	12
2.3	Equivalent probability measures	13
2.4	Arbitrage	13
2.5	Fundamental theorem of asset pricing	15
2.6	Pricing attainable claims	16
2.7	Pricing non-attainable claims in a single-period model	17
2.8	Completeness	17
2.9	Binomial model	17
2.10	Pricing barrier options for a special binomial model	21
2.11	American calls and puts	22
3	Brownian motion	24
3.1	Definition and basic properties	24
3.2	Wiener's theorem	26
3.3	Brownian motion as a limit of random walks	26
3.4	Change of probability measure	27
3.5	Reflection principle	27
3.6	Hitting probabilities	29
3.7	Relation to the heat equation	29
3.8	Transition density for killed Brownian motion	30
4	Black–Scholes model	31
4.1	Black–Scholes pricing formula	31
4.2	Black–Scholes PDE	32
4.3	Binomial approximation to Black–Scholes	33
4.4	Black–Scholes formula for the price of a European call	35
4.5	Sensitivities for contingent claims in the Black–Scholes model	37
4.6	Implied volatility	37
4.7	Pricing of exotic options in the Black–Scholes model	38
5	Appendix	39

1 Martingales

1.1 Conditional probabilities and expectations

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. Recall that, given an event B of positive probability, we define the conditional probability $\mathbb{P}(\cdot|B)$ by

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}, \quad A \in \mathcal{F}.$$

The associated expectation is written $\mathbb{E}(\cdot|B)$. If a random variable X is integrable (with respect to $\mathbb{P}$), then X is also integrable with respect to $\mathbb{P}(\cdot|B)$ and the following formula holds

$$\mathbb{E}(X|B) = \frac{\mathbb{E}(X1_B)}{\mathbb{P}(B)}.$$

In this course we will use also some further notions of conditional probability and conditional expectation. These are associated not to an event B but to a sub- σ -algebra $\mathcal{G}$ of $\mathcal{F}$. In general, these notions have to be approached indirectly, by the following theorem.

Theorem 1.1. *Let $\mathcal{G}$ be a sub- σ -algebra of $\mathcal{F}$ and let X be an integrable random variable. Then there exists an integrable random variable Y with the following properties:*

- (a) Y is $\mathcal{G}$ -measurable,
- (b) $\mathbb{E}(Y1_A) = \mathbb{E}(X1_A)$ for all $A \in \mathcal{G}$.

Moreover, if Y' is another integrable random variable satisfying (a) and (b), then $Y' = Y$ almost surely. Moreover, the same statements are valid if we replace ‘integrable’ by ‘non-negative’, meaning $[0, \infty]$ -valued, throughout.

We call any such random variable Y (a version of) the conditional expectation of X given $\mathcal{G}$ and we write

$$Y = \mathbb{E}(X|\mathcal{G}) \quad \text{almost surely.}$$

In the case where $\mathcal{G} = \sigma(Z)$ for some random variable Z , we also write $Y = \mathbb{E}(X|Z)$ almost surely. In the case where $X = 1_A$ for some event A , we also write $Y = \mathbb{P}(A|\mathcal{G})$ almost surely.

The following argument proves the uniqueness statement in Theorem 1.1, and also establishes a useful monotonicity property. Let X' be another integrable random variable such that $X \leq X'$ almost surely. Suppose that Y' is an integrable random variable which satisfies (a) and (b) with respect to X' . Set $A = \{Y \geq Y'\}$ and consider the non-negative random variable $Z = (Y - Y')1_A$. Then, since $A \in \mathcal{G}$,

$$\mathbb{E}(Y1_A) = \mathbb{E}(X1_A) \leq \mathbb{E}(X'1_A) = \mathbb{E}(Y'1_A)$$

so $\mathbb{E}(Z) = \mathbb{E}(Y1_A) - \mathbb{E}(Y'1_A) \leq 0$ and so $Z = 0$ almost surely, which implies that $Y \leq Y'$ almost surely. In the case $X = X'$, we deduce by symmetry that $Y = Y'$ almost surely, as claimed.

We will omit proof of the existence statement in Theorem 1.1. A proof may be found in lecture notes for Advanced Probability. However, in the case where $\mathcal{G}$ is discrete, we can construct a suitable random variable Y directly, as we now show. Consider the case where

$$\mathcal{G} = \sigma(B_n : n \in \mathbb{N}) = \left\{ \bigcup_{n \in I} B_n : I \subseteq \mathbb{N} \right\}$$

for some sequence $(B_n : n \in \mathbb{N})$ of disjoint events whose union is Ω . Given an integrable random variable X , set

$$Y = \sum_{n \in \mathbb{N}} \mathbb{E}(X|B_n)1_{B_n}$$

where we make the convention that $\mathbb{E}(X|B_n) = 0$ whenever $\mathbb{P}(B_n) = 0$. Then Y is $\mathcal{G}$ -measurable and, by monotone convergence,

$$\mathbb{E}(|Y|) = \mathbb{E} \left(\sum_{n \in \mathbb{N}} |\mathbb{E}(X|B_n)|1_{B_n} \right) = \sum_{n \in \mathbb{N}} \mathbb{E}(|\mathbb{E}(X|B_n)|1_{B_n}) \leq \sum_{n \in \mathbb{N}} \mathbb{E}(|X|1_{B_n}) = \mathbb{E}(|X|) < \infty.$$

so Y is integrable. Moreover, for $A = \bigcup_{n \in I} B_n \in \mathcal{G}$, by dominated convergence,

$$\mathbb{E}(Y1_A) = \mathbb{E} \left(Y \sum_{n \in I} 1_{B_n} \right) = \sum_{n \in I} \mathbb{E}(Y1_{B_n}) = \sum_{n \in I} \mathbb{E}(X|B_n)\mathbb{P}(B_n) = \sum_{n \in I} \mathbb{E}(X1_{B_n}) = \mathbb{E}(X1_A)$$

where we used $|Y|$ and $|X|$ as dominating random variables for the second and last equalities respectively. Hence Y has both properties (a) and (b) of Theorem 1.1, and we have shown that

$$\mathbb{E}(X|\mathcal{G}) = \sum_{n \in \mathbb{N}} \mathbb{E}(X|B_n)1_{B_n} \quad \text{almost surely.}$$

We now examine some general properties of conditional expectation.

Proposition 1.2. *Let $\mathcal{G}$ be a sub- σ -algebra of $\mathcal{F}$ and let X and W be integrable random variables. Then*

- (i) $\mathbb{E}(\mathbb{E}(X|\mathcal{G})) = \mathbb{E}(X)$,
- (ii) if X is $\mathcal{G}$ -measurable, then $\mathbb{E}(X|\mathcal{G}) = X$ almost surely,
- (iii) if X is independent of $\mathcal{G}$, then $\mathbb{E}(X|\mathcal{G}) = \mathbb{E}(X)$ almost surely,
- (iv) if $X \geq 0$ almost surely, then $\mathbb{E}(X|\mathcal{G}) \geq 0$ almost surely,
- (v) $\mathbb{E}(\alpha X + \beta W|\mathcal{G}) = \alpha \mathbb{E}(X|\mathcal{G}) + \beta \mathbb{E}(W|\mathcal{G})$ almost surely, for all $\alpha, \beta \in \mathbb{R}$.

Proof. Properties (i) to (iii) follow easily from the defining properties (a) and (b), while (iv) follows from the monotonicity property shown above (take $X = 0$ and $X' = X$). To show (v), set $Y = \alpha \mathbb{E}(X|\mathcal{G}) + \beta \mathbb{E}(W|\mathcal{G})$. Then Y is $\mathcal{G}$ -measurable and, for all $A \in \mathcal{G}$,

$$\mathbb{E}(Y1_A) = \alpha \mathbb{E}(\mathbb{E}(X|\mathcal{G})1_A) + \beta \mathbb{E}(\mathbb{E}(W|\mathcal{G})1_A) = \alpha \mathbb{E}(X1_A) + \beta \mathbb{E}(W1_A) = \mathbb{E}((\alpha X + \beta W)1_A).$$

Hence $Y = \mathbb{E}(\alpha X + \beta W|\mathcal{G})$ almost surely, as claimed. $\square$

Proposition 1.3 (Tower property). *Let $\mathcal{G}$ and $\mathcal{H}$ be sub- σ -algebras of $\mathcal{F}$ with $\mathcal{H} \subseteq \mathcal{G}$. Let X be an integrable random variable. Then*

$$\mathbb{E}(\mathbb{E}(X|\mathcal{G})|\mathcal{H}) = \mathbb{E}(X|\mathcal{H}) \quad \text{almost surely.}$$

Proof. Choose a version Y of $\mathbb{E}(\mathbb{E}(X|\mathcal{G})|\mathcal{H})$. Then Y is integrable and $\mathcal{H}$ -measurable. Moreover, for all $A \in \mathcal{H}$,

$$\mathbb{E}(Y1_A) = \mathbb{E}(\mathbb{E}(X|\mathcal{G})1_A) = \mathbb{E}(X1_A)$$

where the second equality holds because $A \in \mathcal{G}$. Hence $Y = \mathbb{E}(X|\mathcal{H})$ almost surely. $\square$

Proposition 1.4 (Taking out what is known). *Let $\mathcal{G}$ be a sub- σ -algebra of $\mathcal{F}$. Let X be an integrable random variable and let Z be a $\mathcal{G}$ -measurable random variable such that ZX is integrable. Then*

$$\mathbb{E}(ZX|\mathcal{G}) = Z\mathbb{E}(X|\mathcal{G}) \quad \text{almost surely.}$$

Proof. ¹ Assume for now that $X \geq 0$. Choose a version $Y \geq 0$ of $\mathbb{E}(X|\mathcal{G})$. Consider first the case when $Z = 1_B$ for some $B \in \mathcal{G}$. Then ZY is $\mathcal{G}$ -measurable and, for all $A \in \mathcal{G}$,

$$\mathbb{E}(ZY1_A) = \mathbb{E}(Y1_{A \cap B}) = \mathbb{E}(X1_{A \cap B}) = \mathbb{E}(ZX1_A).$$

This extends to all simple $\mathcal{G}$ -measurable random variables by linearity. Suppose now that Z is any non-negative $\mathcal{G}$ -measurable random variable. Define, for $n \geq 1$,

$$Z_n = (2^{-n} \lfloor 2^n Z \rfloor) \wedge n.$$

Then Z_n is simple and $\mathcal{G}$ -measurable for all n so, for all $A \in \mathcal{G}$,

$$\mathbb{E}(Z_n Y 1_A) = \mathbb{E}(Z_n X 1_A).$$

Also $0 \leq Z_n \uparrow Z$ as $n \rightarrow \infty$. Since $X \geq 0$ and $Y \geq 0$, we can pass to the limit $n \rightarrow \infty$ using monotone convergence to deduce that, for all $A \in \mathcal{G}$,

$$\mathbb{E}(ZY1_A) = \mathbb{E}(ZX1_A).$$

In particular, by taking $A = \Omega$, we see that if ZX is integrable then so is ZY . For a general $\mathcal{G}$ -measurable Z with ZX integrable, we can apply the preceding to $Z^\pm = (\pm Z) \vee 0$ to see that ZY is integrable and, by subtraction, for all $A \in \mathcal{G}$,

$$\mathbb{E}(ZY1_A) = \mathbb{E}(ZX1_A).$$

Hence $ZY = \mathbb{E}(ZX|\mathcal{G})$ almost surely, as claimed. $\square$

¹Text in blue is not examinable as bookwork in this course.

Proposition 1.5 (Averaging over independent variables). *Let X_1, X_2 be random variables taking values in the measurable spaces $(E_1, \mathcal{E}_1), (E_2, \mathcal{E}_2)$ respectively. Let $\mathcal{G}$ be a sub- σ -algebra of $\mathcal{F}$ and suppose that X_1 is $\mathcal{G}$ -measurable, while X_2 is independent of $\mathcal{G}$. Let F be a non-negative measurable function on $E_1 \times E_2$. Then we can define a non-negative measurable function f on E_1 by $f(x) = \mathbb{E}(F(x, X_2))$ and we have*

$$\mathbb{E}(F(X_1, X_2)|\mathcal{G}) = f(X_1), \quad \text{almost surely.}$$

Proof. Define

$$\mathcal{A} = \{B_1 \times B_2 : B_1 \in \mathcal{E}_1 \text{ and } B_2 \in \mathcal{E}_2\}.$$

Then $\mathcal{A}$ is a π -system (see Probability and Measure) and $\mathcal{A}$ generates the product σ -algebra $\mathcal{E} = \mathcal{E}_1 \otimes \mathcal{E}_2$. It is straightforward to check the desired conclusion when $F = 1_B$ with $B \in \mathcal{A}$. Denote by $\mathcal{D}$ the set of elements of $\mathcal{E}$ such that the conclusion holds for $F = 1_B$. Then $\mathcal{A} \subseteq \mathcal{D}$ and it can be checked that $\mathcal{D}$ is a d -system. Hence $\mathcal{D}$ is the whole of $\mathcal{E}$. The conclusion extends to the case where F is a simple function by linearity, and then to the case where F is any non-negative measurable function by monotone convergence. $\square$

1.2 Definitions

Suppose given a measurable space $(\Omega, \mathcal{F})$. A *filtration* is a family of σ -algebras $(\mathcal{F}_n)_{n \geq 0}$ on Ω such that $\mathcal{F}_n \subseteq \mathcal{F}_{n+1} \subseteq \mathcal{F}$ for all n . A *random process* is a family of random variables $(X_n)_{n \geq 0}$ on $(\Omega, \mathcal{F})$. We say that $(X_n)_{n \geq 0}$ is *adapted* (to $(\mathcal{F}_n)_{n \geq 0}$) if X_n is $\mathcal{F}_n$ -measurable for all n . Define

$$\mathcal{F}_n^X = \sigma(X_k : 0 \leq k \leq n).$$

We call $(\mathcal{F}_n^X)_{n \geq 0}$ the *natural filtration* of $(X_n)_{n \geq 0}$. We think of a filtration as representing the emergence of information over time. In particular $\mathcal{F}_n^X$ contains all the events determined by the process $(X_n)_{n \geq 0}$ up to time n .

Suppose now that $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space, which is equipped with a filtration $(\mathcal{F}_n)_{n \geq 0}$. We say that a random process $(X_n)_{n \geq 0}$ is a *martingale* if, for all n ,

- (a) X_n is $\mathcal{F}_n$ -measurable,
- (b) $\mathbb{E}(|X_n|) < \infty$,
- (c) $\mathbb{E}(X_{n+1}|\mathcal{F}_n) = X_n$ almost surely.

When (b) holds, we say that $(X_n)_{n \geq 0}$ is *integrable*. Condition (c) is known as the *martingale property*. Thus, in short, a martingale is an adapted integrable process which satisfies the martingale property. The martingale property can be expressed in terms of (unconditional) expectation as the requirement that

$$\mathbb{E}((X_{n+1} - X_n)1_A) = 0 \quad \text{for all } A \in \mathcal{F}_n.$$

Indeed, it suffices that this relation holds for all A in some π -system generating $\mathcal{F}_n$.

If conditions (a) and (b) hold and, for all n ,

$$\mathbb{E}(X_{n+1}|\mathcal{F}_n) \leq X_n \quad \text{almost surely}$$

then we say that $(X_n)_{n \geq 0}$ is a *supermartingale*. Similarly, if conditions (a) and (b) hold and, for all n ,

$$\mathbb{E}(X_{n+1}|\mathcal{F}_n) \geq X_n \quad \text{almost surely}$$

then we say that $(X_n)_{n \geq 0}$ is a *submartingale*.

Any martingale is also a martingale in its natural filtration. When we refer to a martingale without specifying a filtration, the natural filtration is understood by default.

All the above notions have continuous-time analogues, which are mostly obtained simply by writing t for n , with the convention that $t \in [0, \infty)$ where we had $n \in \mathbb{Z}^+$. Given a filtration $(\mathcal{F}_t)_{t \geq 0}$, we say that $(X_t)_{t \geq 0}$ is a *martingale* if it is adapted, integrable and satisfies, for all $s, t \geq 0$ with $s \leq t$,

$$\mathbb{E}(X_t|\mathcal{F}_s) = X_s \quad \text{almost surely.}$$

The supermartingale and submartingale properties are modified similarly. We say that $(X_t)_{t \geq 0}$ is *continuous* if the map

$$t \mapsto X_t(\omega) : [0, \infty) \rightarrow \mathbb{R}$$

is continuous for all ω . These notions will be relevant when we come to look at Brownian motion, which is itself a continuous martingale.

1.3 Examples

Let $(X_n)_{n \geq 1}$ be a sequence of independent identically distributed random variables. Set $\mathcal{F}_0 = \{\emptyset, \Omega\}$ and for $n \geq 1$ set $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$. Set $S_0 = 0$ and $Z_0 = 1$ and define for $n \geq 1$

$$S_n = \sum_{k=1}^n X_k, \quad Z_n = \prod_{k=1}^n X_k.$$

In the case where X_1 is integrable with $\mathbb{E}(X_1) = 0$, the process $(S_n)_{n \geq 0}$ is a martingale. On the other hand, in the case where $X_1 \geq 0$ with $\mathbb{E}(X_1) = 1$, the process $(Z_n)_{n \geq 0}$ is a martingale. We call $(S_n)_{n \geq 0}$ an *additive martingale* and $(Z_n)_{n \geq 0}$ a *multiplicative martingale*.

Further examples arise in connection with Markov chains. Let $(X_n)_{n \geq 0}$ be a Markov chain with countable state-space S and transition matrix P . Set $\mathcal{F}_n = \sigma(X_0, \dots, X_n)$. Given a bounded or non-negative function f on S , we set

$$Pf(x) = \sum_{y \in S} p_{xy} f(y).$$

Then

$$\mathbb{E}(f(X_{n+1})|\mathcal{F}_n) = Pf(X_n) \quad \text{almost surely.}$$

We say that f is *subharmonic* if $f(x) \leq Pf(x)$ for all x .

Suppose that f is subharmonic and define $M_n = f(X_n)$. Then

$$\mathbb{E}(M_{n+1}|\mathcal{F}_n) = Pf(X_n) \geq f(X_n) = M_n \quad \text{almost surely}$$

so $(M_n)_{n \geq 0}$ has the submartingale property.

1.4 Optional stopping

We say that a random time $T : \Omega \rightarrow \{0, 1, \dots\} \cup \{\infty\}$ is a *stopping time* if, for all n ,

$$\{T \leq n\} \in \mathcal{F}_n.$$

Theorem 1.6 (Optional stopping). *Let $(M_n)_{n \geq 0}$ be a martingale and let T be a bounded stopping time. Then $\mathbb{E}(M_T) = \mathbb{E}(M_0)$.*

Proof. We can choose n so that $T \leq n$. Then

$$M_T = (M_T - M_{T-1}) + \dots + (M_1 - M_0) + M_0 = M_0 + \sum_{k=1}^n (M_k - M_{k-1}) 1_{\{k \leq T\}}.$$

Since T is a stopping time, $\{k \leq T\} = \{T \leq k-1\}^c \in \mathcal{F}_{k-1}$ for all k . Then, since $(M_n)_{n \geq 0}$ is a martingale,

$$\mathbb{E}(M_k 1_{\{k \leq T\}}) = \mathbb{E}(M_{k-1} 1_{\{k \leq T\}}).$$

Hence

$$\mathbb{E}(M_T) = \mathbb{E}(M_0) + \sum_{k=1}^n \mathbb{E}((M_k - M_{k-1}) 1_{\{k \leq T\}}) = \mathbb{E}(M_0).$$

□

This is also called *Doob's optional sampling theorem*. We have stated the basic form of the result. We now show an extension to unbounded stopping times under additional conditions.

Theorem 1.7. *Let $(M_n)_{n \geq 0}$ be a martingale and let T be an almost surely finite stopping time. Suppose that, for some constant $C < \infty$, one of the following two conditions holds:*

- (a) $|M_n| \leq C$ for all $n \leq T$,
- (b) $\mathbb{E}(T) < \infty$ and $|M_n - M_{n-1}| \leq C$ for all $n \in \{1, \dots, T\}$.

Then $\mathbb{E}(M_T) = \mathbb{E}(M_0)$.

Proof. Since T is a stopping time, $T \wedge n$ is a bounded stopping time for all n so, by the optional stopping theorem,

$$\mathbb{E}(M_{T \wedge n}) = \mathbb{E}(M_0).$$

Hence, it will suffice to show that $\mathbb{E}(M_{T \wedge n}) \rightarrow \mathbb{E}(M_T)$ as $n \rightarrow \infty$. Since T is almost surely finite, we have $M_{T \wedge n} \rightarrow M_T$ almost surely as $n \rightarrow \infty$. Condition (a) implies that $|M_{T \wedge n}| \leq C$ for all n , so the desired limit holds by bounded convergence. On the other hand, condition (b) implies that $|M_{T \wedge n}| \leq |M_0| + CT$ for all n , so the desired limit holds by dominated convergence. $\square$

It is instructive to consider two simple examples where the optional stopping theorem does not apply. Consider first the additive martingale $S_n = \sum_{k=1}^n X_k$ associated to a sequence of independent identically distributed random variables $(X_n)_{n \geq 1}$, with

$$\mathbb{P}(X_1 = 1) = \mathbb{P}(X_1 = -1) = 1/2.$$

Set

$$T = \min\{n \geq 0 : S_n = 1\}.$$

Then $(S_n)_{n \geq 0}$ is a simple symmetric random walk, which is recurrent, so $T < \infty$ almost surely. But

$$\mathbb{E}(S_T) = 1 \neq 0 = S_0$$

so the conclusion of the optional stopping theorem is false.

Consider now the multiplicative martingale $Z_n = \prod_{k=1}^n X_k$ associated to a sequence of independent identically distributed random variables $(X_n)_{n \geq 1}$, with

$$\mathbb{P}(X_1 = 0) = \mathbb{P}(X_1 = 2) = 1/2.$$

Set

$$T = \min\{n \geq 0 : Z_n = 0\}.$$

Then $\mathbb{E}(T) = 2$. But

$$\mathbb{E}(Z_T) = 0 \neq 1 = Z_0$$

so the conclusion of the optional stopping theorem is again false. In this and the preceding example, one of the conditions in hypothesis (b) above holds, but not the other.

Here is a further variant of the optional stopping theorem.

Theorem 1.8. *Let $(M_n)_{n \geq 0}$ be a martingale and let T be a stopping time. Then the stopped process $(M_{T \wedge n})_{n \geq 0}$ is also a martingale.*

Proof. Note that

$$M_{T \wedge n} = M_0 + \sum_{k=1}^n (M_k - M_{k-1}) 1_{\{T \geq k\}}$$

and, since T is a stopping time, $\{T \geq k\} = \{T \leq k-1\}^c \in \mathcal{F}_{k-1}$. Since $(M_n)_{n \geq 0}$ is adapted and integrable, this makes it clear that $M_{T \wedge n}$ is $\mathcal{F}_n$ -measurable and integrable for all n . Now

$$M_{T \wedge (n+1)} - M_{T \wedge n} = (M_{n+1} - M_n)1_{\{T \geq n+1\}}$$

so, by taking out what is known,

$$\mathbb{E}(M_{T \wedge (n+1)} - M_{T \wedge n} | \mathcal{F}_n) = 1_{\{T \geq n+1\}} \mathbb{E}(M_{n+1} - M_n | \mathcal{F}_n) = 0.$$

Hence $(M_{T \wedge n})_{n \geq 0}$ has the martingale property. $\square$

We say that a random process $(H_n)_{n \geq 1}$ is *previsible* if H_n is $\mathcal{F}_{n-1}$ -measurable for all n .

Theorem 1.9 (Martingale transform). *Let $(M_n)_{n \geq 0}$ be a martingale and let $(H_n)_{n \geq 1}$ be a bounded previsible process. Define a new random process $(Y_n)_{n \geq 0}$ by*

$$Y_0 = 0, \quad Y_n = \sum_{k=1}^n H_k(M_k - M_{k-1}), \quad n \geq 1.$$

Then $(Y_n)_{n \geq 0}$ is also a martingale.

Proof. Since $(M_n)_{n \geq 0}$ is adapted and integrable, it is clear from the definition that $(Y_n)_{n \geq 0}$ is also adapted and integrable. Now

$$Y_{n+1} - Y_n = H_{n+1}(M_{n+1} - M_n)$$

and H_{n+1} is $\mathcal{F}_n$ -measurable so, by taking out what is known,

$$\mathbb{E}(Y_{n+1} - Y_n | \mathcal{F}_n) = H_{n+1} \mathbb{E}(M_{n+1} - M_n | \mathcal{F}_n) = 0.$$

Hence $(Y_n)_{n \geq 0}$ also has the martingale property. $\square$

The process $(Y_n)_{n \geq 0}$ is called the *martingale transform* of $(M_n)_{n \geq 0}$ by $(H_n)_{n \geq 1}$. Note that the proofs of Theorems 1.8 and 1.9 use essentially the same argument.

The optional stopping theorem and its variants may be interpreted in terms of an investor who is able to buy and sell a risky asset. We model the price of the asset by the martingale $(M_n)_{n \geq 0}$. Suppose that the investor holds one unit of the asset at time 0 and wishes to sell it by time n . The investor is free to choose when to sell, based on the information available at the present time, that is, at any stopping time T of the filtration $(\mathcal{F}_n)_{n \geq 0}$. Then the optional stopping theorem shows that the expected return $\mathbb{E}(M_T)$ is the same no matter what stopping time is chosen. More generally, we may suppose that the investor is free to buy or sell any number of units in each time period, again based on information available at the start of the period. The amount H_k held between times $k-1$ and k will then be $\mathcal{F}_{k-1}$ -measurable, so defining a previsible process, and the total gain by time n is then given by Y_n . Theorem 1.9 shows that no bounded previsible strategy results in an expected gain or loss.

2 Pricing contingent claims

2.1 Multi-period asset price model

Consider a discrete-time random process $(\bar{S}_n)_{0 \leq n \leq T}$ in $\mathbb{R}^{d+1}$, defined on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and adapted to a filtration $(\mathcal{F}_n)_{0 \leq n \leq T}$. Write

$$\bar{S}_n = (S_n^0, S_n^1, \dots, S_n^d) = (S_n^0, S_n).$$

We assume that $(S_n^0)_{0 \leq n \leq T}$ is a *numéraire*, that is to say, $S_n^0 > 0$ for all n . Then we call $(\bar{S}_n)_{0 \leq n \leq T}$ an *asset price model*.

We use this set-up to model the evolution over time of $d + 1$ assets, interpreting S_n^i as the price at time n of the i th asset. There are T time periods: 0 to 1, 1 to 2 and finally $T - 1$ to T . Thus $S_n^i - S_{n-1}^i$ is the change in price of the i th asset over the n th time period.

We will be interested not so much in the absolute prices S_n^i , but in the *discounted prices* X_n^i , given by

$$X_n^i = S_n^i / S_n^0$$

and we write

$$\bar{X}_n = (X_n^0, X_n^1, \dots, X_n^d) = (1, X_n).$$

Often, $(S_n^0)_{0 \leq n \leq T}$ will be interpreted as a bond or bank account. In this case, we will write

$$S_n^0 = (1 + r_n) S_{n-1}^0$$

and call r_n the *interest rate*.

We often take $(\mathcal{F}_n)_{0 \leq n \leq T}$ to be the filtration generated by $(\bar{S}_n)_{0 \leq n \leq T}$ and take $\mathcal{F} = \mathcal{F}_T$. In that case, if say $(S_n^0)_{0 \leq n \leq T}$ is a given deterministic process, and the prices of the other *risky assets* take only a countable set of values, then we will be able to specify $\mathbb{P}$ completely by specifying the probabilities $\mathbb{P}(S_0 = s_0, S_1 = s_1, \dots, S_T = s_T)$ for all possible sequences $(s_n)_{0 \leq n \leq T}$ in $\mathbb{R}^d$.

Let $(\bar{\theta}_n)_{1 \leq n \leq T}$ be a random process in $\mathbb{R}^{d+1}$. Write

$$\bar{\theta}_n = (\theta_n^0, \theta_n^1, \dots, \theta_n^d) = (\theta_n^0, \theta_n).$$

We use such a process to model a portfolio held by an investor, where θ_n^i is the number of units of asset i held in the n th time period, allowing the possibility that this number may take any real value. We say that a portfolio is *self-financing* if

$$\bar{\theta}_n \cdot \bar{S}_n = \bar{\theta}_{n+1} \cdot \bar{S}_n \quad \text{for } n = 1, \dots, T - 1.$$

This expresses that no funds are added or withdrawn from the portfolio at time n , it is simply rebalanced between the $d + 1$ assets. We associate to a self-financing portfolio its *value process* $(V_n)_{0 \leq n \leq T}$ given by

$$V_0 = \bar{\theta}_1 \cdot \bar{X}_0, \quad V_n = \bar{\theta}_n \cdot \bar{X}_n \quad \text{for } n = 1, \dots, T.$$

Thus V_n is the total (discounted) value of the portfolio at time n .

We say that a portfolio is *previsible* if $\bar{\theta}_n$ is $\mathcal{F}_{n-1}$ -measurable for all n . This is a natural condition, since it expresses that $\bar{\theta}_n$ depends only on what is known at the start of the n th period, that is, at time $n - 1$.

Proposition 2.1. *Let $(\theta_n)_{1 \leq n \leq T}$ be a previsible process in $\mathbb{R}^d$ and let $V_0 \in \mathbb{R}$. Then there is a unique previsible process $(\theta_n^0)_{1 \leq n \leq T}$ in $\mathbb{R}$ such that $(\bar{\theta}_n)_{1 \leq n \leq T}$ is a self-financing portfolio with initial value V_0 . Moreover, the value process of this portfolio is given by*

$$V_T = V_0 + \sum_{n=1}^T \theta_n \cdot (X_n - X_{n-1}).$$

Proof. The equations $\bar{\theta}_1 \cdot \bar{X}_0 = V_0$ and $\bar{\theta}_n \cdot \bar{S}_n = \bar{\theta}_{n+1} \cdot \bar{S}_n$ for $n = 1, \dots, T - 1$, which express that the portfolio $(\bar{\theta}_n)_{1 \leq n \leq T}$ has initial value V_0 and is self-financing, may be written in the form

$$\theta_1^0 + \theta_1 \cdot X_0 = V_0, \quad \theta_n^0 S_n^0 + \theta_n \cdot S_n = \theta_{n+1}^0 S_n^0 + \theta_{n+1} \cdot S_n, \quad \text{for } n = 1, \dots, T - 1$$

and then solved uniquely to obtain a previsible process $(\theta_n^0)_{1 \leq n \leq T}$. Then, since $X_n^0 = 1$ for all n , the value of the resulting portfolio changes by $\theta_n \cdot (X_n - X_{n-1})$ over the n th time period, giving the claimed formula for V_T by induction. $\square$

2.2 Contingent claims

By a *contingent claim of maturity T* we mean any non-negative $\mathcal{F}_T$ -measurable random variable. We think of the contingent claim C as a contract which pays the investor the amount C at time T . The fact that the contract is always exercised at the given time T can be emphasised by referring to it as a *European contingent claim*.

Here are some simple examples in the case $d = 1$ of one risky asset:

- (a) $(S_T - K)^+$ is the *call* with *strike price* K , which can be understood as the right but not the obligation to buy one unit of the asset at time T for price K ,
- (b) $(S_T - K)^-$ is the *put* with *strike price* K , which can be understood as the right but not the obligation to sell one unit of the asset at time T for price K .

Calls and puts are also called *options*, emphasising the option to buy or sell. The following are examples of *exotic options*, depending on the entire path $(S_n)_{0 \leq n \leq T}$. They are both examples of *barrier options* which are *knocked out* or *knocked in* when the price hits or passes a given barrier B .

- (c) The *up-and-out call* is a call which is knocked out at level B , given by

$$C = \begin{cases} (S_T - K)^+, & \text{if } \max_{0 \leq n \leq T} S_n < B, \\ 0, & \text{otherwise.} \end{cases}$$

(d) The *down-and-in put* is a put which is knocked in at level B , given by

$$C = \begin{cases} (S_T - K)^-, & \text{if } \min_{0 \leq n \leq T} S_n \leq B, \\ 0, & \text{otherwise.} \end{cases}$$

It is an important problem to find a way to value contingent claims. A natural way to do this, for claims based on an asset price process with a known distribution is to take as value the expected payout relative to the numéraire, given by $\mathbb{E}(C/S_T^0)$. In practice, the distribution of an asset price process may however not be known, or different investors may have different beliefs about the distribution. We will explore instead the notion of no-arbitrage pricing, which is more robust to our assumptions about the asset price model.

2.3 Equivalent probability measures

We say that a probability measure $\tilde{\mathbb{P}}$ on $(\Omega, \mathcal{F})$ is *equivalent* to $\mathbb{P}$, and we write $\tilde{\mathbb{P}} \sim \mathbb{P}$, if there exists a random variable ρ with values in $(0, \infty)$ such that

$$\tilde{\mathbb{P}}(A) = \mathbb{E}(\rho 1_A) \quad \text{for all } A \in \mathcal{F}.$$

Note that $\tilde{\mathbb{P}}(A) = 0$ whenever $\mathbb{P}(A) = 0$. We leave as an exercise to check that any two random variables ρ in $(0, \infty)$ with this property are equal $\mathbb{P}$ -almost surely. We call ρ a *density* for $\tilde{\mathbb{P}}$ with respect to $\mathbb{P}$ and write

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = \rho \quad \text{almost surely.}$$

By a standard argument of measure theory, the expectation $\tilde{\mathbb{E}}$ associated with $\tilde{\mathbb{P}}$ is given by

$$\tilde{\mathbb{E}}(X) = \mathbb{E}(\rho X)$$

for all non-negative random variables X . We leave as an exercise to check that the relation $\tilde{\mathbb{P}} \sim \mathbb{P}$ is symmetric and transitive, with

$$\frac{d\mathbb{P}}{d\tilde{\mathbb{P}}} = 1/\rho \quad \text{almost surely.}$$

2.4 Arbitrage

An arbitrage is a risk-free profit available in the market. The principle of no arbitrage states that prices in the market do not allow this. In mathematical terms, for an asset price model $(\bar{S}_n)_{0 \leq n \leq T}$, by an *arbitrage* we mean a previsible self-financing portfolio $(\bar{\theta}_n)_{1 \leq n \leq T}$ with initial value $V_0 = 0$ such that

$$V_T \geq 0 \quad \text{almost surely}$$

and

$$V_T > 0 \quad \text{with positive probability.}$$

An arbitrage for $(\bar{S}_n)_{0 \leq n \leq T}$ thus delivers a positive return with no downside risk. The value process $(V_n)_{0 \leq n \leq T}$ depends on $(\bar{S}_n)_{0 \leq n \leq T}$ only through $(X_n)_{0 \leq n \leq T}$, and it depends on $(\bar{\theta}_n)_{1 \leq n \leq T}$ only through V_0 and $(\theta_n)_{1 \leq n \leq T}$. We will therefore allow ourselves to refer to $(\theta_n)_{1 \leq n \leq T}$ as an arbitrage for $(X_n)_{0 \leq n \leq T}$ when V_T satisfies the given conditions.

Note that the notion of arbitrage depends only on the equivalence class of the probability measure $\mathbb{P}$.

If there is no arbitrage for $(\bar{S}_n)_{0 \leq n \leq T}$, then we say that $(\bar{S}_n)_{0 \leq n \leq T}$ (or $(X_n)_{0 \leq n \leq T}$) is *arbitrage free*. This is considered a reasonable modelling assumption, on the grounds that the other investors in the market would not collectively give away such a return, even if they held different views about the evolution of the asset prices.

Note that, if $(X_n)_{0 \leq n \leq T}$ is a martingale and $(\theta_n)_{1 \leq n \leq T}$ is a bounded previsible process, then the martingale transform

$$V_T = \sum_{n=1}^T \theta_n \cdot (X_n - X_{n-1})$$

is integrable and $\mathbb{E}(V_T) = 0$. Hence, if $V_T \geq 0$ almost surely, then $V_T = 0$ almost surely, so $(\theta_n)_{1 \leq n \leq T}$ is not an arbitrage. This shows that if the discounted price process $(X_n)_{0 \leq n \leq T}$ is a martingale then there is no bounded arbitrage.

In fact, by the following more elaborate argument, even unbounded processes cannot give an arbitrage in this case.

Proposition 2.2. *Let $(X_n)_{0 \leq n \leq T}$ be a martingale in $\mathbb{R}^d$. Then $(X_n)_{0 \leq n \leq T}$ is arbitrage free.*

Proof. Let $(\theta_n)_{1 \leq n \leq T}$ be a previsible process in $\mathbb{R}^d$ and set

$$V_n = \sum_{k=1}^n \theta_k \cdot (X_k - X_{k-1}).$$

Suppose that $V_T \geq 0$ almost surely. We have $\mathbb{E}(V_T | \mathcal{F}_T) = V_T$ almost surely. Suppose in a backwards induction for $n \leq T$ that $V_n = \mathbb{E}(V_T | \mathcal{F}_n)$ almost surely. We have

$$V_n = V_{n-1} + \theta_n \cdot (X_n - X_{n-1}).$$

Fix $R < \infty$ and set $A = \{|\theta_n| \leq R \text{ and } |V_{n-1}| \leq R\}$. Then $A \in \mathcal{F}_{n-1}$ and

$$1_A V_n = 1_A V_{n-1} + 1_A \theta_n \cdot (X_n - X_{n-1}).$$

Since $V_n = \mathbb{E}(V_T | \mathcal{F}_n) \geq 0$ almost surely and $1_A |V_{n-1}| \leq R$ and $1_A |\theta_n| \leq R$, we can take the conditional expectation on $\mathcal{F}_{n-1}$, taking out what is known, to obtain

$$1_A \mathbb{E}(V_n | \mathcal{F}_{n-1}) = 1_A V_{n-1} + 1_A \theta_n \mathbb{E}(X_n - X_{n-1} | \mathcal{F}_{n-1}) = 1_A V_{n-1} \quad \text{almost surely.}$$

But R was arbitrary and $|\theta_n|$ and $|V_{n-1}|$ are almost surely finite, so we must have

$$V_{n-1} = \mathbb{E}(V_n | \mathcal{F}_{n-1}) = \mathbb{E}(V_T | \mathcal{F}_{n-1}) \quad \text{almost surely}$$

and the induction proceeds. In particular, we see that $\mathbb{E}(V_T | \mathcal{F}_0) = V_0 = 0$ almost surely, so $\mathbb{E}(V_T) = 0$ and so $V_T = 0$ almost surely. Hence there is no arbitrage. $\square$

2.5 Fundamental theorem of asset pricing

The following theorem shows a useful property, namely the existence of an equivalent martingale measure, which characterizes asset price models constrained by the assumption of no arbitrage. Let $(\bar{S}_n)_{0 \leq n \leq T}$ be an asset price model in $\mathbb{R}^{d+1}$. Recall that $X_n = (X_n^1, \dots, X_n^d)$ where $X_n^i = S_n^i/S_n^0$ for all i . We say that a probability measure $\tilde{\mathbb{P}}$ on $\mathcal{F}$ is an *equivalent martingale measure* if $\tilde{\mathbb{P}} \sim \mathbb{P}$ and $(X_n)_{0 \leq n \leq T}$ is a martingale under $\tilde{\mathbb{P}}$. The term *risk-neutral measure* is also used in this case.

Theorem 2.3. *The following are equivalent:*

- (a) $(\bar{S}_n)_{0 \leq n \leq T}$ has no arbitrage,
- (b) $(\bar{S}_n)_{0 \leq n \leq T}$ has an equivalent martingale measure.

We will prove this for the single-period case with non-random initial conditions. A proof of the general case is given in the Appendix.

Proof of the fundamental theorem in the case $T = 1$ and $\mathcal{F}_0 = \{\emptyset, \Omega\}$. Define a random variable Y in $\mathbb{R}^d$ by setting $Y = X_1 - X_0$, where $X_n = S_n/S_n^0$ for $n = 0, 1$. Given a previsible self-financing portfolio $\bar{\theta}$ with initial value 0, set $\theta = \theta_1$ and note that $\theta \in \mathbb{R}^d$ and $V_1 = \theta \cdot Y$. Hence we can reformulate the theorem in this case as the equivalence of the following statements for a random variable Y in $\mathbb{R}^d$.

- (a) there exists no $\theta \in \mathbb{R}^d$ such that $\theta \cdot Y \geq 0$ almost surely and $\theta \cdot Y > 0$ with positive probability,
- (b) there is an equivalent probability measure $\tilde{\mathbb{P}}$ for which Y is integrable with $\tilde{\mathbb{E}}(Y) = 0$.

Suppose (b) holds. Then $\tilde{\mathbb{E}}(\theta \cdot Y) = 0$ for all $\theta \in \mathbb{R}^d$, so if $\theta \cdot Y \geq 0$ almost surely then $\theta \cdot Y = 0$ almost surely. Hence (a) holds.

Suppose on the other hand that (a) holds. It will suffice to consider the case where all exponential moments of Y are finite. For, if this does not already hold, we can replace $\mathbb{P}$ by the equivalent probability measure $\tilde{\mathbb{P}}$ given by $d\tilde{\mathbb{P}}/d\mathbb{P} \propto e^{-|Y|^2}$, for which the exponential moments of Y are all finite.

Define $\phi : \mathbb{R}^d \rightarrow (0, \infty)$ by $\phi(\theta) = \mathbb{E}(e^{\theta \cdot Y})$. Then ϕ is differentiable and

$$\phi'(\theta) = \mathbb{E}(Y e^{\theta \cdot Y}).$$

We will show that ϕ achieves a minimum value at some $\theta^* \in \mathbb{R}^d$. Given such a θ^* , we can define an equivalent probability measure $\tilde{\mathbb{P}}$ by $d\tilde{\mathbb{P}}/d\mathbb{P} = e^{\theta^* \cdot Y}/Z$, where $Z = \phi(\theta^*)$. Then

$$\tilde{\mathbb{E}}(|Y|) = \mathbb{E}(|Y| e^{\theta^* \cdot Y})/Z < \infty, \quad \tilde{\mathbb{E}}(Y) = \mathbb{E}(Y e^{\theta^* \cdot Y})/Z = \phi'(\theta^*)/Z = 0.$$

Hence (b) holds.

Write $\mathbb{R}^d$ as an orthogonal direct sum $E_0 \oplus E_1$, where

$$E_0 = \{\theta \in \mathbb{R}^d : \theta \cdot Y = 0 \text{ almost surely}\}.$$

Then $\phi(\theta_0 + \theta_1) = \phi(\theta_1)$ for all $\theta_0 \in E_0$ and $\theta_1 \in E_1$. Hence it will suffice to show that ϕ achieves a minimum on E_1 . Since $\phi(0) = 1$, it will then suffice to show that $\phi(\theta) \geq 1$ for all $\theta \in E_1$ with $|\theta|$ sufficiently large.

Set $\psi(t) = 0 \vee t \wedge 1$ and define $f(\theta) = \mathbb{E}(\psi(\theta.Y))$. Then ψ is Lipschitz on $\mathbb{R}$ of constant 1 and f is continuous on $\mathbb{R}^d$ by bounded convergence. Define $S = \{\theta \in E_1 : |\theta| = 1\}$. For $\theta \in S$, we have $\mathbb{P}(\theta.Y \neq 0) > 0$ so $\mathbb{P}(\theta.Y > 0) > 0$ and so $f(\theta) > 0$. Set

$$\varepsilon = \frac{1}{2} \inf\{f(\theta) : \theta \in S\}.$$

Then $\varepsilon > 0$ because S is compact. Then, for all $\theta \in S$,

$$\mathbb{P}(\theta.Y \geq \varepsilon) = \mathbb{P}(\theta.Y - \varepsilon \geq 0) \geq \mathbb{E}(\psi(\theta.Y - \varepsilon)) \geq \mathbb{E}(\psi(\theta.Y)) - \varepsilon = f(\theta) - \varepsilon \geq \varepsilon$$

Hence we obtain $\phi(t\theta) \geq \varepsilon e^{t\varepsilon} \geq 1$ for all $t \geq (1/\varepsilon) \log(1/\varepsilon)$, as required. $\square$

2.6 Pricing attainable claims

Let C be a time- T contingent claim. Write $D = C/S_T^0$ for its discounted value. We say that C is *attainable* or *replicable* if there is a previsible self-financing portfolio $(\bar{\theta}_n)_{1 \leq n \leq T}$ such that $C = \bar{\theta}_T \cdot \bar{S}_T$. Equivalently, C (or we sometimes say D) is attainable if there exists an $\mathcal{F}_0$ -measurable random variable V_0 and a previsible process $(\theta_n)_{1 \leq n \leq T}$ in $\mathbb{R}^d$ such that

$$D = V_0 + \sum_{n=1}^T \theta_n \cdot (X_n - X_{n-1}).$$

Then V_0 is called the *fair price* for C and $(\bar{\theta}_n)_{1 \leq n \leq T}$ is called a *replicating portfolio*. We also call $(\bar{\theta}_n)_{1 \leq n \leq T}$ a *hedging portfolio*: it allows the seller of the contingent claim C to deliver it at time T by investing V_0 suitably in the available assets at time 0, and shifting the portfolio among the assets according to $(\bar{\theta}_n)_{1 \leq n \leq T}$ until time T . Note that anyone who sells the claim C at time 0 for a different price to V_0 allows other investors to make risk-free profits by replicating C or $-C$ in the market.

Proposition 2.4. *Assume that $\mathcal{F}_0 = \{\emptyset, \Omega\}$ and $\mathcal{F}_T = \sigma(\bar{S}_1, \dots, \bar{S}_T)$. Let C be an attainable time- T contingent claim and suppose that $\tilde{\mathbb{P}}$ is an equivalent martingale measure. Then the fair price V_0 for C is given by*

$$V_0 = \tilde{\mathbb{E}}(C)/S_T^0.$$

Proof. Since C is attainable, there is a previsible process $(\theta_n)_{1 \leq n \leq T}$ such that

$$D = C/S_T^0 = V_0 + \sum_{n=1}^T \theta_n \cdot (X_n - X_{n-1}).$$

Then, since $D \geq 0$, the argument used in the proof of Proposition 2.2 shows that $\tilde{\mathbb{E}}(D) = V_0$. $\square$

2.7 Pricing non-attainable claims in a single-period model

Consider a single-period asset price model $(\bar{S}_0, \bar{S}_1)$ without arbitrage. Let us consider possible time-0 prices C_0 for a time-1 contingent claim C_1 . Write $X_n = S_n/S_n^0$ as usual and set $C = (C_0, C_1)$. We can think of $(\bar{S}, C)$ itself as a single-period model. Let us say that the price C_0 *presents no arbitrage for C_1* if there is no $\theta \in \mathbb{R}^d$ such that, either

$$C_0 \leq C_1/S_1^0 + \theta \cdot (X_1 - X_0) \quad \text{a.s.} \quad \text{and} \quad C_0 < C_1/S_1^0 + \theta \cdot (X_1 - X_0) \quad \text{w.p.p.}$$

or

$$C_0 \geq C_1/S_1^0 - \theta \cdot (X_1 - X_0) \quad \text{a.s.} \quad \text{and} \quad C_0 > C_1/S_1^0 + \theta \cdot (X_1 - X_0) \quad \text{w.p.p.}$$

It is straightforward to check that this is a reformulation of the property that $(\bar{S}, C)$ has no arbitrage. Hence, by the Fundamental Theorem, there exists an equivalent martingale measure $\tilde{\mathbb{P}}$ for $(\bar{S}, C)$. Note that such measures are precisely the equivalent martingale measures $\tilde{\mathbb{P}}$ for $\bar{S}$ such that $\tilde{\mathbb{E}}(C_1)/S_1^0 = C_0$. Note also that, since any convex combination of equivalent martingale measures for $\bar{S}$ is itself one, the following set is an interval

$$I(C_1) = \{\tilde{\mathbb{E}}(C_1)/S_1^0 : \tilde{\mathbb{P}} \text{ an equivalent martingale measure for } \bar{S}\}.$$

We have shown the following result.

Proposition 2.5. *Under the above conditions, the price C_0 presents no arbitrage for C_1 if and only if $C_0 \in I(C_1)$.*

2.8 Completeness

We say that an asset price model $(\bar{S}_n)_{0 \leq n \leq T}$ is *complete* if all time- T contingent claims are attainable. We will identify a class of complete models in the next section.

Proposition 2.6. *Assume that $\mathcal{F}_0 = \{\emptyset, \Omega\}$ and $\mathcal{F}_T = \sigma(\bar{S}_1, \dots, \bar{S}_T)$. Assume further that $(\bar{S}_n)_{0 \leq n \leq T}$ is complete and that the numéraire is non-random. Then there is at most one equivalent martingale measure on $\mathcal{F}_T$.*

Proof. Let $A \in \mathcal{F}_T$. Then 1_A is a contingent claim, so 1_A is attainable, with fair value $V_0(A) = v(A)/S_T^0$ say. Then, by Proposition 2.4, $\tilde{\mathbb{P}}(A) = v(A)$ for any equivalent martingale measure. $\square$

2.9 Binomial model

Fix parameters $r, a, b \in (-1, \infty)$, $p \in (0, 1)$ and $S_0 \in (0, \infty)$, with $a < b$. We say that $(S_n^0, S_n)_{0 \leq n \leq T}$ is a *binomial model* with interest rate r and parameters $a < b$ and p if

$$S_n^0 = (1 + r)^n, \quad S_n = S_0 \prod_{k=1}^n (1 + R_k)$$

where $R_1, \dots, R_T$ are independent identically distributed random variables with

$$\mathbb{P}(R_1 = a) = 1 - p, \quad \mathbb{P}(R_1 = b) = p.$$

This is also called the *Cox–Ross–Rubinstein model*. Assume that $\mathcal{F} = \mathcal{F}_T = \sigma(R_1, \dots, R_T)$.

Proposition 2.7. *A binomial model with interest rate r and parameters $a < b$ and p has an arbitrage unless $r \in (a, b)$.*

Proof. Consider the self-financing portfolio $(\bar{\theta}_n)_{1 \leq n \leq T}$ with $V_0 = 0$, $\theta_1 = 1$ and $\theta_n = 0$ for $n \geq 2$. Then

$$V_T = X_1 - X_0 = \frac{S_0(1 + R_1)}{1 + r} - S_0 = S_0 \frac{R_1 - r}{1 + r}.$$

If $r \leq a$, then $V_T \geq 0$ and $V_T = S_0(b - r)/(1 + r) > 0$ with probability p , so $(\bar{\theta}_n)_{1 \leq n \leq T}$ is an arbitrage. On the other hand, if $r \geq b$, then $V_T \leq 0$ and $V_T = -S_0(r - a)/(1 + r) < 0$ with probability $1 - p$, so $(-\bar{\theta}_n)_{1 \leq n \leq T}$ is an arbitrage. $\square$

Proposition 2.8. *Let $(S_n^0, S_n)_{0 \leq n \leq T}$ be a binomial model with interest rate r and parameters $a < b$ and p . Suppose that $r \in (a, b)$ and define $p^* \in (0, 1)$ by*

$$p^* = \frac{r - a}{b - a}.$$

Consider the equivalent probability measure $\mathbb{P}^$ on $\mathcal{F}$ given by*

$$\frac{d\mathbb{P}^*}{d\mathbb{P}} = \left(\frac{p^*}{p}\right)^{U_T} \left(\frac{1 - p^*}{1 - p}\right)^{D_T}$$

where $U_T = (T + S_T)/2$ and $D_T = (T - S_T)/2$. Then, under $\mathbb{P}^$, the random variables $R_1, \dots, R_T$ are independent and identically distributed, with*

$$\mathbb{P}(R_1 = a) = 1 - p^*, \quad \mathbb{P}(R_1 = b) = p^*.$$

Moreover, the discounted price process $(X_n)_{0 \leq n \leq T}$ is a martingale under $\mathbb{P}^$.*

Proof. Note that $r = (1 - p^*)a + p^*b$. We have

$$X_n = \frac{S_n}{S_n^0} = S_0 \prod_{k=1}^n \left(\frac{1 + R_k}{1 + r} \right)$$

and

$$\mathbb{E}^* \left(\frac{1 + R_1}{1 + r} \right) = \frac{(1 - p^*)(1 + a) + p^*(1 + b)}{1 + r} = \frac{1 + (1 - p^*)a + p^*b}{1 + r} = 1.$$

Since the random variables $R_1, \dots, R_T$ are independent and identically distributed under $\mathbb{P}^*$, this implies that $(X_n)_{0 \leq n \leq T}$ is a $\mathbb{P}^*$ -martingale. $\square$

Note that Propositions 2.2 and 2.8 show that the binomial model has no arbitrage for $r \in (a, b)$. We will show in Proposition 2.10 that the binomial model is also complete. Hence, the fair price V_0 of any contingent claim of the form $C = f(S_T)$ is given by

$$V_0 = \frac{\mathbb{E}^*(C)}{(1+r)^T} = (1+r)^{-T} \sum_{k=0}^T \binom{T}{k} (1-p^*)^{T-k} p^{*k} f(S_0(1+a)^{T-k}(1+b)^k).$$

More generally, given a contingent claim $C = f(S_0, S_1, \dots, S_T)$, its fair price at time 0 is given by

$$V_0 = \frac{\mathbb{E}^*(C)}{(1+r)^T} = (1+r)^{-T} \sum f(s_0, s_1, \dots, s_T) \mathbb{P}^*(S_1 = s_1, \dots, S_T = s_T)$$

where we sum over the 2^T possible paths $(s_n)_{0 \leq n \leq T}$ starting from S_0 and

$$\mathbb{P}^*(S_1 = s_1, \dots, S_T = s_T) = (1-p^*)^{T-k} p^{*k}$$

where $k = |\{n \in \{1, \dots, T\} : s_n = (1+b)s_{n-1}\}|$. It is sometimes efficient to organize this calculation over a tree. Set

$$f_T(s_0, \dots, s_T) = f(s_0, \dots, s_T)$$

and carry out the backwards recursion for $n \leq T-1$

$$f_n(s_0, \dots, s_n) = (1-p^*)f_{n+1}(s_0, \dots, s_n, (1+a)s_n) + p^*f_{n+1}(s_0, \dots, s_n, (1+b)s_n).$$

The nodes of the tree are the partial paths $(s_0, \dots, s_n)$, each with two upward edges to $(s_0, \dots, s_n, (1+a)s_n)$ and $(s_0, \dots, s_n, (1+b)s_n)$. We start with the values at the leaves $(s_0, \dots, s_T)$ and compute down the tree to the root s_0 .

Proposition 2.9. *For $n = 0, 1, \dots, T$, almost surely,*

$$\mathbb{E}^*(f(S_0, \dots, S_T) | \mathcal{F}_n) = f_n(S_0, \dots, S_n)$$

and in particular

$$\mathbb{E}^*(C) = f_0(S_0).$$

Proof. The claim holds for $n = T$. Let $n \leq T-1$ and suppose for a reverse induction that the claim holds for $n+1$. Fix $s_0, \dots, s_n$ and define $A = \{S_0 = s_0, \dots, S_n = s_n\}$. Then

$$\begin{aligned} \mathbb{E}^*(f(S_0, \dots, S_T) | A) &= \mathbb{E}^*(f_{n+1}(S_0, \dots, S_n, S_{n+1}) | A) \\ &= (1-p^*)f_{n+1}(s_0, \dots, s_n, (1+a)s_n) + p^*f_{n+1}(s_0, \dots, s_n, (1+b)s_n) \\ &= f_n(s_0, \dots, s_n) = \mathbb{E}^*(f_n(S_0, \dots, S_n) | A). \end{aligned}$$

Now $f_n(S_0, \dots, S_n)$ is $\mathcal{F}_n$ -measurable, and every element of $\mathcal{F}_n$ is a finite union of such sets A . Hence, almost surely,

$$\mathbb{E}^*(f(S_0, \dots, S_T) | \mathcal{F}_n) = f_n(S_0, \dots, S_n)$$

and the induction proceeds. $\square$

We now show completeness for the binomial model by identifying a replicating portfolio for a general contingent claim $C = f(S_0, \dots, S_T)$. Define

$$\Delta_n(s_0, \dots, s_{n-1}) = \frac{f_n(s_0, \dots, s_{n-1}, (1+b)s_{n-1}) - f_n(s_0, \dots, s_{n-1}, (1+a)s_{n-1})}{(1+r)^{T-n}(b-a)s_{n-1}}.$$

Proposition 2.10. *The contingent claim $C = f(S_0, \dots, S_T)$ has a replicating portfolio $(\theta_n)_{1 \leq n \leq T}$ given by*

$$\theta_n = \Delta_n(S_0, \dots, S_{n-1}).$$

Proof. Define, for $n = 0, 1, \dots, T$,

$$V_n = \frac{\mathbb{E}^*(C|\mathcal{F}_n)}{(1+r)^T}.$$

Fix a path $(s_0, s_1, \dots, s_{n-1})$ starting from S_0 , set $\phi(x) = f_n(s_0, \dots, s_{n-1}, (1+x)s_{n-1})$ and consider the events

$$\Omega_0 = \{S_1 = s_1, \dots, S_{n-1} = s_{n-1}\}, \quad \Omega_a = \Omega_0 \cap \{R_n = a\}, \quad \Omega_b = \Omega_0 \cap \{R_n = b\}.$$

On Ω_a , we have

$$V_n = \frac{\phi(a)}{(1+r)^T}, \quad V_{n-1} = \frac{f_{n-1}(s_0, \dots, s_{n-1})}{(1+r)^T} = \frac{(1-p^*)\phi(a) + p^*\phi(b)}{(1+r)^T}$$

so

$$V_n - V_{n-1} = \frac{p^*(\phi(a) - \phi(b))}{(1+r)^T}$$

and

$$X_n - X_{n-1} = \frac{1+a}{1+r}X_{n-1} - X_{n-1} = \frac{(a-r)s_{n-1}}{(1+r)^n}$$

so

$$\theta_n(X_n - X_{n-1}) = \left(\frac{a-r}{b-a}\right) \frac{\phi(b) - \phi(a)}{(1+r)^T} = V_n - V_{n-1}.$$

By a similar calculation, the same identity holds on Ω_b , and hence everywhere, since $s_1, \dots, s_{n-1}$ are arbitrary. On summing over n , we obtain

$$D = V_0 + \sum_{n=1}^T \theta_n(X_n - X_{n-1}).$$

showing that $(\theta_n)_{1 \leq n \leq T}$ is a replicating portfolio for C , as claimed. $\square$

2.10 Pricing barrier options for a special binomial model

A reflection trick allows to obtain the joint distribution of a simple random walk and its maximum.

Proposition 2.11. *Let $(W_n)_{0 \leq n \leq T}$ be a simple random walk on the integers starting from 0, with $\mathbb{P}(W_1 = 1) = p$. Set $M_T = \max_{0 \leq n \leq T} W_n$. Then, for all integers $m, k \geq 0$, with $k \leq T$ and $2k - T \leq m \leq k$,*

$$\mathbb{P}(M_T = m \text{ and } W_T = 2k - T) = \left(\binom{T}{k-m} - \binom{T}{k-m-1} \right) p^k (1-p)^{T-k}.$$

Proof. Write E for the set of paths taken by the random walk. Thus

$$E = \{w = (w_n)_{0 \leq n \leq T} : w_0 = 0 \text{ and } w_n = w_{n-1} \pm 1 \text{ for } n = 1, \dots, T\}.$$

Consider the map $\phi : E \rightarrow E$ which reflects about level m the portion of the path (if any) after it first hits level m . Then $\phi \circ \phi$ is the identity, so ϕ is a bijection. Define

$$A = \{w \in E : \max_{0 \leq n \leq T} w_n \geq m \text{ and } w_T = 2k - T\}.$$

Then

$$\phi(A) = \{w \in E : w_T = 2m - 2k + T\}.$$

Hence

$$|A| = |\phi(A)| = \binom{T}{k-m}.$$

But the walk takes every path in A with equal probability $p^k (1-p)^{T-k}$. Hence

$$\mathbb{P}(M_T \geq m \text{ and } W_T = 2k - T) = \binom{T}{k-m} p^k (1-p)^{T-k}$$

and the result follows by subtracting the corresponding formula for $m+1$. $\square$

In the special case where $(1+a)(1+b) = 1$, we can realise the binomial model as a function of a simple random walk $(W_n)_{0 \leq n \leq T}$ on the integers by setting

$$S_n = S_0(1+b)^{W_n}.$$

Assume that the interest rate $r \in (a, b)$ and set $p^* = (r-a)/(b-a)$ as usual. Proposition 2.11 then gives an explicit formula for the fair price at time 0 of any contingent claim of the form

$$C = F \left(S_T, \max_{0 \leq n \leq T} S_n \right)$$

which is given by

$$V_0 = \frac{\mathbb{E}^*(C)}{(1+r)^T} = \sum_{k=0}^T \sum_{\substack{m=0 \\ m \geq 2k-T}}^k \frac{f(k, m)}{(1+r)^T} \left(\binom{T}{k-m} - \binom{T}{k-m-1} \right) p^{*k} (1-p^*)^{T-k}$$

where $f(k, m) = F(S_0(1+b)^{2k-T}, S_0(1+b)^m)$.

2.11 American calls and puts

An *American call* of expiry T confers the right but not the obligation to buy one unit of the underlying asset $(S_t)_{0 \leq n \leq T}$ for price K at any stopping time $\tau \leq T$ chosen by the holder. An *American put* of expiry T confers the right but not the obligation to sell one unit of the underlying asset $(S_t)_{0 \leq n \leq T}$ for price K at any stopping time $\tau \leq T$ chosen by the holder.

American options are not simple contingent claims such as discussed above, since the stopping time can be chosen by the holder, so the no-arbitrage pricing theory does not apply in general. However, if the asset price model is complete and arbitrage-free, then the holder of the option can never do better than choose the stopping time τ to maximise $\mathbb{E}^*(C_\tau/S_\tau^0)$, where $\mathbb{E}^*$ denotes expectation under the unique equivalent martingale measure. This is because any difference $C_{\tau_1}/S_{\tau_1}^0 - C_{\tau_2}/S_{\tau_2}^0$ can be replicated in the market for an initial cost of $\mathbb{E}^*(C_{\tau_1}/S_{\tau_1}^0) - \mathbb{E}^*(C_{\tau_2}/S_{\tau_2}^0)$. Hence, in this context, the pricing of American calls and puts reduces to the problem of maximizing $\mathbb{E}^*(C_\tau/S_\tau^0)$ over τ .

We will consider the pricing of these options when $(S_n)_{0 \leq n \leq T}$ is a binomial model of parameters $a < b$ with interest rate $r \in (a, b)$. Write, as usual, $\mathbb{P}^*$ for the equivalent martingale measure, corresponding to the parameter choice $p^* = (r - a)/(b - a)$.

Consider first the American call. Fix a stopping time $\tau \leq T$ and fix $n \in \{0, 1, \dots, T\}$. Note that

$$\mathbb{E}^*(S_T | \mathcal{F}_n) = (1 + r)^{T-n} S_n, \quad \text{almost surely.}$$

Consider the event

$$A = \{S_\tau \geq K \text{ and } \tau = n\}.$$

Then

$$\begin{aligned} \mathbb{E}^*(C_T 1_{\{\tau=n\}}) &\geq \mathbb{E}^*((S_T - K) 1_A) = \mathbb{E}^*((1 + r)^{T-n} S_n - K) 1_A \\ &\geq \mathbb{E}^*((1 + r)^{T-n} (S_n - K) 1_A) = \mathbb{E}^*(C_\tau 1_{\{\tau=n\}}). \end{aligned}$$

On summing over n , we see that $\mathbb{E}^*(C_T) \geq \mathbb{E}^*(C_\tau)$. Hence the choice $\tau = T$ is always optimal, making the American and European calls equivalent.

In order to find the fair price V_0 of the American put, we must solve the following optimal stopping problem: maximize

$$\mathbb{E}^*((1 + r)^{T-\tau} (K - S_\tau)^+)$$

over all stopping times $\tau \leq T$. At each time $n \in \{0, \dots, T-1\}$, the investor has two possible actions: to stop, receiving reward $(1 + r)^{T-n} (K - S_n)$, or to continue. If he has not already stopped, at time T he receives a reward of $(K - S_T)^+$. Define

$$E_n = \{S_0(1 + a)^{n-k}(1 + b)^k : k = 0, \dots, n\}.$$

The Bellman equation for this problem is then

$$\begin{aligned} v(T, x) &= (K - x)^+, \quad \text{for } x \in E_T, \\ v(n, x) &= \max\{(1 + r)^{T-n} (K - x), (1 - p^*)v(n + 1, x(1 + a)) + p^*v(n + 1, x(1 + b))\}, \\ &\quad \text{for } x \in E_n \text{ and } n = 0, \dots, T - 1. \end{aligned}$$

This can be solved by backwards recursion to find $v(0, S_0)$. Then

$$V_0 = \sup_{\tau} \mathbb{E}^*((1+r)^{T-\tau}(K - S_{\tau})^+) = v(0, S_0)$$

and the optimal stopping time τ is the smallest $n \in \{0, 1, \dots, T\}$ such that

$$(1+r)^{T-n}(K - S_n)^+ = v(n, S_n).$$

3 Brownian motion

3.1 Definition and basic properties

A real-valued random process $(B_t)_{t \geq 0}$ is said to be a *Brownian motion* if $B_0 = 0$ and

- (a) for all $s, t \geq 0$, the random variable $B_{s+t} - B_s$ is Gaussian, of mean 0 and variance t , and is independent of $\sigma(B_r : r \leq s)$,
- (b) for all $\omega \in \Omega$, the map $t \mapsto B_t(\omega) : [0, \infty) \rightarrow \mathbb{R}$ is continuous.

We sometimes replace the condition $B_0 = 0$ by $B_0 = x$, when we will refer to $(B_t)_{t \geq 0}$ as a Brownian motion starting from x . We often write $\mathbb{P}_x$ and $\mathbb{E}_x$ in this case as a reminder. If the starting point is not mentioned, then the default is always to start at 0.

Proposition 3.1. *Let $(B_t)_{t \geq 0}$ be a continuous random process starting from 0. Then the following are equivalent:*

- (a) $(B_t)_{t \geq 0}$ is a Brownian motion,
- (b) $(B_t)_{t \geq 0}$ is a zero-mean Gaussian process with $\mathbb{E}(B_s B_t) = s \wedge t$ for all $s, t \geq 0$.

Proof. Suppose (a) holds. To see that $(B_t)_{t \geq 0}$ is zero-mean Gaussian, it suffices to note that, for $0 = t_0 \leq t_1 \leq \dots \leq t_n$, the random variable $(B_{t_1}, \dots, B_{t_n})$ is a linear function of the independent zero-mean Gaussian random variables $(B_{t_1} - B_{t_0}, \dots, B_{t_n} - B_{t_{n-1}})$ and hence is Gaussian. For the covariance, we have, for $s \leq t$,

$$\mathbb{E}(B_s B_t) = \mathbb{E}(B_s^2) + \mathbb{E}(B_s(B_t - B_s)) = s$$

since $B_s \sim N(0, s)$ and $B_t - B_s$ is independent of B_s . Hence (b) holds.

Suppose on the other hand that (b) holds. Then, for $s, t \geq 0$, $B_{s+t} - B_s$ is a zero-mean Gaussian and

$$\mathbb{E}((B_{s+t} - B_s)^2) = (s+t) - 2s + s = t$$

so $B_{s+t} - B_s \sim N(0, t)$. To show that $B_{s+t} - B_s$ is independent of $\sigma(B_r : r \leq s)$, it suffices by a result from Probability and Measure to show that it is independent of $(B_{r_1}, \dots, B_{r_n})$ for all n and all $0 \leq r_1 \leq \dots \leq r_n \leq s$. Then, given that $(B_t)_{t \geq 0}$ is zero-mean Gaussian, it suffices to note that

$$\mathbb{E}(B_{r_k}(B_{t+s} - B_s)) = 0$$

for all k . Hence $(B_t)_{t \geq 0}$ is a Brownian motion. $\square$

The following proposition is left as an exercise.

Proposition 3.2 (Scaling property). *Let $(B_t)_{t \geq 0}$ be a Brownian motion and let $c \in (0, \infty)$. Set $B_t^{(c)} = c^{-1} B_{c^2 t}$. Then $(B_t^{(c)})_{t \geq 0}$ is also a Brownian motion.*

Proposition 3.3. *Let $(B_t)_{t \geq 0}$ be a Brownian motion. Then $(B_t)_{t \geq 0}$ exits every finite interval almost surely.*

Proof. Fix an interval I , of length L say. Consider the sequence of independent events $(A_n)_{n \geq 1}$ given by

$$A_n = \{|B_n - B_{n-1}| > L\}.$$

Then $\mathbb{P}(A_n) = p$ for all n , where $p = \mathbb{P}(|B_1| > L) > 0$. Then, by independence,

$$\mathbb{P}(A_1 \cup \dots \cup A_n) = 1 - (1 - p)^n.$$

So $\mathbb{P}(\cup_n A_n) = 1$. But, on A_n , if $(B_t)_{t \geq 0}$ has not left I by time $n - 1$, it must do so by time n . Hence $(B_t)_{t \geq 0}$ leaves I almost surely. $\square$

Write $\mathcal{F}_t = \sigma(B_s : s \in [0, t])$. A random time $T : \Omega \rightarrow [0, \infty]$ is a *stopping time* if

$$\{T \leq t\} \in \mathcal{F}_t \quad \text{for all } t.$$

Write $\mathcal{F}_T$ for the set of events $A \in \mathcal{F}_\infty$ such that

$$A \cap \{T \leq t\} \in \mathcal{F}_t \quad \text{for all } t.$$

We will need the following proposition which is proved in Advanced Probability.

Proposition 3.4 (Strong Markov property). *Let $(B_t)_{t \geq 0}$ be a Brownian motion and let T be an almost surely finite stopping time. Define $(\tilde{B}_t)_{t \geq 0}$ by*

$$\tilde{B}_t = B_{T+t} - B_T.$$

Then $(\tilde{B}_t)_{t \geq 0}$ is also a Brownian motion and is independent of $\mathcal{F}_T$.

Proposition 3.5. *Let $(B_t)_{t \geq 0}$ be a Brownian motion and let $a \in \mathbb{R}$. Define*

$$T_a = \inf\{t \geq 0 : B_t = a\}.$$

Then T_a is an almost surely finite stopping time.

Proof. It suffices to consider the case $a > 0$. Since $(B_t)_{t \geq 0}$ is continuous, we have

$$\{T_a \leq t\} = \{B_t = a\} \cup \bigcap_{r < a, r \in \mathbb{Q}} \bigcup_{s < t, s \in \mathbb{Q}} \{B_s > r\} \in \mathcal{F}_t.$$

Hence T_a is a stopping time. Set $T = T_1 \wedge T_{-1}$. Then T is a stopping time and, since Brownian motion leaves every interval, $T < \infty$ almost surely. By symmetry

$$\mathbb{P}(B_T = 1) = \mathbb{P}(B_T = -1) = 1/2.$$

Then, by the strong Markov property, the sequence of integers hit by $(B_t)_{t \geq 0}$, omitting immediate repeats, is a simple symmetric random walk. Since the random walk is recurrent, it follows that $T_a < \infty$ almost surely. $\square$

3.2 Wiener's theorem

We make the set of continuous paths $C[0, \infty)$ into a metric space by setting

$$d(w_1, w_2) = \sum_{n=1}^{\infty} 2^{-n} \left(\sup_{t \leq n} |w_1(t) - w_2(t)| \wedge 1 \right).$$

The Borel σ -algebra generated by the resulting open sets then makes $C[0, \infty)$ into a measurable space. For $t \geq 0$, define the *coordinate function* $X_t : C[0, \infty) \rightarrow \mathbb{R}$ by $X_t(w) = w(t)$. These are then continuous and together they generate the Borel σ -algebra. We call $(X_t)_{t \geq 0}$ the *coordinate process*. The following is proved in Advanced Probability.

Theorem 3.6 (Wiener's theorem). *There is a unique probability measure μ_W on $C[0, \infty)$ such that the coordinate process is a Brownian motion.*

The measure μ_W is called Wiener measure. We can consider any Brownian motion $(B_t)_{t \geq 0}$, as a $C[0, \infty)$ -valued random variable B , such that

$$X_t(B(\omega)) = B(\omega)(t) = B_t(\omega).$$

Then the law of B on $C[0, \infty)$ makes the coordinate process into a Brownian motion. Thus, any construction of a Brownian motion shows existence in Wiener's theorem, and uniqueness in Wiener's theorem establishes that all Brownian motions have the same law on $C[0, \infty)$.

3.3 Brownian motion as a limit of random walks

Brownian motion can be understood as a limit object, associated to random walks having steps of mean 0 and variance 1. We take a limit which scales down the time-step and the spatial steps of the walk in a coordinated way, so that the variance of the walk at time 1 is always 1. It turns out that, in this limit, all such random walks 'look almost the same' and all 'approximate' the behaviour of Brownian motion. The following is proved in Advanced Probability.

Theorem 3.7 (Donsker's invariance principle). *Let $(W_n)_{n \geq 0}$ be a random walk in $\mathbb{R}$ with steps of mean 0 and variance 1. Write $(W_t)_{t \geq 0}$ for the linear interpolation given by $W_{n+t} = (1-t)W_n + tW_{n+1}$ for $t \in [0, 1]$. Set*

$$W_t^{(k)} = k^{-1/2} W_{kt}.$$

Consider the sequence of random variables $W^{(k)} = (W_t^{(k)})_{t \geq 0}$ in $C[0, \infty)$. Then $W^{(k)}$ converges weakly to Wiener measure as $k \rightarrow \infty$. Thus, for all continuous bounded functions F on $C[0, \infty)$,

$$\mathbb{E}(F(W^{(k)})) \rightarrow \int_{C[0, \infty)} F(w) \mu_W(dw).$$

3.4 Change of probability measure

We now show that, on a finite time interval, the distribution of a Brownian motion with constant drift is absolutely continuous with respect to Wiener measure and we identify its density. This is a special case of the Cameron–Martin theorem.

Proposition 3.8. *Let $T \geq 0$ and $c \in \mathbb{R}$. Let $B = (B_t)_{t \in [0, T]}$ be a Brownian motion and define $\tilde{B} = (\tilde{B}_t)_{t \in [0, T]}$ by $\tilde{B}_t = B_t + ct$. Then, for all measurable sets $A \subseteq C[0, T]$, we have*

$$\mathbb{P}(\tilde{B} \in A) = \mathbb{E}(1_{\{B \in A\}} e^{cB_T - c^2 T/2}).$$

Proof. Consider the set $\mathcal{A}$ of all subsets of $C[0, T]$ of the form

$$A = \{x \in C[0, T] : x_{t_k} - x_{t_{k-1}} \in I_k \text{ for } k = 1, \dots, n\}$$

where $n \in \mathbb{N}$ and $0 = t_0 \leq t_1 \leq \dots \leq t_n = T$ and $I_1, \dots, I_n$ are intervals in $\mathbb{R}$. By a standard argument from Probability and Measure, it will suffice to prove the formula in the case $A \in \mathcal{A}$.

For $X \sim N(0, s)$ and $\tilde{X} = X + cs$, we have $\tilde{X} \sim N(cs, s)$ so, for any interval I ,

$$\begin{aligned} \mathbb{P}(\tilde{X} \in I) &= \int_I \frac{1}{\sqrt{2\pi s}} e^{-(x-cs)^2/(2s)} dx \\ &= \int_I \frac{1}{\sqrt{2\pi s}} e^{-x^2/(2s)} e^{cx - c^2 s/2} dx = \mathbb{E}(1_{\{X \in I\}} e^{cX - c^2 s/2}). \end{aligned}$$

Set $s_k = t_k - t_{k-1}$ and $X_k = B_{t_k} - B_{t_{k-1}}$ and $\tilde{X}_k = \tilde{B}_{t_k} - \tilde{B}_{t_{k-1}}$. Then $X_1, \dots, X_n$ are independent, with $X_k \sim N(0, s_k)$ and $\tilde{X}_k = X_k + cs_k$. Hence

$$\begin{aligned} \mathbb{P}(\tilde{B} \in A) &= \mathbb{P}\left(\prod_{k=1}^n 1_{\{\tilde{X}_k \in I_k\}}\right) = \prod_{k=1}^n \mathbb{P}(\tilde{X}_k \in I_k) \\ &= \prod_{k=1}^n \mathbb{E}(1_{\{X_k \in I_k\}} e^{cX_k - c^2 s_k/2}) = \mathbb{E}\left(\prod_{k=1}^n 1_{\{X_k \in I_k\}} e^{cX_k - c^2 s_k/2}\right) = \mathbb{E}\left(1_{\{B \in A\}} e^{cB_T - c^2 T/2}\right) \end{aligned}$$

where we used that $X_1 + \dots + X_n = B_T$ and $s_1 + \dots + s_n = T$. $\square$

3.5 Reflection principle

The reflection trick used in the proof of Proposition 2.11 has the following continuum analogue.

Proposition 3.9 (Reflection principle). *Let $(B_t)_{t \geq 0}$ be a Brownian motion and let $a \geq 0$. Set*

$$T_a = \inf\{t \geq 0 : B_t = a\}$$

and define $(\tilde{B}_t)_{t \geq 0}$ by

$$\tilde{B}_t = \begin{cases} B_t, & \text{if } t \leq T_a, \\ 2a - B_t, & \text{if } t > T_a. \end{cases}$$

Then $(\tilde{B}_t)_{t \geq 0}$ is also a Brownian motion.

Proof. Write T for T_a in the proof. We know that $T < \infty$ almost surely and T is a stopping time. Define processes $Y = (Y_t)_{t \geq 0}$ and $Z = (Z_t)_{t \geq 0}$ by

$$Y_t = B_{T \wedge t}, \quad Z_t = B_{T+t} - B_T.$$

Then Y_t is $\mathcal{F}_T$ -measurable for all t and, by the strong Markov property, Z is a Brownian motion independent of $\mathcal{F}_T$. So $-Z$ is also a Brownian motion independent of $\mathcal{F}_T$. Hence (Y, Z) and $(Y, -Z)$ have the same distribution on $C[0, \infty) \times C_0[0, \infty)$.

Consider the measurable map $F : C[0, \infty) \times C_0[0, \infty) \rightarrow C[0, \infty)$ given by

$$F(y, z) = \begin{cases} y(t), & \text{if } t \leq \tau(y), \\ y(\tau(y)) + z(t - \tau(y)), & \text{if } t > \tau(y) \end{cases}$$

where $\tau(y) = \inf\{t \geq 0 : y(t) = a\}$. Then $F(Y, Z) = B$ and $F(Y, -Z) = \tilde{B}$, so B and $\tilde{B}$ have the same distribution. $\square$

The reflection principle facilitates some useful calculations for Brownian motion. For example, consider the maximum process

$$M_t = \sup_{0 \leq s \leq t} B_s.$$

Then, for all $a \geq 0$ and $x \leq a$,

$$\{M_t \geq a \text{ and } B_t \leq x\} = \{\tilde{B}_t \geq 2a - x\}$$

so

$$\mathbb{P}(M_t \geq a \text{ and } B_t \leq x) = \mathbb{P}(B_t \geq 2a - x).$$

In particular, taking $x = a$, we see that

$$\mathbb{P}(M_t \geq a) = \mathbb{P}(B_t \geq a) + \mathbb{P}(M_t \geq a \text{ and } B_t \leq a) = 2\mathbb{P}(B_t \geq a).$$

Hence M_t has the same distribution as $|B_t|$. We can moreover compute the moment generating function. For

$$\mathbb{E}(e^{uM_1}) = 2 \int_0^\infty e^{ux} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx = 2e^{u^2/2} \int_0^\infty \frac{1}{\sqrt{2\pi}} e^{-(x-u)^2/2} dx = 2e^{u^2/2} \Phi(u)$$

and so, by scaling,

$$\mathbb{E}(e^{uM_t}) = \mathbb{E}(e^{u\sqrt{t}M_1}) = 2e^{u^2t/2} \Phi(u\sqrt{t}).$$

3.6 Hitting probabilities

Proposition 3.10. *Let $(B_t)_{t \geq 0}$ be a Brownian motion and let $a \geq 0$. Then T_a has a density function h_a on $[0, \infty)$ given by*

$$h_a(t) = \frac{a}{\sqrt{2\pi t^3}} e^{-a^2/(2t)}.$$

Proof. By the reflection principle,

$$\mathbb{P}(T_a \leq t) = \mathbb{P}(M_t \geq a) = 2\mathbb{P}(B_t \geq a).$$

But

$$\mathbb{P}(B_t \geq a) = \mathbb{P}(\sqrt{t}B_1 \geq a) = \int_{a/\sqrt{t}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy$$

and we can now differentiate in t to see that T_a has the claimed density. $\square$

3.7 Relation to the heat equation

Given a continuous bounded function f on $\mathbb{R}$, define for $t \geq 0$ and $x \in \mathbb{R}$

$$u(t, x) = \mathbb{E}(f(x + B_t))$$

where $(B_t)_{t \geq 0}$ is a Brownian motion. For $t > 0$, $x + B_t \sim N(x, t)$ so has density function $p(t, x, \cdot)$, where

$$p(t, x, y) = \frac{1}{\sqrt{2\pi t}} e^{-|x-y|^2/(2t)}.$$

Thus

$$u(t, x) = \int_{\mathbb{R}} p(t, x, y) f(y) dy.$$

Now p is the fundamental solution of the heat equation

$$\frac{\partial p}{\partial t} = \frac{1}{2} \frac{\partial^2 p}{\partial x^2}.$$

So, by differentiating under the integral sign,

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}$$

while, as $t \rightarrow 0$, since $B_t \rightarrow 0$, by bounded convergence,

$$u(t, x) \rightarrow f(x).$$

Hence u solves the Cauchy problem for the heat equation starting from f .

3.8 Transition density for killed Brownian motion

Proposition 3.11. *Let $x, a \in \mathbb{R}$ with $x \leq a$. Let $(B_t)_{t \geq 0}$ be a Brownian motion starting from x . Then, for all non-negative measurable functions f ,*

$$\mathbb{E}_x(f(B_t)1_{\{T_a > t\}}) = \int_{-\infty}^a f(y)p_t^a(x, y)dy$$

where

$$p_t^a(x, y) = p_t(x, y) - p_t(x, 2a - y).$$

Proof. By a standard argument, it suffices to consider the case where $f = 1_{(-\infty, b]}$ for some $b \leq a$. Set $M_t = \sup_{s \leq t} B_s$. Then, using the reflection principle,

$$\begin{aligned} \mathbb{E}_x(f(B_t)1_{\{T_a > t\}}) &= \mathbb{P}_x(M_t < a \text{ and } B_t \leq b) \\ &= \mathbb{P}_x(B_t \leq b) - \mathbb{P}_x(M_t \geq a \text{ and } B_t \leq b) \\ &= \mathbb{P}_x(B_t \leq b) - \mathbb{P}_x(B_t \geq 2a - b) \\ &= \int_{-\infty}^b p_t(x, y)dy - \int_{-\infty}^b p_t(x, 2a - y)dy = \int_{-\infty}^a f(y)p_t^a(x, y)dy. \end{aligned}$$

□

By change of measure we can deduce an analogous result for Brownian motion with drift. Fix $c \in \mathbb{R}$ and set

$$\tilde{B}_t = B_t + ct, \quad \tilde{T}_a = \inf\{t \geq 0 : \tilde{B}_t = a\}.$$

Then

$$\mathbb{E}_x(f(\tilde{B}_t)1_{\{\tilde{T}_a > t\}}) = \mathbb{E}_x(f(B_t)e^{c(B_t - x) - c^2 t/2}1_{\{T_a > t\}}) = \int_{-\infty}^a f(y)\tilde{p}_t^a(x, y)dy.$$

where

$$\tilde{p}_t^a(x, y) = e^{c(y - x) - c^2 t/2}p_t^a(x, y).$$

4 Black–Scholes model

4.1 Black–Scholes pricing formula

By a *Black–Scholes model* we mean any pair of processes $(S_t^0)_{t \geq 0}$ and $(S_t)_{t \geq 0}$ of the form

$$S_t^0 = e^{rt}, \quad S_t = S_0 e^{\sigma B_t + \mu t}$$

where $(B_t)_{t \geq 0}$ is a Brownian motion and where $r, \mu \in \mathbb{R}$ and $\sigma, S_0 \in (0, \infty)$. We interpret $(S_t^0)_{t \geq 0}$ as the price of a *riskless bond of interest rate r* . We interpret $(S_t)_{t \geq 0}$ as the price of a *risky asset*. We call μ the *drift* and σ the *volatility*.

The process $(e^{\sigma B_t - \sigma^2 t/2})_{t \geq 0}$ is a martingale. Hence for the special choice of drift

$$\mu^* = r - \sigma^2/2.$$

the discounted asset price $(e^{-rt} S_t)_{t \geq 0}$ is a martingale.

Proposition 4.1. *Let $(S_t^0, S_t)_{t \geq 0}$ be a Black–Scholes model of interest rate r , drift μ and volatility σ . Fix $T \in (0, \infty)$ and consider the equivalent probability measure $\mathbb{P}^*$ given by*

$$\frac{d\mathbb{P}^*}{d\mathbb{P}} = e^{\lambda B_T - \lambda^2 T/2}, \quad \sigma \lambda = \mu^* - \mu = r - \sigma^2/2 - \mu.$$

Then, under $\mathbb{P}^$, the discounted asset price $(e^{-rt} S_t)_{0 \leq t \leq T}$ is a martingale.*

Proof. Set $B_t^* = B_t - \lambda t$. By Proposition 3.8, under $\mathbb{P}^*$, $(B_t)_{0 \leq t \leq T}$ is a Brownian motion with drift λ , so $(B_t^*)_{0 \leq t \leq T}$ is a Brownian motion. Now

$$\sigma B_t^* + \mu^* t = \sigma B_t - \sigma \lambda t + \mu^* t = \sigma B_t + \mu t$$

so

$$e^{-rt} S_t = S_0 e^{\sigma B_t + \mu t - rt} = S_0 e^{\sigma B_t^* + \mu^* t - rt} = S_0 e^{\sigma B_t^* - \sigma^2 t/2}.$$

Hence, under $\mathbb{P}^*$, $(e^{-rt} S_t)_{0 \leq t \leq T}$ is a martingale. $\square$

We will abuse notation in writing $\mathbb{P}^*$ instead of $\mathbb{P}$ when considering a Black–Scholes model with drift μ^* .

By a *time- T contingent claim*, we mean a non-negative $\mathcal{F}_T$ -measurable random variable C , considered as an amount payable to the investor at time T . Let us assume that an investor is free to trade in the asset and the bond. The *Black–Scholes price* V_0 for the claim C is defined by

$$V_0 = e^{-rT} \mathbb{E}^*(C). \tag{1}$$

An argument which we will not develop shows that V_0 is the unique fair price for C , in the sense that any other price presents an opportunity to make risk-free profits by buying or selling the claim and trading continuously in the asset and the bond. We have not made precise what it means for the investor to trade freely in the asset and the bond. However, the following examples give some plausibility to the assertion that V_0 is the unique fair

price for C . In the case $C = S_T$, the investor can replicate C by buying one unit of the asset at time 0, so S_T must have time-0 price S_0 . On the other hand, in the case of a constant claim $C = K$, the investor can replicate C by investing $e^{-rT}K$ in the bond at time 0, so K (at time T) must have time-0 price $e^{-rT}K$. In both cases, the price is the Black–Scholes price.

More generally, by a *simple replicable claim* we mean any claim C such that

$$e^{-rT}C = C_0 + \sum_{k=1}^n \theta_k (X_{t_k} - X_{t_{k-1}})$$

for some constant C_0 , some $n \in \mathbb{N}$ and $0 = t_0 \leq t_1 \leq \dots \leq t_n = T$, with θ_k a bounded $\mathcal{F}_{t_{k-1}}$ -measurable random variable for all k . Here X_t is the discounted asset price $e^{-rt}S_t$. We leave as an exercise to show that any such claim C can be replicated at time T for cost C_0 at time 0. Since $(X_t)_{0 \leq t \leq T}$ is a martingale under $\mathbb{P}^*$, we can take expectations to obtain

$$V_0 = e^{-rT} \mathbb{E}^*(C) = C_0.$$

Hence V_0 is the unique fair price for C .

In fact, by the Brownian martingale representation theorem, every $\mathcal{F}_T$ -measurable contingent claim is the limit in probability of simple replicable claims, and this can be used to justify the validity of the pricing formula in general.

4.2 Black–Scholes PDE

For some simple types of option, the Black–Scholes pricing theory can be cast in terms of PDE. In this section, we show how the stochastic theory reduces to a PDE for options depending only on the final asset price. Consider a Black–Scholes model with interest rate r and risky asset $(S_t)_{t \geq 0}$ having drift μ and volatility σ . We derive first the form of the backward equation for the transition density of $(S_t)_{t \geq 0}$. Consider the case $S_0 = s$. Then

$$\log S_t = \log s + \sigma B_t + \mu t$$

so $\log S_t$ has density $p(\sigma^2 t, x(t), \cdot)$ on $\mathbb{R}$, where

$$p(t, x, z) = \frac{1}{\sqrt{2\pi t}} e^{-|x-z|^2/(2t)}, \quad x(t) = \log s + \mu t.$$

Write $\rho(t, s, \cdot)$ for the density function of S_t on $(0, \infty)$ when $S_0 = s$. We will write $\dot{\rho}$ and ρ' for derivatives in the first and second argument respectively. We have

$$y\rho(t, s, y) = p(\sigma^2 t, x(t), z), \quad z = \log y$$

so

$$sy\rho'(t, s, y) = p'(\sigma^2 t, x(t), z), \quad sy(\rho' + s\rho'')(t, s, y) = p''(\sigma^2 t, x(t), z)$$

and so

$$\dot{\rho}(t, s, y) = y^{-1} (\sigma^2 \dot{p} + \mu p') = y^{-1} \left(\frac{1}{2} \sigma^2 p'' + \mu p' \right) = \frac{1}{2} \sigma^2 s^2 \rho''(t, s, y) + \left(\mu + \frac{1}{2} \sigma^2 \right) s \rho'(t, s, y).$$

Proposition 4.2. *Let F be a continuous function on $(0, \infty)$ of polynomial growth. For $t \in [0, T]$ and $s \in (0, \infty)$, write $V(t, s)$ for the time- t value of the time- T contingent claim $F(S_T)$, conditional on $S_t = s$, as given by the Black–Scholes pricing formula*

$$V(t, s) = e^{-r(T-t)} \mathbb{E}^*(F(S_T) | S_t = s) = e^{-r(T-t)} \mathbb{E}(F(se^{\sigma B_{T-t} + \mu^*(T-t)})).$$

Then V is continuous on $[0, T] \times (0, \infty)$ and $C^{1,2}$ on $[0, T) \times (0, \infty)$ and satisfies the Black–Scholes PDE

$$\mathcal{L}V = \dot{V} + \frac{1}{2}\sigma^2 s^2 V'' + rsV' - rV = 0$$

with terminal value $V(., T) = F$.

Proof. Set

$$v(t, s, y) = e^{-r(T-t)} \rho(T - t, s, y)$$

where ρ is the transition density for the case of drift $\mu^* = r - \sigma^2/2$. Then

$$\dot{v}(t, s, y) = e^{-r(T-t)} (r\rho - \dot{\rho})(T - t, s, y)$$

and

$$\dot{\rho} = \frac{1}{2}\sigma^2 s^2 \rho'' + rs\rho'$$

so

$$\dot{v} = rv - \frac{1}{2}\sigma^2 s^2 v'' - rsv'$$

and so

$$\mathcal{L}v = \dot{v} + \frac{1}{2}\sigma^2 s^2 v'' + rsv' - rv = 0.$$

Now

$$V(t, s) = \int_0^\infty F(y) v(t, s, y) dy$$

and the growth condition on F , combined with super-polynomial decay in $v(t, s, .)$, allows to differentiate under the integral sign to obtain

$$\mathcal{L}V(t, s) = \int_0^\infty F(y) \mathcal{L}v(t, s, y) dy = 0.$$

Finally, continuity of V on $[0, T] \times (0, \infty)$ follows from continuity of Brownian motion by dominated convergence. $\square$

4.3 Binomial approximation to Black–Scholes

Fix $\mu, r \in \mathbb{R}$ and $\sigma \in (0, \infty)$ and consider the Black–Scholes model

$$S_t^0 = e^{rt}, \quad S_t = S_0 e^{\sigma B_t + \mu t}.$$

We will show, using Donsker's invariance principle, that this model may be viewed as a weak limit of suitably chosen binomial models, and that the Black–Scholes pricing formula then matches the limiting binomial pricing formula.

Let $(W_n)_{n \geq 0}$ be a simple symmetric random walk on the integers. Form the rescaled linear interpolation $(W_t^{(k)})_{t \geq 0}$ as in Donsker's invariance principle. For k sufficiently large, we can determine $a_k < r_k < b_k$ by

$$1 + a_k = \exp\left(-\frac{\sigma}{\sqrt{k}} + \frac{\mu}{k}\right), \quad 1 + b_k = \exp\left(\frac{\sigma}{\sqrt{k}} + \frac{\mu}{k}\right), \quad 1 + r_k = \exp\left(\frac{r}{k}\right).$$

Set

$$S_t^{(k)} = S_0 \exp\left(\sigma W_t^{(k)} + \mu t\right).$$

Then $(S_{n/k}^0, S_{n/k}^{(k)})_{n \geq 0}$ is a binomial model of parameters $a_k < r_k < b_k$ and $p = 1/2$. By Donsker's invariance principle, $W^{(k)} \rightarrow B$ weakly on $C[0, \infty)$. Hence $S^{(k)} \rightarrow S$ weakly on $C[0, \infty)$, as desired.

We will now show that the equivalent martingale measures for these binomial models converge to that for the Black–Scholes model. First we recall the change of probability formula for the binomial model. Fix a time interval $[0, T]$ and a convergent sequence $\lambda_k \rightarrow \lambda$ in $\mathbb{R}$. For convenience, we will assume that T is an integer. Set

$$p = \frac{1}{2}, \quad p_k = \frac{1}{2} \left(1 + \frac{\lambda_k}{\sqrt{k}}\right).$$

We can and do choose k sufficiently large that $p_k \in (0, 1)$. Write

$$U_T^{(k)} = \frac{kT + \sqrt{k}W_T^{(k)}}{2}, \quad D_T^{(k)} = \frac{kT - \sqrt{k}W_T^{(k)}}{2}$$

for the numbers of up-steps and down-steps and define

$$Z_T^{(k)} = \left(\frac{1-p_k}{1-p}\right)^{D_T^{(k)}} \left(\frac{p_k}{p}\right)^{U_T^{(k)}} = \left(1 - \frac{\lambda_k}{\sqrt{k}}\right)^{D_T^{(k)}} \left(1 + \frac{\lambda_k}{\sqrt{k}}\right)^{U_T^{(k)}}.$$

Consider the equivalent probability measure $\tilde{\mathbb{P}}^{(k)}$ given by

$$\frac{d\tilde{\mathbb{P}}^{(k)}}{d\mathbb{P}} = Z_T^{(k)}.$$

Under $\tilde{\mathbb{P}}^{(k)}$, $(S_{n/k}^0, S_{n/k}^{(k)})_{0 \leq n \leq kT}$ is a binomial model of parameters $a_k < r_k < b_k$ and p_k . Consider the limit $k \rightarrow \infty$. We have

$$Z_T^{(k)} = \left(1 - \frac{\lambda_k^2}{k}\right)^{kT} \left(\left(1 - \frac{\lambda_k}{\sqrt{k}}\right)^{-\sqrt{k}/2} \left(1 + \frac{\lambda_k}{\sqrt{k}}\right)^{\sqrt{k}/2}\right)^{W_T^{(k)}}$$

and

$$\left(1 - \frac{\lambda_k^2}{k}\right)^{kT} \rightarrow e^{-\lambda^2 T}, \quad \left(1 \pm \frac{\lambda_k}{\sqrt{k}}\right)^{\pm\sqrt{k}/2} \rightarrow e^{\lambda/2}$$

so $Z_T^{(k)} \rightarrow Z_T$ weakly, where $Z_T = e^{\lambda B_T - \lambda^2 T/2}$. Consider the change of measure for B given by

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = Z_T.$$

Then, for any continuous bounded function F on $C[0, T]$,

$$\tilde{\mathbb{E}}^{(k)}(F(S^{(k)})) = \mathbb{E}(Z_T^{(k)} F(S^{(k)})) \rightarrow \mathbb{E}(Z_T F(S)) = \tilde{\mathbb{E}}(F(S)).$$

We have seen that the equivalent martingale measures are obtained when $\lambda = \lambda^*$ for Black–Scholes and $\lambda_k = \lambda_k^*$ for the binomial model, where

$$\sigma\lambda^* = r - \frac{\sigma^2}{2} - \mu, \quad \frac{1}{2} \left(1 + \frac{\lambda_k^*}{\sqrt{k}}\right) = \frac{r_k - a_k}{b_k - a_k}.$$

It is left as an exercise, for the choices of a_k, b_k and r_k made above, to check that $\lambda_k^* \rightarrow \lambda^*$ so that

$$\tilde{\mathbb{E}}^{(k)*}(F(S^{(k)})) \rightarrow \tilde{\mathbb{E}}^*(F(S))$$

giving an exact tie-up between the discrete-time and continuous-time theories.

4.4 Black–Scholes formula for the price of a European call

The *European call* of *maturity* T and *strike* $K \geq 0$ is the contingent claim $(S_T - K)^+$. This may be thought of as conferring the right but not the obligation to buy one unit of stock at time T at price K . We will compute a formula for the fair value of this claim at time 0, in the Black–Scholes model. The fair value is given by

$$EC(S_0, K, \sigma, r, T) = \mathbb{E}^*(e^{-rT}(S_T - K)^+).$$

Recall that we write Φ for the standard normal distribution function, given by

$$\Phi(a) = \int_{-\infty}^a \phi(y) dy, \quad \phi(y) = \frac{1}{\sqrt{2\pi}} e^{-y^2/2}$$

and that

$$\bar{\Phi}(a) = 1 - \Phi(a) = \Phi(-a).$$

Proposition 4.3 (Black–Scholes formula). *We have*

$$EC(S_0, K, \sigma, r, T) = S_0 \Phi(d_+) - e^{-rT} K \Phi(d_-)$$

where

$$d_{\pm} = \frac{\log(S_0/K) + rT}{\sigma\sqrt{T}} \pm \frac{\sigma\sqrt{T}}{2}.$$

Proof. It suffices to consider the case $\mu = \mu^*$, when $\mathbb{P}^* = \mathbb{P}$. Then $B_T \sim \sqrt{T}B_1$ and $B_1 \sim N(0, 1)$. We compute

$$\begin{aligned} EC(S_0, K, \sigma, r, T) &= \mathbb{E} \left(e^{-rT} (S_T - K)^+ \right) = \mathbb{E} \left((S_0 e^{\sigma B_T - \sigma^2 T/2} - e^{-rT} K)^+ \right) \\ &= \int_a^\infty (S_0 e^{\sigma \sqrt{T}y - \sigma^2 T/2} - e^{-rT} K) \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \end{aligned}$$

where a is given by $S_0 e^{\sigma \sqrt{T}a - \sigma^2 T/2} = e^{-rT} K$ so

$$\log(S_0/K) + \sigma \sqrt{T}a + rT = \sigma^2 T/2$$

and so $a = -d_-$. Hence

$$\begin{aligned} EC(S_0, K, \sigma, r, T) &= S_0 \int_a^\infty \frac{1}{\sqrt{2\pi}} e^{-(y - \sigma \sqrt{T})^2/2} dy - e^{-rT} K \int_a^\infty \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \\ &= S_0 \bar{\Phi}(a - \sigma \sqrt{T}) - e^{-rT} K \bar{\Phi}(a) \\ &= S_0 \Phi(-a + \sigma \sqrt{T}) - e^{-rT} K \Phi(-a) \\ &= S_0 \Phi(d_+) - e^{-rT} K \Phi(d_-). \end{aligned}$$

□

The *European put of maturity* $T > 0$ and *strike* $K \geq 0$ on $(S_t)_{t \geq 0}$ is the contingent claim $(S_T - K)^-$. This may be thought of as conferring the right but not the obligation to sell one unit of stock at time T at price K . In the Black-Scholes model $(S_t^0, S_t)_{t \geq 0}$ the fair value of this claim at time 0 is given by

$$EP(S_0, K, \sigma, r, T) = \mathbb{E}^* \left(e^{-rT} (S_T - K)^- \right).$$

Note the *put-call parity equation*

$$(S_T - K)^+ - (S_T - K)^- = S_T - K.$$

Thus a portfolio long in the call and short in the put is equivalent to the forward contract $S_T - K$ of maturity T and strike K . The fair value of the forward contract is

$$\mathbb{E}^* \left(e^{-rT} (S_T - K) \right) = S_0 - e^{-rT} K.$$

Hence we obtain

$$EP(S_0, K, \sigma, r, T) = EC(S_0, K, \sigma, r, T) - S_0 + e^{-rT} K = e^{-rT} K \bar{\Phi}(d_-) - S_0 \bar{\Phi}(d_+).$$

4.5 Sensitivities for contingent claims in the Black–Scholes model

Recall that the value of a contingent claim C at time T in the Black–Scholes model $(S_t^0, S_t)_{t \geq 0}$ is given by

$$v(S_0) = v(S_0, C, \sigma, r, T) = \mathbb{E}^*(e^{-rT}C).$$

where $\mathbb{E}^*$ denotes expectation with respect to the equivalent martingale measure $\mathbb{P}^*$. The derivatives of v with respect to parameters of the model are known as *sensitivities*. The following terminology is widely used

$$\text{Delta} = \Delta = \frac{\partial v}{\partial S_0}, \quad \text{Gamma} = \Gamma = \frac{\partial^2 v}{\partial S_0^2}, \quad \text{Vega} = \mathcal{V} = \frac{\partial v}{\partial \sigma}, \quad \text{Rho} = \rho = \frac{\partial v}{\partial r}.$$

These and other sensitivities are also known as *Greeks*.

In the case of a European call of maturity T and strike K , the Black–Scholes formula can be differentiated explicitly to obtain

$$\Delta = \Phi(d_+), \quad \mathcal{V} = S_0 \phi(d_+) \sqrt{T}.$$

4.6 Implied volatility

Proposition 4.4. *The map $\sigma \mapsto EC(S_0, K, \sigma, r, T)$ is an increasing bijection $(0, \infty) \rightarrow ((S_0 - e^{-rT}K)^+, S_0)$.*

Proof. It is straightforward to check that, for all values of S_0, K, r and T , we have

$$\lim_{\sigma \rightarrow 0} EC(S_0, K, \sigma, r, T) = (S_0 - e^{-rT}K)^+, \quad \lim_{\sigma \rightarrow \infty} EC(S_0, K, \sigma, r, T) = S_0.$$

Since

$$\frac{\partial}{\partial \sigma} EC(S_0, K, \sigma, r, T) = \mathcal{V} = S_0 \phi(d_+) \sqrt{T} > 0$$

this implies the claim. $\square$

European calls are widely traded in financial markets, so the actual market price may be observed, for a range of maturities T and strikes K . Since the current stock price S_0 and interest rate r are also known, if we accept the Black–Scholes model, this determines a unique *implied volatility* $\sigma_{\text{implied}}(K, T)$ such that

$$EC(S_0, K, \sigma_{\text{implied}}(K, T), r, T) = EC_{\text{market}}(K, T).$$

In principle, $\sigma_{\text{implied}}(K, T)$ should be independent of K and T . In practice, the implied volatility surface is not flat, indicating that the Black–Scholes model is not an exact fit. However, implied volatility, acting as an encoding of the call price $EC_{\text{market}}(K, T)$, is a convenient and widely used means to quote these market prices.

4.7 Pricing of exotic options in the Black–Scholes model

The reflection principle allows us to obtain pricing formulas for certain barrier options in the Black–Scholes model. We illustrate this by the case of the up-and-out call

$$C = h(S_T)1_{\{\sup_{0 \leq t \leq T} S_t < A\}}, \quad h(s) = (s - K)^+$$

where $A \geq \max\{S_0, K\}$. The fair price at time 0 for C is given by the Black–Scholes pricing formula

$$V_0 = e^{-rT} \mathbb{E}^*(C)$$

where $\mathbb{P}^*$ is the equivalent martingale measure, corresponding to drift $\mu^* = r - \sigma^2/2$. In calculating V_0 , we will assume that $\mu = \mu^*$ and drop the stars. Then

$$S_t = S_0 e^{\sigma B_t + rt - \sigma^2 t/2} = S_0 e^{\sigma \tilde{B}_t}, \quad \tilde{B}_t = B_t + ct, \quad c = (r - \sigma^2/2)/\sigma$$

so

$$\left\{ \sup_{0 \leq t \leq T} S_t < A \right\} = \left\{ \sup_{0 \leq t \leq T} \tilde{B}_t < a \right\} = \{\tilde{T}_a > t\}, \quad \tilde{T}_a = \inf\{t \geq 0 : \tilde{B}_t = a\}$$

where $a > 0$ is determined by $S_0 e^{\sigma a} = A$. We showed in Section 3.8 that

$$\mathbb{E} \left(f(\tilde{B}_t) 1_{\{\tilde{T}_a > t\}} \right) = \int_{-\infty}^a f(y) e^{cy - c^2 t/2} p_t^a(0, y) dy$$

where

$$p_t^a(0, y) = p_t(0, y) - p_t(0, 2a - y).$$

Hence

$$V_0 = e^{-rT} \int_{-\infty}^a h(S_0 e^{\sigma y}) e^{cy - c^2 T/2} p_T^a(0, y) dy.$$

In the special case $h(s) = (s - K)^+$, the right-hand side can be expressed in terms of the normal distribution function Φ by using the identity

$$\int_a^b e^{\lambda y} p_t(0, y) dy = e^{\lambda^2 t/2} \left(\Phi \left(\frac{b - \lambda t}{\sqrt{t}} \right) - \Phi \left(\frac{a - \lambda t}{\sqrt{t}} \right) \right).$$

5 Appendix

Recall the statement of Theorem 2.3 for an asset price model $(\bar{S}_n)_{0 \leq n \leq T}$. *The following are equivalent:*

- (a) $(\bar{S}_n)_{0 \leq n \leq T}$ has no arbitrage,
- (b) $(\bar{S}_n)_{0 \leq n \leq T}$ has an equivalent martingale measure.

This is the discrete-time version of the fundamental theorem of asset pricing. We will now prove this statement – the proof is not examinable. The main step is to show the following conditional version of the equivalence shown in the proof given above of the fundamental theorem in the case $T = 1$.

Proposition 5.1. *Let Y be a random variable in $\mathbb{R}^d$ and let $\mathcal{F}_0$ be a sub- σ -algebra of $\mathcal{F}$. Then the following are equivalent:*

- (a) *there exists no $\mathcal{F}_0$ -measurable random variable Θ in $\mathbb{R}^d$ such that $\Theta \cdot Y \geq 0$ almost surely and $\Theta \cdot Y > 0$ with positive probability,*
- (b) *there exists an equivalent probability measure $\tilde{\mathbb{P}}$ such that $\tilde{\mathbb{P}} = \mathbb{P}$ on $\mathcal{F}_0$ and, almost surely,*

$$\tilde{\mathbb{E}}(|Y| | \mathcal{F}_0) < \infty, \quad \tilde{\mathbb{E}}(Y | \mathcal{F}_0) = 0.$$

Moreover, given any finite non-negative random variable U , we can choose $\tilde{\mathbb{P}}$ in (b) so that $\tilde{\mathbb{E}}(U | \mathcal{F}_0) < \infty$ almost surely.

Proof. Suppose that (a) holds. It will suffice to consider the case where, almost surely, for all $\theta \in \mathbb{R}^d$,

$$\mathbb{E}(e^{\theta \cdot Y} | \mathcal{F}_0) < \infty, \quad \mathbb{E}(U e^{\theta \cdot Y} | \mathcal{F}_0) < \infty.$$

For the general case, we can replace $\mathbb{P}$ by the equivalent probability measure $\tilde{\mathbb{P}}$ given by $d\tilde{\mathbb{P}}/d\mathbb{P} = W/\mathbb{E}(W | \mathcal{F}_0)$ where $W = e^{-|Y|^2} e^{-U}$ for which $\tilde{\mathbb{P}} = \mathbb{P}$ on $\mathcal{F}_0$ and, almost surely,

$$\tilde{\mathbb{E}}(e^{\theta \cdot Y} | \mathcal{F}_0) < \infty, \quad \tilde{\mathbb{E}}(U e^{\theta \cdot Y} | \mathcal{F}_0) < \infty.$$

We will assume that this has been done and drop the tildes.

There are $\mathcal{F}_0$ -measurable kernels M and N on $\Omega \times \mathcal{B}(\mathbb{R}^d)$ such that, for all $B \in \mathcal{B}(\mathbb{R}^d)$, almost surely,

$$M(B) = \mathbb{P}(Y \in B | \mathcal{F}_0), \quad N(B) = \mathbb{E}(U 1_{\{Y \in B\}} | \mathcal{F}_0).$$

We write here $M(B)$ for the random variable $M(\cdot, B)$ and will use similar notation elsewhere. Define $\mathcal{F}_0$ -measurable fields of subspaces of $\mathbb{R}^d$ by

$$E_0(\omega) = \{\theta \in \mathbb{R}^d : M(\omega, \{\theta \cdot y = 0\}) = 1\}, \quad E_1(\omega) = \text{span}(\text{supp } M(\omega, \cdot)) = E_0(\omega)^\perp.$$

Write $P(\omega)$ for the orthogonal projection $\mathbb{R}^d \rightarrow E_1(\omega)$. Fix $\theta \in \mathbb{R}^d$ and consider the event

$$B(\theta) = \{P\theta \neq 0\} = \{\theta \notin E_0\}.$$

Fix $B \in \mathcal{F}_0$ with $B \subseteq B(\theta)$ and $\mathbb{P}(B) > 0$. Consider the $\mathcal{F}_0$ -measurable random variable $\Theta = 1_B \theta$. Define

$$W = M(\{y \in \mathbb{R}^d : \theta \cdot y \neq 0\}), \quad W' = M(\{y \in \mathbb{R}^d : \theta \cdot y > 0\}).$$

Then, almost surely,

$$W = \mathbb{P}(\theta \cdot Y \neq 0 | \mathcal{F}_0), \quad W' = \mathbb{P}(\theta \cdot Y > 0 | \mathcal{F}_0).$$

Now $W > 0$ on $B(\theta)$, so $\mathbb{P}(\Theta \cdot Y \neq 0) = \mathbb{E}(1_B W) > 0$. Condition (a) then implies that $\mathbb{E}(1_B W') = \mathbb{P}(\Theta \cdot Y > 0) > 0$ so, since B is arbitrary, $W' > 0$ almost surely on $B(\theta)$.

Set $\psi(t) = 0 \vee t \wedge 1$ and define $F : \Omega \times \mathbb{R}^d \rightarrow [0, \infty)$ by

$$F(\omega, \theta) = \int_{\mathbb{R}^d} \psi(\theta \cdot y) M(\omega, dy).$$

Then $F(\omega, \cdot)$ is continuous on $\mathbb{R}^d$ for all ω , $F(\theta) = \mathbb{E}(\psi(\theta \cdot Y) | \mathcal{F}_0)$ almost surely, and $F(\theta) > 0$ almost surely on $B(\theta)$ for all θ . Hence, by Fubini's theorem, $F(\theta) > 0$ for Lebesgue almost all $\theta \in \mathbb{R}^d \setminus E_0$, almost surely. Since F is continuous on $\mathbb{R}^d$ and $\mathbb{R}^d \setminus E_0$ is open, this implies that $F(\theta) > 0$ for all $\theta \in \mathbb{R}^d \setminus E_0$ almost surely. Set $S(\omega) = \{\theta \in E_1(\omega) : |\theta| = 1\}$. Then $S(\omega)$ is compact. Define

$$\varepsilon(\omega) = \frac{1}{2} \inf\{F(\omega, \theta) : \theta \in S(\omega)\}, \quad \Omega_0 = \{\omega \in \Omega : \varepsilon(\omega) > 0\}.$$

Then ε is $\mathcal{F}_0$ -measurable, $\mathbb{P}(\Omega_0) = 1$ and, for all $\omega \in \Omega_0$ and $\theta \in S(\omega)$,

$$\begin{aligned} \int_{\mathbb{R}^d} 1_{\{\theta \cdot y \geq \varepsilon(\omega)\}} M(\omega, dy) &\geq \int_{\mathbb{R}^d} \psi(\theta \cdot y - \varepsilon(\omega)) M(\omega, dy) \\ &\geq \int_{\mathbb{R}^d} \psi(\theta \cdot y) M(\omega, dy) - \varepsilon(\omega) = F(\omega, \theta) - \varepsilon(\omega) \geq \varepsilon(\omega). \end{aligned}$$

Define functions $\Phi : \Omega \times \mathbb{R}^d \rightarrow (0, \infty)$ and $\Psi : \Omega \times \mathbb{R}^d \rightarrow [0, \infty)$ by

$$\Phi(\omega, \theta) = \int_{\mathbb{R}^d} e^{\theta \cdot y} M(\omega, dy), \quad \Psi(\omega, \theta) = \int_{\mathbb{R}^d} e^{\theta \cdot y} N(\omega, dy).$$

Then Φ is $\mathcal{F}_0$ -measurable on Ω , differentiable on $\mathbb{R}^d$, and strictly convex on E_1 , with

$$\Phi'(\omega, \theta) = \int_{\mathbb{R}^d} y e^{\theta \cdot y} M(\omega, dy)$$

and, almost surely,

$$\Phi(\theta) = \mathbb{E}(e^{\theta \cdot Y} | \mathcal{F}_0), \quad \Phi'(\theta) = \mathbb{E}(Y e^{\theta \cdot Y} | \mathcal{F}_0).$$

Similarly, Ψ is $\mathcal{F}_0$ -measurable on Ω , continuous on $\mathbb{R}^d$ and, almost surely,

$$\Psi(\theta) = \mathbb{E}(U e^{\theta \cdot Y} | \mathcal{F}_0).$$

Then, for $\omega \in \Omega_0$, $\theta_0 \in E_0(\omega)$, $\theta_1 \in S(\omega)$ and $t \geq (1/\varepsilon(\omega)) \log(1/\varepsilon(\omega))$,

$$\Phi(\omega, \theta_0 + t\theta_1) = \Phi(\omega, t\theta_1) \geq \varepsilon e^{t\varepsilon} \geq 1 = \Phi(\omega, 0).$$

Hence, for all $\omega \in \Omega_0$, there exists a unique $\Theta^*(\omega) \in E_1(\omega)$ such that $\Phi(\omega, \theta) \geq \Phi(\omega, \Theta^*(\omega))$ for all $\theta \in \mathbb{R}^d$. Then $\Phi'(\omega, \Theta^*(\omega)) = 0$. Set $\Phi^*(\omega) = 0$ for $\omega \notin \Omega_0$. Define recursively a sequence of $\mathcal{F}_0$ -measurable random variables $(\Theta_n : n \geq 0)$ in $\mathbb{R}^d$ by

$$\Theta_0 = 0, \quad \Theta_{n+1} = \Theta_n - \gamma \Phi'(\Theta_n).$$

Then $\Theta_n \in E_1$ for all n almost surely, so by the properties of gradient descent, for γ sufficiently small, we have $\Theta_n \rightarrow \Theta^*$ almost surely. Hence Θ^* is an $\mathcal{F}_0$ -measurable random variable.

Define an equivalent probability measure $\tilde{\mathbb{P}}$ by $d\tilde{\mathbb{P}}/d\mathbb{P} \propto e^{\Theta^* \cdot Y}$. Then, almost surely,

$$\tilde{\mathbb{E}}(U|\mathcal{F}_0) = \mathbb{E}(Ue^{\Theta^* \cdot Y}|\mathcal{F}_0) = \Psi(\Theta^*) < \infty$$

and similarly $\tilde{\mathbb{E}}(|Y||\mathcal{F}_0) < \infty$ almost surely. For $\omega \in \Omega_0$, since $\Phi(\omega, \cdot)$ is minimized at $\Theta^*(\omega)$, we have $\Phi'(\omega, \Theta^*(\omega)) = 0$. So, for all $B \in \mathcal{F}_0$,

$$\mathbb{E}(Y1_B e^{\Theta^* \cdot Y}|\mathcal{F}_0) = 1_B \mathbb{E}(Y e^{\Theta^* \cdot Y}|\mathcal{F}_0) = 1_B \Phi'(\Theta^*) = 0$$

and so

$$\tilde{\mathbb{E}}(Y1_B) = \frac{\mathbb{E}(Y1_B e^{\Theta^* \cdot Y})}{\mathbb{E}(e^{\Theta^* \cdot Y})} = 0.$$

Hence $\tilde{\mathbb{E}}(Y|\mathcal{F}_0) = 0$ almost surely and (b) holds.

Suppose on the other hand that (b) holds and let Θ be an $\mathcal{F}_0$ -measurable random variable in $\mathbb{R}^d$. Then, almost surely,

$$\tilde{\mathbb{E}}(\Theta \cdot Y|\mathcal{F}_0) = \Theta \cdot \tilde{\mathbb{E}}(Y|\mathcal{F}_0) = 0$$

and so $\tilde{\mathbb{E}}(\Theta \cdot Y) = 0$. Hence, if $\Theta \cdot Y \geq 0$ almost surely, then $\Theta \cdot Y = 0$ almost surely. Hence (a) holds. $\square$

Proof of Theorem 2.3. Suppose that (a) holds. Write $Y_n = X_n - X_{n-1}$. Consider the following hypothesis: there is a random variable ρ_n such that, almost surely, $\rho_n > 0$ and $\mathbb{E}(\rho_n|\mathcal{F}_{n-1}) = 1$ and $U_n < \infty$ and $\mathbb{E}(Y_k \rho_n|\mathcal{F}_{k-1}) = 0$ for $k = n, \dots, T$, where we write U_n for a version of

$$\sum_{k=n}^T \mathbb{E}(|Y_k| \rho_n|\mathcal{F}_{n-1}).$$

The hypothesis holds for $n = T+1$ with $\rho_{T+1} = 1$, the third and fourth conditions holding vacuously. Suppose, for a reverse induction, that the hypothesis holds for $n+1$. Given an $\mathcal{F}_{n-1}$ -measurable random variable Θ in $\mathbb{R}^d$ such that $\Theta \cdot Y_n \geq 0$ almost surely, we obtain a

previsible process $(\theta_1, \dots, \theta_T)$ in $\mathbb{R}^d$ by setting $\theta_k = \Theta 1_{\{k=n\}}$. The associated value process satisfies

$$V_T = \sum_{k=1}^T \theta_k \cdot Y_k = \Theta \cdot Y_n \geq 0.$$

The no-arbitrage condition (a) then forces $\Theta \cdot Y_n = 0$ almost surely. Hence, condition (a) of Proposition 5.1 holds for the $\mathcal{F}_n$ -measurable random variable Y_n and the sub- σ -algebra $\mathcal{F}_{n-1}$. Then, by Proposition 5.1, there is an $\mathcal{F}_n$ -measurable random variable α_n such that, almost surely, $\alpha_n > 0$ and $\mathbb{E}(\alpha_n | \mathcal{F}_{n-1}) = 1$ and

$$\mathbb{E}(|Y_n| \alpha_n | \mathcal{F}_{n-1}) < \infty, \quad \mathbb{E}(Y_n \alpha_n | \mathcal{F}_{n-1}) = 0$$

and moreover we can choose α_n so that $\mathbb{E}(U_n \alpha_n | \mathcal{F}_{n-1}) < \infty$ almost surely. Define

$$\rho_n = \alpha_n \rho_{n+1}.$$

Then, almost surely, $\rho_n > 0$ and

$$\mathbb{E}(\rho_n | \mathcal{F}_{n-1}) = \mathbb{E}(\alpha_n \mathbb{E}(\rho_{n+1} | \mathcal{F}_n) | \mathcal{F}_{n-1}) = \mathbb{E}(\alpha_n | \mathcal{F}_{n-1}) = 1$$

and

$$U_n = \mathbb{E}(U_{n+1} \alpha_n | \mathcal{F}_{n-1}) + \mathbb{E}(|Y_n| \alpha_n | \mathcal{F}_{n-1}) < \infty$$

and, for $k = n+1, \dots, T$,

$$\mathbb{E}(Y_k \rho_n | \mathcal{F}_{k-1}) = \rho_n \mathbb{E}(Y_k | \mathcal{F}_{k-1}) = 0$$

while

$$\mathbb{E}(Y_n \rho_n | \mathcal{F}_{n-1}) = \mathbb{E}(Y_n \alpha_n \mathbb{E}(\rho_{n+1} | \mathcal{F}_n) | \mathcal{F}_{n-1}) = \mathbb{E}(Y_n \alpha_n | \mathcal{F}_{n-1}) = 0.$$

Hence the hypothesis holds for n and the induction proceeds.

Since the random variables $\mathbb{E}(|Y_k| \rho_1 | \mathcal{F}_0)$ are all finite-valued and $\mathcal{F}_0$ -measurable, and so is X_0 , there is an $\mathcal{F}_0$ -measurable random variable α_0 such that $\alpha_0 > 0$ almost surely, $\mathbb{E}(\alpha_0) = 1$ and

$$\alpha_0 \left(|X_0| + \sum_{k=1}^T \mathbb{E}(|Y_k| \rho_1 | \mathcal{F}_0) \right)$$

is integrable. We can define an equivalent probability measure $\tilde{\mathbb{P}}$ by $d\tilde{\mathbb{P}}/d\mathbb{P} = \alpha_0 \rho_1$. Then, under $\tilde{\mathbb{P}}$, the random variable $|X_0| + |Y_1| + \dots + |Y_T|$ is integrable and $\tilde{\mathbb{E}}(Y_k | \mathcal{F}_{k-1}) = 0$ for $k = 1, \dots, T$. Hence $(X_n)_{0 \leq n \leq T}$ is a martingale under $\tilde{\mathbb{P}}$. Hence (b) holds. The reverse implication is the content of Proposition 2.2. $\square$