

ANALYSIS 1 EXAMPLES SHEET 2

Lent Term 2014

W. T. G.

1. Let (a_n) and (b_n) be two real sequences. Suppose that (a_n) is a subsequence of (b_n) and (b_n) is a subsequence of (a_n) . Suppose also that (a_n) converges. Does it follow that they are the same sequence?

2. Let $H : \mathbb{R} \rightarrow \mathbb{R}$ be defined as follows: if $x < 0$ then $H(x) = 0$ and if $x \geq 0$ then $H(x) = 1$. Prove carefully that H is not continuous (i) by directly using the definition of continuity and (ii) by using the sequence definition.

3. Suppose that $f(x) \rightarrow \ell$ as $x \rightarrow a$ and $g(y) \rightarrow k$ as $y \rightarrow \ell$. Does it follow that $g(f(x)) \rightarrow k$ as $x \rightarrow a$?

4. For each natural number n , let $f_n : [0, 1] \rightarrow [0, 1]$ be a continuous function, and for each n let h_n be defined by $h_n(x) = \max\{f_1(x), \dots, f_n(x)\}$. Show that for each n the function h_n is continuous on $[0, 1]$. Must the function h defined by $h(x) = \sup\{f_n(x) : n \in \mathbb{N}\}$ be continuous?

5. Let $g : [0, 1] \rightarrow [0, 1]$ be a continuous function. Prove that there exists some $c \in [0, 1]$ such that $g(c) = c$. Such a c is called a *fixed point* of g .

Give an example of a bijection $h : [0, 1] \rightarrow [0, 1]$ with no fixed point.

Give an example of a continuous bijection $p : (0, 1) \rightarrow (0, 1)$ with no fixed point.

6. Prove that the function $q(x) = 2x^5 + 3x^4 + 2x + 16$ (defined on the reals) takes the value 0 exactly once, and that the number where it takes that value is somewhere in the interval $[-2, -1]$.

7. Assuming standard properties of trigonometric functions, prove rigorously that there are exactly 143 solutions to the simultaneous equations $x = \sin(6\pi y)$ and $y = \sin(6\pi x)$. That is, prove that there are exactly 143 ordered pairs (x, y) such that the two equations are satisfied. (Hint: you should start by drawing a diagram to understand where the number 143 comes from, but this diagram does not constitute part of the proof.)

8. Let $f : [0, 1] \rightarrow \mathbb{R}$ be continuous, with $f(0) = f(1) = 0$. Suppose that for every $x \in (0, 1)$ there exists $\delta > 0$ such that both $x + \delta$ and $x - \delta$ belong to $(0, 1)$ and $f(x) = \frac{1}{2}(f(x - \delta) + f(x + \delta))$. Prove that $f(x) = 0$ for every $x \in [0, 1]$.
9. Define a function $f : \mathbb{R} \rightarrow \mathbb{R}$ as follows. If x is irrational, then $f(x) = 0$, while if x is rational, then $f(x) = 1/q$, where q is the denominator of x . (That is, $x = p/q$, with p and q coprime integers and $q > 0$.) Prove that f is continuous at every irrational and discontinuous at every rational.
10. Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) . Which of the following statements are always true and which are sometimes false?
- (i) If f is increasing, then $f'(x) \geq 0$ for every $x \in (a, b)$.
 - (ii) If $f'(x) \geq 0$ for every $x \in (a, b)$, then f is increasing.
 - (iii) If f is strictly increasing, then $f'(x) > 0$ for every $x \in (a, b)$.
 - (iv) If $f'(x) > 0$ for every $x \in (a, b)$, then f is strictly increasing.
11. (i) Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function such that $g(0) = g'(0) = 0$ and $g''(0)$ exists and is positive. Prove that there exists $x > 0$ such that $g(x) > 0$.
- (ii) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function such that $f(0) = 0$, and $f''(0)$ exists and is positive. Prove that there exists $x > 0$ such that $f(2x) > 2f(x)$.
12. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable everywhere. Prove that if $f'(x) \rightarrow \ell$ as $x \rightarrow \infty$, then $f(x)/x \rightarrow \ell$. If $f(x)/x \rightarrow \ell$ as $x \rightarrow \infty$, does it follow that $f'(x) \rightarrow \ell$?
13. Find a function $f : \mathbb{R} \rightarrow \mathbb{R}$ that takes every value in every interval. That is, for every $a < b$ and every t there should exist $x \in (a, b)$ such that $f(x) = t$.
14. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function that has the intermediate value property: that is, if $f(a) < c < f(b)$ then there exists $x \in (a, b)$ such that $f(x) = c$. Suppose also that for every rational r the set $S_r = \{x : f(x) = r\}$ is closed. (This means that if (x_n) is any convergent sequence in S_r , then its limit also belongs to S_r .) Prove that f is continuous.