

1. Let  $(x_n)$  be a real sequence.
  - (i) Show that  $x_n \rightarrow -\infty$  if and only if  $-x_n \rightarrow \infty$ .
  - (ii) Show that if  $x_n \neq 0$  for all  $n$  and  $x_n \rightarrow \infty$  then  $\frac{1}{x_n} \rightarrow 0$ .
  - (iii) If  $x_n \neq 0$  for all  $n$  and  $\frac{1}{x_n} \rightarrow 0$ , does it follow that  $x_n \rightarrow \infty$ ?
2. Let  $x_1 > y_1 > 0$  and for every  $n \geq 1$  let  $x_{n+1} = (x_n + y_n)/2$  and  $y_{n+1} = 2x_n y_n / (x_n + y_n)$ . Show that  $x_n > x_{n+1} > y_{n+1} > y_n$ . Deduce that  $(x_n)$  and  $(y_n)$  converge to a common limit. What is that limit?
3. For each  $n \in \mathbb{N}$  a closed interval  $[x_n, y_n]$  is given. Assume that  $[x_m, y_m] \cap [x_n, y_n] \neq \emptyset$  for all  $m, n \in \mathbb{N}$ . Show that  $\bigcap_{n=1}^{\infty} [x_n, y_n] \neq \emptyset$ .
4. Give an example of a divergent sequence  $(x_n)$  with  $x_n - x_{n+1} \rightarrow 0$  as  $n \rightarrow \infty$ . Can such a sequence be bounded?
5. Let  $(x_n)$  and  $(y_n)$  be sequences such that  $(x_n)$  is a subsequence of  $(y_n)$  and  $(y_n)$  is a subsequence of  $(x_n)$ . Does it follow that  $x_n = y_n$  for all  $n$ ? Does your answer change if we further assume that  $(x_n)$  is convergent?
6. Let  $x$  be a real or complex number. Assume that every subsequence of a sequence  $(x_n)$  has a further subsequence that converges to  $x$ . Deduce that  $(x_n)$  converges to  $x$ .
7. Let  $(x_n)$  be a real sequence. Let  $L$  be the set of those  $x \in \mathbb{R}$  for which there is a subsequence of  $(x_n)$  that converges to  $x$ . Which of the following subsets of  $\mathbb{R}$  can occur as  $L$ :  $\emptyset$ ,  $\{0\}$ ,  $\{0, 1\}$ ,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$ ? Give examples or proofs as appropriate. Show further that if  $(x_n)$  is bounded but not convergent then  $L$  contains at least two elements.
8. The two series  $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$  and  $1 + \frac{1}{3} - \frac{1}{2} + \frac{1}{5} + \frac{1}{7} - \frac{1}{4} + \dots$  have the same terms in different orders. Let  $s_n$  and, respectively,  $t_n$  be the  $n^{\text{th}}$  partial sums of these series. Set  $h_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$ . Show that  $s_{2n} = h_{2n} - h_n$  and  $t_{3n} = h_{4n} - \frac{1}{2}h_{2n} - \frac{1}{2}h_n$ . Show that  $(s_n)$  converges to a limit  $s$  and  $(t_n)$  tends to  $3s/2$ .

9. Investigate the convergence of the following series. For each expression that contains the variable  $z$ , find all complex numbers  $z$  for which the series converges.

$$\sum_n \frac{\sin n}{n^2} \quad \sum_n \frac{n^2 z^n}{5^n} \quad \sum_n \frac{(-1)^n}{4 + \sqrt{n}} \quad \sum_n \frac{z^n(1-z)}{n} \quad \sum_{n \geq 3} \frac{n^2}{(\log \log n)^{\log n}}$$

10. Show that  $\sum_{n \geq 2} \frac{1}{n(\log n)^\alpha}$  converges for  $\alpha > 1$  and diverges otherwise. Does  $\sum_{n \geq 3} \frac{1}{n \log n \log \log n}$  converge?

11. Let  $x_n > 0$  and  $y_n > 0$  for all  $n \in \mathbb{N}$ . Assume that for some  $N \in \mathbb{N}$  we have

$$\frac{x_{n+1}}{x_n} \leq \frac{y_{n+1}}{y_n} \quad \text{for all } n \geq N.$$

Show that if  $\sum y_n$  converges, then so does  $\sum x_n$ .

12. Can you enumerate  $\mathbb{Q}$  as  $q_1, q_2, \dots$  so that the series  $\sum (q_n - q_{n+1})^2$  is convergent? How about  $\sum |q_n - q_{n+1}|$ ?

13. Let  $(x_n)$  and  $(y_n)$  be real sequences.

(i) Suppose  $x_n \rightarrow 0$  as  $n \rightarrow \infty$ . Show that there is a sequence  $(\varepsilon_n)$  of signs (*i.e.*,  $\varepsilon_n \in \{-1, +1\}$  for all  $n$ ) such that  $\sum \varepsilon_n x_n$  is convergent.

(ii) Suppose  $x_n \rightarrow 0$  and  $y_n \rightarrow 0$ . Must there be a sequence  $(\varepsilon_n)$  of signs such that  $\sum \varepsilon_n x_n$  and  $\sum \varepsilon_n y_n$  are both convergent?

14. Let  $S$  be a (possibly infinite) set of odd positive integers. Prove that there exists a real sequence  $(x_n)$  such that, for each positive integer  $k$ , the series  $\sum_{n=1}^{\infty} x_n^k$  converges when  $k$  belongs to  $S$  and diverges otherwise.