

1. Let $f: [a, b] \rightarrow \mathbb{R}$ be an integrable function with $f(x) \geq 0$ for every x . Assume further that f is continuous. Show that $\int_a^b f(t) dt = 0$ iff $f(x) = 0$ for every x . Does this hold without the assumption of continuity?
2. Give an example of a function $f: [0, 1] \rightarrow \mathbb{R}$ such that $|f|$ is integrable but f is not. Give an example of a sequence $f_n: [0, 1] \rightarrow [0, 1]$ of integrable functions such that the function $f(x) = \sup_n f_n(x)$ is not integrable.
3. Let $f: [0, 1] \rightarrow \mathbb{R}$ be defined by $f(x) = 0$ when x is irrational, and $f(x) = 1/q$ when $x = p/q$ is a rational written in its lowest terms. Prove that f is integrable on $[0, 1]$. What is $\int_0^1 f(x) dx$?
4. Let $a < b$ and f be an integrable function on $[a, b]$. Show that for every closed subinterval $I \subset [a, b]$ of positive length and every $\varepsilon > 0$ there exists a closed subinterval $J \subset I$ of positive length such that $\sup_J f - \inf_J f < \varepsilon$. Use this to show that if $f(x) > 0$ for every x then $\int_a^b f(x) dx > 0$.
5. Let f be a continuous function on $[a, b]$ and let $a < c < d < b$. Prove that

$$\lim_{h \rightarrow 0} \int_c^d \frac{f(x+h) - f(x)}{h} dx = f(d) - f(c) .$$

6. Let $f: [0, 1] \rightarrow \mathbb{R}$ be continuous. Let $G(x, t) = t(x-1)$ when $t \leq x$ and $x(t-1)$ when $t \geq x$. Let $g(x) = \int_0^1 G(x, t)f(t) dt$. Show that $g''(x)$ exists for $x \in (0, 1)$ and equals $f(x)$.
7. Which of the following improper integrals converge?
 - (i) $\int_1^\infty \frac{\log x}{1+x^2} dx$.
 - (ii) $\int_0^\infty x^p \exp(-x^q) dx$ (where $p, q > 0$).
 - (iii) $\int_0^\infty \sin(x^2) dx$.

8. Give an example of a continuous function $f: [0, \infty) \rightarrow [0, \infty)$ such that $\int_0^\infty f(x) dx$ exists but f is unbounded.

9. Prove that $\frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \rightarrow \log 2$ as $n \rightarrow \infty$, and find the limit of

$$\frac{1}{n+1} - \frac{1}{n+2} + \cdots + \frac{(-1)^{n-1}}{2n} .$$

10. For each non-negative integer n let $I_n(\theta) = \int_{-1}^1 (1-x^2)^n \cos(\theta x) dx$. Prove that $\theta^2 I_n = 2n(2n-1)I_{n-1} - 4n(n-1)I_{n-2}$ for all $n \geq 2$, and hence that $\theta^{2n+1} I_n(\theta) = n!(P_n(\theta) \sin \theta + Q_n(\theta) \cos \theta)$ for some pair P_n, Q_n of polynomials of degree at most $2n$ with integer coefficients. Deduce that π is irrational.

11. For each $n \in \mathbb{N}$ let $u_n = \int_0^{\pi/2} \sin 2nx \cot x dx$ and $v_n = \int_0^{\pi/2} \frac{\sin 2nx}{x} dx$. Prove that $u_n = \pi/2$. By considering the limit of v_n and of $u_n - v_n$ as $n \rightarrow \infty$, show that $\int_0^\infty \frac{\sin x}{x} dx = \frac{\pi}{2}$.

12. A function $f: [a, b] \rightarrow \mathbb{R}$ is said to have *bounded variation* if there exists $K \geq 0$ such that for every dissection $\mathcal{D}: a = x_0 < x_1 < \cdots < x_n = b$ of $[a, b]$, we have $\sum_{k=1}^n |f(x_k) - f(x_{k-1})| \leq K$. Prove that a function of bounded variation is integrable. Is the converse true?

13. For a dissection $\mathcal{D}: a = x_0 < x_1 < \cdots < x_n = b$ of a closed bounded interval $[a, b]$ define $|\mathcal{D}| = \max_k (x_k - x_{k-1})$. Assume that $f: [a, b] \rightarrow \mathbb{R}$ is integrable and (\mathcal{D}_n) is a sequence of dissections of $[a, b]$ with $|\mathcal{D}_n| \rightarrow 0$ as $n \rightarrow \infty$. Prove that $S_{\mathcal{D}_n}(f) - s_{\mathcal{D}_n}(f) \rightarrow 0$ as $n \rightarrow \infty$. Deduce that if \mathcal{D}_n is the sequence $a = x_0^{(n)} < x_1^{(n)} < \cdots < x_{m_n}^{(n)} = b$ and $\xi_k^{(n)} \in [x_{k-1}^{(n)}, x_k^{(n)}]$ for each $1 \leq k \leq m_n$, then

$$\sum_{k=1}^{m_n} f(\xi_k^{(n)}) (x_k^{(n)} - x_{k-1}^{(n)}) \rightarrow \int_a^b f(t) dt \quad \text{as } n \rightarrow \infty .$$

14⁺. Let $f: [0, 1] \rightarrow \mathbb{R}$ be a function that is differentiable everywhere (with right and left derivatives at the end points) with a derivative f' that is bounded. Must f' be integrable?