

## ANALYSIS II (Michaelmas 2010): EXAMPLES 2

The questions are not equally difficult and the ‘additional’ ones are marked with \*. Unless stated otherwise, the norm on  $\mathbb{R}^n$  may be taken to be the Euclidean norm  $\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$ , and the spaces  $\ell_0$  and  $\ell_\infty$  may be assumed to have the sup-norm  $\|x\|_\infty = \sup_i |x_i|$ . ( $\ell_0$  denotes the space of real sequences  $(x_n)_{n=1}^\infty$  such that all but finitely many  $x_n$  are zero.) Comments, corrections are welcome at any time and may be sent to [a.j.scholl@dpmmms.cam.ac.uk](mailto:a.j.scholl@dpmmms.cam.ac.uk).

1. Let  $(x^{(m)})$  and  $(y^{(m)})$  be sequences in  $\mathbb{R}^n$  converging to  $x$  and  $y$  respectively. Show that  $x^{(m)} \cdot y^{(m)}$  converges to  $x \cdot y$ . Deduce that if  $f : \mathbb{R}^n \rightarrow \mathbb{R}^p$  and  $g : \mathbb{R}^n \rightarrow \mathbb{R}^p$  are continuous at  $x \in \mathbb{R}^n$ , then so is the pointwise scalar product function  $f \cdot g : \mathbb{R}^n \rightarrow \mathbb{R}$ .

2. Show that  $\|x\|_1 = \sum_{i=1}^n |x_i|$  defines a norm on  $\mathbb{R}^n$ . Show directly that it is Lipschitz equivalent to the Euclidean norm.

3. (a) Show that  $\|f\|_1 = \int_0^1 |f(x)|dx$  defines a norm on the space  $C[0, 1]$ . Is it Lipschitz equivalent to the uniform norm?

(b) Let  $R[0, 1]$  denote the vector space of all integrable functions on  $[0, 1]$ . Does  $\|f\| = \int_0^1 |f(x)|dx$  define a norm on  $R[0, 1]$ ?

4. Which of the following subsets of  $\mathbb{R}^2$  are open? Which are closed? (And why?)

- (i)  $\{(x, 0) : 0 \leq x \leq 1\}$ ;
- (ii)  $\{(x, 0) : 0 < x < 1\}$ ;
- (iii)  $\{(x, y) : y \neq 0\}$ ;
- (iv)  $\{(x, y) : x \in \mathbb{Q} \text{ or } y \in \mathbb{Q}\}$ ;
- (v)  $\{(x, y) : y = nx \text{ for some } n \in \mathbb{N}\} \cup \{(x, y) : x = 0\}$ ;
- (vi)  $\{(x, f(x)) : x \in \mathbb{R}\}$ , where  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function.

5. Is the set  $\{f : f(1/2) = 0\}$  closed in the space  $C[0, 1]$  with the uniform norm? What about the set  $\{f : \int_0^1 f(x)dx = 0\}$ ? In each case, does the answer change if we replace the uniform norm with the norm  $\|\cdot\|_1$  defined in Question 3?

6. Which of the following functions  $f$  are continuous?

- (i) The linear map  $f : \ell_\infty \rightarrow \mathbb{R}$  defined by  $f(x) = \sum_{n=1}^\infty x_n/n^2$ .
- (ii) The identity map from the space  $C[0, 1]$  with the uniform norm to the space  $C[0, 1]$  with the norm  $\|\cdot\|_1$  defined in Question 3.
- (iii) The identity map from  $C[0, 1]$  with the norm  $\|\cdot\|_1$  to  $C[0, 1]$  with the uniform norm.
- (iv) The linear map  $f : \ell_0 \rightarrow \mathbb{R}$  defined by  $f(x) = \sum_{i=1}^\infty x_i$ .

7. If  $A$  and  $B$  are subsets of  $\mathbb{R}^n$ , we write  $A + B$  for the set  $\{a + b : a \in A, b \in B\}$ . Show that if  $A$  and  $B$  are both closed and one of them is bounded then  $A + B$  is closed. Give an example in  $\mathbb{R}^1$  to show that the boundedness condition cannot be omitted. If  $A$  and  $B$  are both open, is  $A + B$  necessarily open? Justify your answer.

8. (a) Show that the space  $\ell_\infty$  is complete. Show also that  $c_0 = \{x \in \ell_\infty : x_n \rightarrow 0\}$ , the vector subspace of  $\ell_\infty$  consisting of all sequences converging to 0, is complete.

(b) Is the space  $R[0, 1]$  of integrable functions on  $[0, 1]$ , equipped with the uniform norm, complete?

9. Let  $\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear map. Show that  $\|x\|' = \|x\| + \|\alpha x\|$  defines a norm on  $\mathbb{R}^n$ . Using the fact that all norms on a finite-dimensional space are Lipschitz equivalent, deduce that  $\alpha$  is continuous.

**10.\*** Which of the following vector spaces of functions, considered with the uniform norm, are complete? (Justify your answer.)

- (i) The space  $C_b(\mathbb{R})$  of bounded continuous functions  $f : \mathbb{R} \rightarrow \mathbb{R}$ .
- (ii) The space  $C_0(\mathbb{R})$  of continuous functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) \rightarrow 0$  as  $|x| \rightarrow \infty$ .
- (iii) The space  $C_c(\mathbb{R})$  of continuous functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = 0$  for  $|x|$  sufficiently large.

**11.** In lectures we proved that if  $E$  is a closed and bounded set in  $\mathbb{R}^n$ , then any continuous function defined on  $E$  has bounded image. Prove the converse: if every continuous real-valued function on  $E \subseteq \mathbb{R}^n$  is bounded, then  $E$  is closed and bounded.

**12.** Let  $(x^{(m)})_{m \geq 1}$  be a bounded sequence in  $\ell_\infty$ . Show that there is a subsequence  $(x^{(m_j)})_{j \geq 1}$  which converges in every coordinate; that is to say, the sequence  $(x_i^{(m_j)})_{j \geq 1}$  of real numbers converges for each  $i$ . Why does this not show that every bounded sequence in  $\ell_\infty$  has a convergent subsequence?

**13.** Show that  $\|x\|_1 = \sum_{i=1}^\infty |x_i|$  defines a norm on  $\ell_0$  and that this norm is not Lipschitz equivalent to the uniform norm  $\|\cdot\|$ . Find a third norm on  $\ell_0$  which is neither Lipschitz equivalent to  $\|\cdot\|_1$ , nor to  $\|\cdot\|$ . Is it possible to find uncountably many norms on  $\ell_0$  such that no two are Lipschitz equivalent?

**14.** Let  $V$  be a normed space in which every bounded sequence has a convergent subsequence. (a) Show that  $V$  must be complete. (b)\* Show further that  $V$  must be finite-dimensional.

[Hint for (b): Show first that for every finite-dimensional subspace  $V_0$  of  $V$  there exists an  $x \in V$  with  $\|x + y\| > \|x\|/2$  for each  $y \in V_0$ .]

**15.\*** Recall from the lectures the normed space  $\ell_2$ . The Hilbert cube is the subset of  $\ell_2$  consisting of all the sequences  $(x_n)_{n=1}^\infty$  such that for each  $n$ ,  $|x_n| \leq 1/n$ . Show that the Hilbert cube is closed in  $\ell_2$ , and that it has the Bolzano–Weierstrass property, that is, any sequence in the Hilbert cube has a convergent subsequence. (So the Hilbert cube is *compact*.)