

EXAMPLE SHEET 2

1. Let V be a normed space.
 - (a) If $(\mathbf{v}_n) \rightarrow \mathbf{v} \in V$, show that any subsequence of (\mathbf{v}_n) converges to \mathbf{v} .
 - (b) If (\mathbf{v}_n) is a Cauchy sequence in V , show that it is bounded.
2. Suppose $(f_n) \rightarrow f$ pointwise, where $f_n, f \in C[0, 1]$.
 - (a) If $(f_n) \rightarrow f$ uniformly and (x_m) is a sequence of points in $[0, 1]$ converging to x , show that $(f_n(x_m)) \rightarrow f(x)$.
 - (b) If (f_n) does not converge uniformly, show that there is a convergent sequence $(x_m) \rightarrow x \in [0, 1]$ such that $(f_n(x_m))$ does not converge to $f(x)$. (Hint: Bolzano-Weierstrass.)
3. If A and B are subsets of \mathbb{R}^n , let $A + B = \{a + b \mid a \in A, b \in B\}$. Show that if A and B are both closed and one of them is bounded, then $A + B$ is closed. Give an example in \mathbb{R} to show that the boundedness condition cannot be omitted. If A and B are both open, is $A + B$ necessarily open? Justify your answer.
4. Let $(V, \|\cdot\|)$ be a complete normed space, and let W be a linear subspace of V . Show that $(W, \|\cdot\|)$ is complete if and only if W is a closed subset of V . Which of the following vector spaces of functions on \mathbb{R} are complete with respect to the uniform norm?
 - (a) The space $C_b(\mathbb{R})$ of bounded continuous functions on \mathbb{R} .
 - (b) The space $C_0(\mathbb{R})$ of continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) \rightarrow 0$ as $|x| \rightarrow \infty$.
 - (c) The space $C_c(\mathbb{R})$ of continuous functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = 0$ for $|x|$ sufficiently large.
5. Let $(V, \|\cdot\|)$ be a normed space, and let (\mathbf{v}_n) be a sequence in V .
 - (a) If V is complete, show that $\sum_{n=1}^{\infty} \mathbf{v}_n$ converges whenever $\sum_{n=1}^{\infty} \|\mathbf{v}_n\|$ converges.
 - (b)* If $\sum_{n=1}^{\infty} \mathbf{v}_n$ converges whenever $\sum_{n=1}^{\infty} \|\mathbf{v}_n\|$ converges, show that V is complete. (Hint: If (\mathbf{v}_n) is Cauchy, there is a subsequence (\mathbf{v}_{n_i}) such that $\sum_{i=1}^{\infty} \|\mathbf{v}_{n_{i+1}} - \mathbf{v}_{n_i}\|$ converges.)
6. Which of the following functions $f : [0, \infty) \rightarrow \mathbb{R}$ are uniformly continuous?
 - (a) $f(x) = \sin x^2$
 - (b) $f(x) = \inf\{|x - n^2| \mid n \in \mathbb{Z}\}$
 - (c) $(\sin x^3)/(x + 1)$
7. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable, and suppose that f' is bounded. Show that f is uniformly continuous. Let $g : [-1, 1] \rightarrow \mathbb{R}$ be given by $g(x) = x^2 \sin(1/x^2)$ for $x \neq 0$ and $g(0) = 0$. Show that g is differentiable, but that its derivative is unbounded. Is g uniformly continuous?
8. Suppose that $f : [0, \infty) \rightarrow \mathbb{R}$ is continuous, and that $f(x)$ tends to a finite limit as $x \rightarrow \infty$. Must f be uniformly continuous on $[0, \infty)$? Give a proof or a counterexample.

9. Let ℓ^1 be the set of real sequences (x_n) such that $\sum_{n=1}^{\infty} |x_n|$ is convergent. Show that with addition and scalar multiplication defined termwise, ℓ^1 is a vector space. Define $\|\cdot\|_1 : \ell^1 \rightarrow \mathbb{R}$ by $\|(x_n)\|_1 = \sum_{n=1}^{\infty} |x_n|$. Show that $\|\cdot\|_1$ is a norm on ℓ^1 , and that $(\ell^1, \|\cdot\|_1)$ is complete. Is $\overline{B}_1(0, \|\cdot\|_1)$ sequentially compact?
10. Let ℓ^∞ be the vector space of bounded real sequences $\mathbf{x} = (x_n)$, with the norm given by $\|(x_n)\| = \sup |x_n|$. Let (\mathbf{x}_k) be a bounded sequence in ℓ^∞ , and let $x_{i,k} \in \mathbb{R}$ be the i th entry in the sequence \mathbf{x}_k . Show that (\mathbf{x}_k) has a subsequence (\mathbf{x}_{k_j}) such that for each fixed value of i , the sequence (x_{i,k_j}) converges as $j \rightarrow \infty$. (Hint: use a diagonal argument.) Must \mathbf{x}_{k_j} converge in ℓ^∞ ? How is this example related to the notions of pointwise and uniform convergence?
11. Show that $A = \{(x_n) \in \ell^1 \mid |x_n| \leq 1/n^2 \text{ for all } n\}$ is a sequentially compact subset of $(\ell^1, \|\cdot\|_1)$, but that $B = \{(x_n) \in \ell^1 \mid |x_n| \leq 1/n \text{ for all } n\}$ is not.
12. Let V be a complete normed space, and let $\mathcal{B}(V, V)$ be the space of continuous linear maps from V to itself. Show that $\mathcal{B}(V, V)$ is complete with respect to the operator norm defined in problem 6 of the first example sheet. Show that if $\phi \in \mathcal{B}(V, V)$ satisfies $\|\phi\| < 1$, then $I - \phi$ is invertible, where I is the identity map. (Hint: consider $I + \phi + \phi^2 + \dots$) Deduce that the set of invertible maps is an open subset of $\mathcal{B}(V, V)$.
13. Suppose V is a normed vector space, and let $C_V = \{\mathbf{v}_n \mid \mathbf{v}_n \in V, (\mathbf{v}_n) \text{ is Cauchy}\}$. The operations $\lambda(\mathbf{v}_n) = (\lambda\mathbf{v}_n)$ and $(\mathbf{v}_n) + (\mathbf{w}_n) = (\mathbf{v}_n + \mathbf{w}_n)$ make C_V into a vector space.
- If $(\mathbf{v}_n) \in C_V$, show that $\lim_{n \rightarrow \infty} \|\mathbf{v}_n\|$ exists.
 - Define $\|(\mathbf{v}_n)\| = \lim_{n \rightarrow \infty} \|\mathbf{v}_n\|$, and let $C_0 = \{(\mathbf{v}_n) \mid \|(\mathbf{v}_n)\| = 0\}$. Show that C_0 is a linear subspace of C_V .
 - Show that $\overline{V} = C_V/C_0$ is a complete normed space containing V as a dense linear subspace. If W is another normed space satisfying these properties, show that $W \simeq \overline{V}$.
- 14.* (*A space-filling curve*) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function with the following properties: $f(t) = 0$ for $t \in [0, 1/3]$; $f(t) = 1$ for $t \in [2/3, 1]$; $0 \leq f(t) \leq 1$ for all $t \in \mathbb{R}$; and $f(t+2) = f(t)$ for all $t \in \mathbb{R}$. Define $\Phi(t) = (x(t), y(t))$, where

$$x(t) = \sum_{n=1}^{\infty} 2^{-n} f(3^{2n-1}t), \quad y(t) = \sum_{n=1}^{\infty} 2^{-n} f(3^{2n}t).$$

Prove that Φ is continuous and maps the unit interval $I = [0, 1]$ onto the unit square $I^2 \subset \mathbb{R}^2$. (Hint: show that $\Phi(I)$ contains all points of the form $(a/2^n, b/2^n)$, where $a, b \in \mathbb{Z}$, $0 \leq a, b \leq 2^n$.)

- 15.* Suppose C is a sequentially compact subset of a normed space V . If $\{U_\alpha \mid \alpha \in A\}$ is a family of open subsets of V , we say that the U_α cover C if $C \subset \cup_{\alpha \in A} U_\alpha$.
- Given $\epsilon > 0$, show that there is a finite set of points $\{\mathbf{v}_1, \dots, \mathbf{v}_n\} \subset C$ such that $\{B_\epsilon(\mathbf{v}_1), \dots, B_\epsilon(\mathbf{v}_n)\}$ cover C .
 - Show that there exists some $\epsilon > 0$ such that for every $\mathbf{v} \in C$, $B_\epsilon(\mathbf{v}) \subset U_\alpha$ for some α . Deduce that there is a finite set $\{\alpha_1, \dots, \alpha_n\} \subset A$ such that $\{U_{\alpha_1}, \dots, U_{\alpha_n}\}$ cover C .

J.Rasmussen@dpmms.cam.ac.uk