

1. Which of the following subsets of \mathbb{R}^2 are (a) connected, (b) path-connected?
 - (i) $D_1((-1, 0)) \cup D_1((1, 0))$ (ii) $D_1((-1, 0)) \cup B_1((1, 0))$
 - (iii) $\{(x, y) : x = 0 \text{ or } y/x \in \mathbb{Q}\}$ (iv) $\{(x, y) : x = 0 \text{ or } y/x \in \mathbb{Q}\} \setminus \{(0, 0)\}$.
2. Let $f: X \rightarrow S$ be a function from a connected space X to a set S . Assume f is *locally constant*: every $x \in X$ has a neighbourhood on which f is constant. Show that f is constant.
3. Show that homeomorphic spaces have the same number of connected components. Show that no two of $[0, 1]$, $[0, 1)$ and $(0, 1)$ are homeomorphic. Show also that the letters A and H drawn in the plane are not homeomorphic.
4. Find the connected components of the subspace $X = \{(0, 0), (0, 1)\} \cup \bigcup_{n \in \mathbb{N}} \{1/n\} \times [0, 1]$ of \mathbb{R}^2 . Show that there exist $x, y \in X$ that belong to different components but there are no open sets U and V disconnecting X with $x \in U$ and $y \in V$.
5. Let $A \subset \mathbb{R}^n$ be such that every continuous function $f: A \rightarrow \mathbb{R}$ is bounded. Show that A is compact.
6. Show that if A and B are closed subsets of \mathbb{R}^n and if A or B is bounded, then $A + B$ is closed. Give an example in \mathbb{R} to show that the boundedness condition cannot be omitted.
7. Let R be the equivalence relation on $Q = [0, 1]^2$ defined as follows: $(x_1, y_1) \sim (x_2, y_2)$ if and only if EITHER $(x_1, y_1) = (x_2, y_2)$ OR $\{x_1, x_2\} = \{0, 1\}$ and $y_1 = y_2$ OR $y_1 = y_2 = 0$ OR $y_1 = y_2 = 1$. Show that Q/R is homeomorphic to the sphere $S^2 = \{x \in \mathbb{R}^3 : \|x\| = 1\}$.
8. Show that a continuous real-valued function on a sequentially compact topological space is bounded and attains its bounds. Show also that a continuous function from a compact metric space to an arbitrary metric space is uniformly continuous.
9. (a) Let R_1 be an equivalence relation on a topological space X and let R_2 be an equivalence relation on the quotient space X/R_1 . Define

$$R = \{(x, y) \in X \times X : (q(x), q(y)) \in R_2\}$$

where $q: X \rightarrow X/R_1$ is the quotient map. Show that R is an equivalence relation on X and that X/R is homeomorphic to $(X/R_1)/R_2$.

(b) For a topological space X and for $A \subset X$, we let X/A denote the quotient space of X by the relation identifying the points of A : $x \sim y$ if and only if either $x = y$ or $x, y \in A$. Now consider the subset $A = \{(0, 0, 1), (0, 0, -1)\}$ of the two-dimensional sphere S^2 , and the subset $B = \{(2 + \cos \theta, 0, \sin \theta) : \theta \in [0, 2\pi]\}$ of T^2 (as defined in lectures). Show that S^2/A and T^2/B are homeomorphic.

10. (a) Show that the coordinate projections π_X and π_Y on a product space $X \times Y$ are open maps. Show that if Y is compact, then π_X is a *closed map*: for a closed subset F of $X \times Y$, its projection $\pi_X(F)$ is closed in X . Give an example of a closed set in \mathbb{R}^2 whose projections are not closed in \mathbb{R} .

(b) Let $f: X \rightarrow Y$ be a function between topological spaces. The *graph of f* is the set $\Gamma = \{(x, y) \in X \times Y : y = f(x)\}$. Show that if f is continuous and Y is Hausdorff, then Γ is closed in the product topology. Conversely, show that if Γ is closed and Y is compact, then f is continuous.

11. Let M be a non-empty compact metric space and $f: M \rightarrow M$ be a function.

(a) Show that if $d(f(x), f(y)) < d(x, y)$ for all $x \neq y$ in M , then f has a unique fixed point.

(b) Show that if f is isometric, *i.e.*, $d(f(x), f(y)) = d(x, y)$ for all $x, y \in M$, then f is surjective.

12. (a) A topological space is *normal* if disjoint closed subsets can be separated by open sets: given disjoint closed subsets A and B , there exist disjoint open sets U and V such that $A \subset U$ and $B \subset V$. Show that a compact Hausdorff space is normal.

(b) Let (C_n) be a decreasing sequence of compact connected subsets of a Hausdorff space. Show that $\bigcap_{n \in \mathbb{N}} C_n$ is connected. (Part (a) will be useful here.) Give an example in \mathbb{R}^2 of a decreasing sequence of closed connected sets whose intersection is disconnected.

13. Show that $C[0, 1]$ in the uniform metric D is separable. Let $B = \{f \in C[0, 1] : D(0, f) \leq 1\}$ and $B' = \{f \in B : f \text{ differentiable and } f' \in B\}$. Show that B is closed but not compact. On the other hand, show that every sequence in B' has a subsequence convergent in $C[0, 1]$. Deduce that $\overline{B'}$ is compact.

14. Given topological spaces X and Y and continuous bijections $f: X \rightarrow Y$ and $g: Y \rightarrow X$, show that X and Y need not be homeomorphic.

+15. Let $f: \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a function under which the image of any path-connected set is path-connected and the image of any compact set is compact. Show that f must be continuous.