

- 1 Use the residue theorem to give a proof of Cauchy's derivative formula: if f is holomorphic on $D(a, R)$, and $|w - a| < r < R$, then

$$f^{(n)}(w) = \frac{n!}{2\pi i} \int_{|z-a|=r} \frac{f(z)}{(z-w)^{n+1}} dz.$$

- 2 Show that if f and g are holomorphic and non-constant, then $\deg_{z=a}(g \circ f) = \deg_{z=a}(f) \cdot \deg_{z=f(a)}(g)$.

- 3 Use Rouché's Theorem to give another proof of the Fundamental Theorem of Algebra.

- 4 Let $p(z) = z^5 + z$. Find all z such that $|z| = 1$ and $\operatorname{Im} p(z) = 0$. Calculate $\operatorname{Re} p(z)$ for such z . Hence sketch the curve $p \circ \gamma$, where $\gamma(t) = e^{2\pi i t}$ and use your sketch to determine the number of z (counted with multiplicity) such that $|z| < 1$ and $p(z) = x$ for each real number x .

- 5 Evaluate:

(a) $\int_0^\pi \frac{d\theta}{4 + \sin^2 \theta}$;

(b) $\int_{-\infty}^\infty \frac{\sin mx}{x(a^2 + x^2)} dx$ where $a, m \in \mathbb{R}^+$;

(c) $\int_0^\infty \frac{x^2}{(x^2 + 4)^2(x^2 + 9)} dx$;

(d) $\int_0^{2\pi} \frac{\cos^3 3t}{1 - 2a \cos t + a^2} dt$ where $a \in (0, 1)$.

- 6 By integrating $z/(a - e^{-iz})$ round the rectangle with vertices $\pm\pi, \pm\pi + iR$, prove that

$$\int_0^\pi \frac{x \sin x}{1 - 2a \cos x + a^2} dx = \frac{\pi}{a} \log(1 + a) \quad \text{for } a \in (0, 1).$$

- 7 Evaluate:

(a) $\int_0^\infty \sin x^2 dx$

(b) $\int_{-\infty}^\infty e^{-ax^2} e^{-itx} dx$ where $a > 0, t \in \mathbb{R}$

(c) $\int_0^\infty \frac{\ln(x^2 + 1)}{x^2 + 1} dx$

(d) $\int_0^\infty \frac{\cosh ax}{\cosh x} dx, a \in (-1, 1)$

(e) $\int_{-\infty}^\infty \frac{\sin x}{x} e^{-itx} dx, t \in \mathbb{R}$.

- 8 Assuming $\alpha \geq 0$ and $\beta \geq 0$ prove that

$$\int_0^\infty \frac{\cos \alpha x - \cos \beta x}{x^2} dx = \frac{\pi}{2}(\beta - \alpha),$$

and deduce the value of

$$\int_0^\infty \left(\frac{\sin x}{x} \right)^2 dx.$$

9 (i) For a positive integer N , let γ_N be the square contour with vertices $(\pm 1 \pm i)(N + 1/2)$. Show that there exists $C > 0$ such that for every N , $|\cot \pi z| < C$ on γ_N .

(ii) By integrating $\frac{\pi \cot \pi z}{z^2 + 1}$ around γ_N , show that

$$\sum_{n=0}^{\infty} \frac{1}{n^2 + 1} = \frac{1 + \pi \coth \pi}{2}.$$

(iii) Evaluate $\sum_{n=0}^{\infty} (-1)^n / (n^2 + 1)$.

10 (i) Show that the Taylor expansion of $z/(e^z - 1)$ near the origin has the form

$$1 - \frac{z}{2} + \sum_{k=1}^{\infty} \frac{(-1)^{k-1} B_k}{(2k)!} z^{2k}$$

where the numbers B_k (the *Bernoulli numbers*) are rational.

(ii) If k is a positive integer show that

$$\sum_{n=1}^{\infty} \frac{1}{n^{2k}} = \frac{2^{2k-1} \pi^{2k} B_k}{(2k)!}.$$

11 Prove that $z^5 + 2 + e^z$ has exactly three zeros in the half-plane $\{z \mid \operatorname{Re}(z) < 0\}$.

12 Show that $z^4 + 26z + 2 = 0$ has exactly three zeroes with $5/2 < |z| < 3$.

13 Show that the equation $z^4 + z + 1 = 0$ has one solution in each quadrant. Prove that all solutions lie inside the circle $\{z \mid |z| = 3/2\}$.

14 Show that the equation $z \sin z = 1$ has only real solutions.

15 Suppose f is holomorphic when $|z| \leq 1$ and satisfies $|f(z)| < 1$ when $|z| = 1$. Show there is exactly one complex number w such that $|w| < 1$ and $f(w) = w$.

16 Let f be a meromorphic function on \mathbb{C} such that $|f(z)| \rightarrow \infty$ as $|z| \rightarrow \infty$. Show that f cannot have poles at all integer points.