

### Complex Analysis IB – 2013 – Sheet 3

The symbols  $\Re(z)$  and  $\Im(z)$  denote the real respectively imaginary parts of  $z$ .

1. Let  $f$  be a meromorphic function on  $\mathbb{C}$  for which  $|f(z)| \rightarrow \infty$  as  $|z| \rightarrow \infty$ . Show that  $f$  cannot have poles at all integer points.
2. Let  $g(z) = p(z)/q(z)$  be a rational function with  $\deg(q) \geq \deg(p) + 2$ . Show that the sum of the residues of  $g$  over all its singularities is zero.
3. Prove that the group of conformal automorphisms of the Riemann sphere  $\mathbb{C} \cup \{\infty\} = \mathbb{CP}^1$  is the Möbius group. [*Hint: take an automorphism  $g$  fixing 0 and  $\infty$  and consider  $z \mapsto g(z)/z$ .]*
4. Evaluate the following:

$$\begin{array}{ll} (a) \int_0^\pi \frac{d\theta}{4 + \sin^2 \theta}; & (b) \int_0^\infty \frac{x^2 dx}{(x^2 + 4)^2(x^2 + 9)}; \\ (c) \int_0^\infty \sin x^2 dx; & (d) \int_0^\infty \frac{\ln(x^2 + 1)}{x^2 + 1} dx. \end{array}$$

5. For  $-1 < \alpha < 1$  and  $\alpha \neq 0$ , compute

$$\int_0^\infty \frac{x^\alpha}{1 + x + x^2} dx.$$

Letting  $\alpha \rightarrow 0$ , compute

$$\int_0^\infty \frac{1}{1 + x + x^2} dx.$$

6. Establish the following refinement of the Fundamental Theorem of Algebra. Let  $p(z) = z^n + a_{n-1}z^{n-1} + \dots + a_0$  be a polynomial of degree  $n$ , and let  $A = \max\{|a_i|, 0 \leq i \leq n-1\}$ . Then  $p(z)$  has  $n$  roots (counted with multiplicity) in the disk  $\{|z| < A + 1\}$ .
7. Let  $p(z) = z^5 + z$ . Find all  $z$  such that  $|z| = 1$  and  $\Im p(z) = 0$ . Calculate  $\Re p(z)$  for such  $z$ . Sketch the curve  $p \circ \gamma$ , where  $\gamma(t) = e^{2\pi it}$ , and hence determine the number of  $z$  (counted with multiplicity) such that  $|z| < 1$  and  $p(z) = x$  for each  $x \in \mathbb{R}$ .
8. (i) For a positive integer  $N$ , let  $\gamma_N$  be the square contour with vertices  $(\pm 1 \pm i)(N + 1/2)$ . Show that there exists  $C > 0$  such that for every  $N$ ,  $|\cot \pi z| < C$  on  $\gamma_N$ .  
(ii) By integrating  $\frac{\pi \cot \pi z}{z^2 + 1}$  around  $\gamma_N$ , show that

$$\sum_{n=0}^{\infty} \frac{1}{n^2 + 1} = \frac{1 + \pi \coth \pi}{2}.$$

- (iii) Evaluate  $\sum_{n=0}^{\infty} (-1)^n / (n^2 + 1)$ .

9. Show that the Taylor expansion of  $z/(e^z - 1)$  near the origin has the form

$$1 - \frac{z}{2} + \sum_{k=1}^{\infty} \frac{(-1)^{k-1} B_k}{(2k)!} z^{2k}$$

where the numbers  $B_k$  (the *Bernoulli numbers*) are rational.

10. (i) Show that  $z^4 + 12z + 1$  has exactly three zeroes in the annulus  $\{1 < |z| < 4\}$ .  
(ii) Prove that  $z^5 + 2 + e^z$  has exactly three zeroes in the half-plane  $\{z \mid \Re(z) < 0\}$ .  
(iii) Show that the equation  $z^4 + z + 1 = 0$  has one solution in each quadrant. Prove that all solutions lie inside the circle  $\{z \mid |z| = 3/2\}$ .
11. Show that the equation  $z \sin z = 1$  has only real solutions.  
[Hint: Find the number of real roots in the interval  $[-(n + 1/2)\pi, (n + 1/2)\pi]$  and compare with the number of zeroes of  $z \sin z - 1$  in a square box  $\{|\Re(z)|, |\Im(z)| < (n + 1/2)\pi\}$ .]
- 11\* (Additional) Let  $f : U \rightarrow \mathbb{C}$  be holomorphic and suppose  $a \in U$  with  $f'(a) \neq 0$ . Show that for  $r > 0$  sufficiently small,

$$g(w) = \frac{1}{2\pi i} \int_{|z-a|=r} \frac{z f'(z)}{f(z) - w} dz$$

defines a holomorphic function  $g$  in a neighbourhood of  $f(a)$  which is inverse to  $f$ .

The following integrals are *not* part of the question sheet, but may provide a good start for revision or a first port of call for the addicted.

- (i)  $\int_{-\infty}^{\infty} \frac{\sin mx}{x(a^2 + x^2)} dx$  where  $a, m \in \mathbb{R}^+$ ;      (ii)  $\int_0^{2\pi} \frac{\cos^3 3t}{1 - 2a \cos t + a^2} dt$  where  $a \in (0, 1)$ ;  
(iii)  $\int_{-\infty}^{\infty} e^{-ax^2} e^{-itx} dx$  where  $a > 0, t \in \mathbb{R}$ ;      (iv)  $\int_{-\infty}^{\infty} \frac{\sin x}{x} e^{-itx} dx$ , where  $t \in \mathbb{R}$ .

(v) By integrating  $z/(a - e^{-iz})$  round the rectangle with vertices  $\pm\pi, \pm\pi + iR$ , prove that

$$\int_0^{\pi} \frac{x \sin x}{1 - 2a \cos x + a^2} dx = \frac{\pi}{a} \log(1 + a) \quad \text{for } a \in (0, 1).$$

(vi) Assuming  $\alpha \geq 0$  and  $\beta \geq 0$  prove that

$$\int_0^{\infty} \frac{\cos \alpha x - \cos \beta x}{x^2} dx = \frac{\pi}{2}(\beta - \alpha),$$

and deduce the value of

$$\int_0^{\infty} \left(\frac{\sin x}{x}\right)^2 dx.$$

Ivan Smith  
is200@cam.ac.uk