

Complex Analysis IB – 2014 – Sheet 1

- Let $T : \mathbb{C} = \mathbb{R}^2 \rightarrow \mathbb{R}^2 = \mathbb{C}$ be a real linear map. Show that T can be written $Tz = Az + B\bar{z}$ for unique $A, B \in \mathbb{C}$. Show that T is complex differentiable if and only if $B = 0$.
- (i) Let $f : D \rightarrow \mathbb{C}$ be a holomorphic function defined on a domain (path-connected open subset) D . Show that f is constant if any of its real part, modulus or argument is constant.
 (ii) Find all holomorphic functions on \mathbb{C} of the form $f(x + iy) = u(x) + iv(y)$ where u and v are both real valued.
 (iii) Find all holomorphic functions $f(z)$ on \mathbb{C} which have real part $x^3 - 3xy^2$.
- Define $f : \mathbb{C} \rightarrow \mathbb{C}$ by $f(0) = 0$, and

$$f(z) = \frac{(1+i)x^3 - (1-i)y^3}{x^2 + y^2} \quad \text{for } z = x + iy \neq 0.$$

Show that f satisfies the Cauchy-Riemann equations at 0 but is not differentiable there.

- (i) Verify directly that e^z and $\cos z$ satisfy the Cauchy-Riemann equations everywhere.
 (ii) Find the set of $z \in \mathbb{C}$ for which $|e^{iz}| > 1$, and the set of those for which $|e^z| \leq e^{|z|}$.
 (iii) Find the zeros of $1 + e^z$ and $\cosh z$.
- (i) Defining $z^\alpha = e^{\alpha \text{Log } z}$, for Log the principal branch of the logarithm and $z \notin \mathbb{R}_{\leq 0}$, show that $\frac{d}{dz} z^\alpha = \alpha z^{\alpha-1}$. Does $(zw)^\alpha = z^\alpha w^\alpha$ always hold?
 (ii) If $z \in \mathbb{C}$, show that $n \text{Log}(1 + z/n)$ is defined if n is sufficiently large, and that it tends to z as n tends to ∞ . Deduce

$$\lim_{n \rightarrow \infty} \left(1 + \frac{z}{n}\right)^n = e^z \quad \forall z \in \mathbb{C}.$$

- Find conformal equivalences between the following pairs of domains:
 - the sector $\{z \in \mathbb{C} \mid -\pi/4 < \arg(z) < \pi/4\}$ and the open unit disc D ;
 - the lune $\{z \in \mathbb{C} : |z - 1| < \sqrt{2} \text{ and } |z + 1| < \sqrt{2}\}$ and the open unit disk D ;
 - the strip $S = \{z \in \mathbb{C} : 0 < \text{Im } z < 1\}$ and the quadrant $Q = \{z \in \mathbb{C} : \text{Re } z > 0, \text{Im } z > 0\}$.
 By considering a suitable bounded solution of the Laplace equation $u_{xx} + u_{yy} = 0$ on the strip S , find a non-constant harmonic function on Q which is constant on each of the two boundaries of the quadrant.
- (i) Show that the general Möbius transformation which takes the unit disk to itself has the form $z \mapsto \lambda \frac{z-a}{\bar{a}z-1}$, with $|a| < 1$, $|\lambda| = 1$. [*Hint: first show these maps form a group.*]
 (ii) Find a Möbius transformation taking the region between $\{|z| = 1\}$ and $\{|z - 1| = 5/2\}$ to an annulus $\{1 < |z| < R\}$. [*Hint: A circle can be described by an equation of the shape $|z - a|/|z - b| = l$.*]
 (iii) Find a conformal map from an infinite strip onto an annulus. Can such a map be the restriction to the strip of a Möbius map?

8. Prove that each of the following series converges uniformly on the corresponding subset of \mathbb{C} :

$$(a) \sum_{n=1}^{\infty} \sqrt{n} e^{-nz}, \quad \text{on } \{z \mid 0 < r \leq \operatorname{Re}(z)\}; \quad (b) \sum_{n=1}^{\infty} \frac{2^n}{z^n + z^{-n}}, \quad \text{on } \{z \mid |z| \leq r < \frac{1}{2}\}.$$

9. Calculate $\int_{\gamma} z \sin z \, dz$ when γ is the straight line joining 0 to i .

10. Show that the following functions do not have antiderivatives (i.e. functions of which they are the derivative) on the domains indicated:

$$(a) \frac{1}{z} - \frac{1}{z-1} \quad (0 < |z| < 1); \quad (b) \frac{z}{1+z^2} \quad (1 < |z| < \infty).$$

Ivan Smith
is200@cam.ac.uk