

Complex Analysis IB – 2014 – Sheet 3

- Let f be a meromorphic function on \mathbb{C} for which $|f(z)| \rightarrow \infty$ as $|z| \rightarrow \infty$. Show that f cannot have poles at all integer points.
- Let $g(z) = p(z)/q(z)$ be a rational function with $\deg(q) \geq \deg(p) + 2$. Show that the sum of the residues of g over all its singularities is zero.
- Prove that the group of conformal automorphisms of the Riemann sphere $\mathbb{C} \cup \{\infty\} = \mathbb{C}\mathbb{P}^1$ is the Möbius group. [Hint: take an automorphism g fixing 0 and ∞ and consider $z \mapsto g(z)/z$.]
- Evaluate the following:

$$(a) \int_0^\pi \frac{d\theta}{4 + \sin^2 \theta}; \quad (b) \int_0^\infty \frac{x^2 dx}{(x^2 + 4)^2(x^2 + 9)};$$

$$(c) \int_0^\infty \sin x^2 dx; \quad (d) \int_0^\infty \frac{\ln(x^2 + 1)}{x^2 + 1} dx.$$

- For $-1 < \alpha < 1$ and $\alpha \neq 0$, compute

$$\int_0^\infty \frac{x^\alpha}{1 + x + x^2} dx.$$

- Establish the following refinement of the Fundamental Theorem of Algebra. Let $p(z) = z^n + a_{n-1}z^{n-1} + \dots + a_0$ be a polynomial of degree n , and let $A = \max\{|a_i|, 0 \leq i \leq n-1\}$. Then $p(z)$ has n roots (counted with multiplicity) in the disk $\{|z| < A + 1\}$.
- Let $p(z) = z^5 + z$. Find all z such that $|z| = 1$ and $\operatorname{Im} p(z) = 0$. Calculate $\operatorname{Re} p(z)$ for such z . Sketch the curve $p \circ \gamma$, where $\gamma(t) = e^{2\pi it}$, and hence determine the number of z (counted with multiplicity) such that $|z| < 1$ and $p(z) = x$ for each $x \in \mathbb{R}$.
- (i) For a positive integer N , let γ_N be the square contour with vertices $(\pm 1 \pm i)(N + 1/2)$. Show that there exists $C > 0$ such that for every N , $|\cot \pi z| < C$ on γ_N .
 (ii) By integrating $\frac{\pi \cot \pi z}{z^2 + 1}$ around γ_N , show that

$$\sum_{n=0}^{\infty} \frac{1}{n^2 + 1} = \frac{1 + \pi \coth \pi}{2}.$$

- (iii) Evaluate $\sum_{n=0}^{\infty} (-1)^n / (n^2 + 1)$.

- Show that the Taylor expansion of $z/(e^z - 1)$ near the origin has the form

$$1 - \frac{z}{2} + \sum_{k=1}^{\infty} \frac{(-1)^{k-1} B_k}{(2k)!} z^{2k}$$

where the numbers B_k (the *Bernoulli numbers*) are rational.

10. Let $w \in \mathbb{C}$, and let $\gamma, \delta: [0, 1] \rightarrow \mathbb{C}$ be closed curves such that for all $t \in [0, 1]$, $|\gamma(t) - \delta(t)| < |\gamma(t) - w|$. By computing the winding number $n_\sigma(0)$ about the origin for the closed curve $\sigma(t) = (\delta(t) - w)/(\gamma(t) - w)$, show that $n_\gamma(w) = n_\delta(w)$.
- (ii) If $w \in \mathbb{C}$, $r > 0$, and γ is a closed curve which does not meet $B_w(r)$, show that $n_\gamma(w) = n_\gamma(z)$ for every $z \in B_w(r)$. Deduce that if γ is a closed curve in \mathbb{C} and U is the complement of (the image of) γ , then the function $w \mapsto n_\gamma(w)$ is a locally constant function on U .
11. (i) Show that $z^4 + 12z + 1$ has exactly three zeroes in the annulus $\{1 < |z| < 4\}$.
- (ii) Prove that $z^5 + 2 + e^z$ has exactly three zeros in the half-plane $\{z \mid \operatorname{Re}(z) < 0\}$.
- (iii) Show that the equation $z^4 + z + 1 = 0$ has one solution in each quadrant. Prove that all solutions lie inside the circle $\{z \mid |z| = 3/2\}$.
12. Show that the equation $z \sin z = 1$ has only real solutions.
- [Hint: Find the number of real roots in the interval $[-(n + 1/2)\pi, (n + 1/2)\pi]$ and compare with the number of zeroes of $z \sin z - 1$ in a square box $\{|\operatorname{Re}(z)|, |\operatorname{Im}(z)| < (n + 1/2)\pi\}$.]
- 13* (Additional) Let $f : U \rightarrow \mathbb{C}$ be holomorphic and suppose $a \in U$ with $f'(a) \neq 0$. Show that for $r > 0$ sufficiently small,

$$g(w) = \frac{1}{2\pi i} \int_{|z-a|=r} \frac{z f'(z)}{f(z) - w} dz$$

defines a holomorphic function g in a neighbourhood of $f(a)$ which is inverse to f .

The following integrals are *not* part of the question sheet, but may provide a good start for revision or a first port of call for the addicted.

- (i) $\int_{-\infty}^{\infty} \frac{\sin mx}{x(a^2 + x^2)} dx$ where $a, m \in \mathbb{R}^+$; (ii) $\int_0^{2\pi} \frac{\cos^3 3t}{1 - 2a \cos t + a^2} dt$ where $a \in (0, 1)$;
- (iii) $\int_{-1}^1 \frac{\sqrt{1-x^2}}{1+x^2} dx$ (“dog-bone” contour); (iv) $\int_{-\infty}^{\infty} \frac{\sin x}{x} e^{-itx} dx$, where $t \in \mathbb{R}$.

(v) By integrating $z/(a - e^{-iz})$ round the rectangle with vertices $\pm\pi, \pm\pi + iR$, prove that

$$\int_0^\pi \frac{x \sin x}{1 - 2a \cos x + a^2} dx = \frac{\pi}{a} \log(1 + a) \quad \text{for } a \in (0, 1).$$

(vi) Assuming $\alpha \geq 0$ and $\beta \geq 0$ prove that

$$\int_0^\infty \frac{\cos \alpha x - \cos \beta x}{x^2} dx = \frac{\pi}{2}(\beta - \alpha),$$

and deduce the value of

$$\int_0^\infty \left(\frac{\sin x}{x} \right)^2 dx.$$

Ivan Smith
is200@cam.ac.uk