

IB Groups, Rings and Modules: Example Sheet 4

All rings in this course are commutative with a multiplicative identity.

1. How many abelian groups are there of order 6? Of order 60? Of order 6000?
2. Let M be a module over an integral domain R . An element $m \in M$ is a *torsion* element if $rm = 0$ for some non-zero $r \in R$. Show that the set T of all torsion elements in M is a submodule of M , and that the quotient M/T is *torsion-free* – that is, contains no non-zero torsion elements.
3. We say that an R -module satisfies condition (N) on submodules if any submodule is finitely generated. Show that this condition is equivalent to condition (ACC) : every increasing chain of submodules terminates.
4. (i) Is the abelian group \mathbb{Q} torsion-free? Is it free? Is it finitely generated?
 (ii) What are the torsion elements in the abelian group \mathbb{Q}/\mathbb{Z} ? In \mathbb{R}/\mathbb{Z} ? In \mathbb{R}/\mathbb{Q} ?
 (iii) Prove that \mathbb{R} is not finitely generated as a module over the ring \mathbb{Q} .
5. Let M be a module over a ring R , and let N be a submodule of M .
 - (i) Show that if M is finitely generated then so is M/N .
 - (ii) Show that if N and M/N are finitely generated then so is M .
 - (iii) Show that if M/N is free, then $M \cong N \oplus M/N$.
 - (iv) Show that if R is a PID and M is a finitely-generated free module, then N is free.
6. Use elementary operations to bring the integer matrix $A = \begin{pmatrix} -4 & -6 & 7 \\ 2 & 2 & 4 \\ 6 & 6 & 15 \end{pmatrix}$ to Smith normal form D .
 Check your result using minors. Explain how to find invertible matrices P, Q for which $D = QAP$.
7. Work out the invariant factors of the matrices over $\mathbb{R}[X]$:

$$\begin{pmatrix} 2X-1 & X & X-1 & 1 \\ X & 0 & 1 & 0 \\ 0 & 1 & X & X \\ 1 & X^2 & 0 & 2X-2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} X^2+2X & 0 & 0 & 0 \\ 0 & X^2+3X+2 & 0 & 0 \\ 0 & 0 & X^3+2X^2 & 0 \\ 0 & 0 & 0 & X^4+X^3 \end{pmatrix}.$$

8. Let G be the abelian group with generators a, b, c , and relations $6a+10b=0$, $6a+15c=0$, $10b+15c=0$. (That is, G is the free abelian group on generators a, b, c quotiented by the subgroup generated by the elements $6a+10b$, $6a+15c$, $10b+15c$). Determine the structure of G as a direct sum of cyclic groups.
9. Prove that a finitely-generated abelian group G is finite if and only if $G/pG = 0$ for some prime p .
 Give a non-trivial abelian group G such that $G/pG = 0$ for all primes p , and prove that your example is not finitely generated.
10. Let A be a complex matrix with characteristic polynomial $(X+1)^6(X-2)^3$ and minimal polynomial $(X+1)^3(X-2)^2$. Write down the possible Jordan normal forms for A .
11. Find a 2×2 matrix over $\mathbb{Z}[X]$ that is not equivalent to a diagonal matrix.
12. Let M be a finitely-generated module over a ring R , and let f be an R -module homomorphism from M to itself. Does f injective imply f surjective? Does f surjective imply f injective?

Additional Questions

13. Write $f(n)$ for the number of distinct abelian groups of order n .
- (i) Show that if $n = p_1^{a_1} p_2^{a_2} \cdots p_k^{a_k}$ with the p_i distinct primes and $a_i \in \mathbb{N}$ then $f(n) = f(p_1^{a_1}) \cdots f(p_k^{a_k})$.
 - (ii) Show that $f(p^a)$ equals the number $p(a)$ of partitions of a , that is, $p(a)$ is the number of ways of writing a as a sum of positive integers, where the order of summands is unimportant. (For example, $p(5) = 7$, since $5 = 4 + 1 = 3 + 2 = 3 + 1 + 1 = 2 + 2 + 1 = 2 + 1 + 1 + 1 = 1 + 1 + 1 + 1 + 1$.)
14. A real $n \times n$ matrix A satisfies the equation $A^2 + I = 0$. Show that n is even and A is similar to a block matrix $\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ with each block an $m \times m$ matrix (where $n = 2m$).
15. Let R be a Noetherian ring and M be a finitely generated R -module. Show that all submodules of M are finitely generated.
16. Show that a complex number α is an algebraic integer if and only if the additive group of the ring $\mathbb{Z}[\alpha]$ is finitely generated (i.e. $\mathbb{Z}[\alpha]$ is a finitely generated \mathbb{Z} -module). Furthermore if α and β are algebraic integers show that the subring $\mathbb{Z}[\alpha, \beta]$ of \mathbb{C} generated by α and β also has a finitely generated additive group and deduce that $\alpha - \beta$ and $\alpha\beta$ are algebraic integers. Show that the algebraic integers form a subring of \mathbb{C} .
17. What is the rational canonical form of a matrix?

Show that the group $GL_2(\mathbb{F}_2)$ of non-singular 2×2 matrices over the field \mathbb{F}_2 of 2 elements has three conjugacy classes of elements.

Show that the group $GL_3(\mathbb{F}_2)$ of non-singular 3×3 matrices over the field \mathbb{F}_2 has six conjugacy classes of elements, corresponding to minimal polynomials $X + 1$, $(X + 1)^2$, $(X + 1)^3$, $X^3 + 1$, $X^3 + X^2 + 1$, $X^3 + X + 1$, one each of elements of orders 1, 2, 3 and 4, and two of elements of order 7.

Comments and corrections should be sent to rdc26@dpmmms.cam.ac.uk.