

Linear Algebra: Example Sheet 1 of 4

The first twelve questions cover the relevant part of the course and should ensure a good understanding. The remaining questions may or may not be harder; they should only be attempted after completion of the first part.

1. Let $\mathbb{R}^{\mathbb{R}}$ be the vector space of all functions $f : \mathbb{R} \rightarrow \mathbb{R}$, with addition and scalar multiplication defined pointwise. Which of the following sets of functions form a vector subspace of $\mathbb{R}^{\mathbb{R}}$?
 - (a) The set C of continuous functions.
 - (b) The set $\{f \in C : |f(t)| \leq 1 \text{ for all } t \in [0, 1]\}$.
 - (c) The set $\{f \in C : f(t) \rightarrow 0 \text{ as } t \rightarrow \infty\}$.
 - (d) The set $\{f \in C : f(t) \rightarrow 1 \text{ as } t \rightarrow \infty\}$.
 - (e) The set of solutions of the differential equation $\ddot{x}(t) + (t^2 - 3)\dot{x}(t) + t^4x(t) = 0$.
 - (f) The set of solutions of $\ddot{x}(t) + (t^2 - 3)\dot{x}(t) + t^4x(t) = \sin t$.
 - (g) The set of solutions of $(\dot{x}(t))^2 - x(t) = 0$.
 - (h) The set of solutions of $(\ddot{x}(t))^4 + (x(t))^2 = 0$.

2. Suppose that the vectors $\mathbf{e}_1, \dots, \mathbf{e}_n$ form a basis for V . Which of the following are also bases?
 - (a) $\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2 + \mathbf{e}_3, \dots, \mathbf{e}_{n-1} + \mathbf{e}_n, \mathbf{e}_n$;
 - (b) $\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2 + \mathbf{e}_3, \dots, \mathbf{e}_{n-1} + \mathbf{e}_n, \mathbf{e}_n + \mathbf{e}_1$;
 - (c) $\mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2 - \mathbf{e}_3, \dots, \mathbf{e}_{n-1} - \mathbf{e}_n, \mathbf{e}_n - \mathbf{e}_1$;
 - (d) $\mathbf{e}_1 - \mathbf{e}_n, \mathbf{e}_2 + \mathbf{e}_{n-1}, \dots, \mathbf{e}_n + (-1)^n \mathbf{e}_1$.

3. Let V be a vector space over a field F .
 - (i) Describe a procedure for picking vectors in V that produces **either** a finite basis for V **or** an infinite linearly independent subset of V .
 - (ii) Show that V is finite dimensional if and only if every linearly independent subset $S \subset V$ is finite.
 - (iii) Deduce that a subspace of a finite dimensional vector space is always finite dimensional.
[Although it is true that every vector space V has a basis, this is only proved in lectures for V finite dimensional. It would not be reasonable to quote the more general result in answering this question.]

4. Let T, U and W be subspaces of V .
 - (i) Show that $T \cup U$ is a subspace of V only if either $T \leq U$ or $U \leq T$.
 - (ii) Give explicit counter-examples to the following statements:
 - (a) $T + (U \cap W) = (T + U) \cap (T + W)$;
 - (b) $(T + U) \cap W = (T \cap W) + (U \cap W)$.
 - (iii) Show that each of the equalities in (ii) can be replaced by a valid inclusion of one side in the other.

5. For each of the following pairs of vector spaces (V, W) over \mathbb{R} , either give an isomorphism $V \rightarrow W$ or show that no such isomorphism can exist. *[Here P denotes the space of polynomial functions $\mathbb{R} \rightarrow \mathbb{R}$, and $C[a, b]$ denotes the space of continuous functions defined on the closed interval $[a, b]$.]*
 - (a) $V = \mathbb{R}^4, W = \{\mathbf{x} \in \mathbb{R}^5 : x_1 + x_2 + x_3 + x_4 + x_5 = 0\}$.
 - (b) $V = \mathbb{R}^5, W = \{p \in P : \deg p \leq 5\}$.
 - (c) $V = C[0, 1], W = C[-1, 1]$.
 - (d) $V = C[0, 1], W = \{f \in C[0, 1] : f(0) = 0, f \text{ continuously differentiable}\}$.
 - (e) $V = \mathbb{R}^2, W = \{\text{solutions of } \ddot{x}(t) + x(t) = 0\}$.
 - (f) $V = \mathbb{R}^4, W = C[0, 1]$.
 - (g) (Harder:) $V = P, W = \mathbb{R}^{\mathbb{N}}$.

6. (i) If α and β are linear maps from U to V show that $\alpha + \beta$ is linear. Give explicit counter-examples to the following statements:
 - (a) $\text{Im}(\alpha + \beta) = \text{Im}(\alpha) + \text{Im}(\beta)$;
 - (b) $\text{Ker}(\alpha + \beta) = \text{Ker}(\alpha) \cap \text{Ker}(\beta)$.

Show that each of these equalities can be replaced by a valid inclusion of one side in the other.

(ii) Let α be a linear map from V to V . Show that if $\alpha^2 = \alpha$ then $V = \text{Ker}(\alpha) \oplus \text{Im}(\alpha)$. Does your proof work if V is infinite dimensional? Is the result still true?

7. Let

$$U = \{\mathbf{x} \in \mathbb{R}^5 : x_1 + x_3 + x_4 = 0, 2x_1 + 2x_2 + x_5 = 0\}, \quad W = \{\mathbf{x} \in \mathbb{R}^5 : x_1 + x_5 = 0, x_2 = x_3 = x_4\}.$$

Find bases for U and W containing a basis for $U \cap W$ as a subset. Give a basis for $U + W$ and show that

$$U + W = \{\mathbf{x} \in \mathbb{R}^5 : x_1 + 2x_2 + x_5 = x_3 + x_4\}.$$

8. Recall that F^n has standard basis $\mathbf{e}_1, \dots, \mathbf{e}_n$. Let U be a subspace of F^n . Show that there is a subset I of $\{1, 2, \dots, n\}$ for which the subspace $W = \langle \{\mathbf{e}_i : i \in I\} \rangle$ is a complementary subspace to U in F^n .

9. Let $\alpha : U \rightarrow V$ be a linear map between two finite dimensional vector spaces and let W be a vector subspace of U . Show that the restriction of α to W is a linear map $\alpha|_W : W \rightarrow V$ which satisfies

$$r(\alpha) \geq r(\alpha|_W) \geq r(\alpha) - \dim(U) + \dim(W).$$

Give examples (with $W \neq U$) to show that either of the two inequalities can be an equality.

10. Let $\alpha : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear map given by $\alpha : \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \mapsto \begin{pmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$. Find the matrix

representing α relative to the basis $\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ for both the domain and the range.

Write down bases for the domain and range with respect to which the matrix of α is the identity.

11. Let U_1, \dots, U_k be subspaces of a vector space V and let B_i be a basis for U_i . Show that the following statements are equivalent:

(i) $U = \sum_i U_i$ is a direct sum, i.e. every element of U can be written uniquely as $\sum_i u_i$ with $u_i \in U_i$.

(ii) $U_j \cap \sum_{i \neq j} U_i = \{0\}$ for all j .

(iii) The B_i are pairwise disjoint and their union is a basis for $\sum_i U_i$.

Given an example where $U_i \cap U_j = \{0\}$ for all $i \neq j$, yet $U_1 + \dots + U_k$ is not a direct sum.

12. Let Y and Z be subspaces of the finite dimensional vector spaces V and W , respectively. Show that $R = \{\alpha \in L(V, W) : \alpha(Y) \leq Z\}$ is a subspace of the space $L(V, W)$ of all linear maps from V to W . What is the dimension of R ?

13. Let V be a vector space over F and let W be a subspace. Show that there is a natural way in which the quotient group V/W is a vector space over F . Show that if the dimension of V is finite, then so is the dimension of V/W , and

$$\dim V = \dim W + \dim V/W.$$

14. Suppose X and Y are linearly independent subsets of a vector space V ; no member of X is expressible as a linear combination of members of Y , and no member of Y is expressible as a linear combination of members of X . Is the set $X \cup Y$ necessarily linearly independent? Give a proof or counterexample.

15. Show that any two subspaces of the same dimension in a finite dimensional vector space have a common complementary subspace. [You may wish to consider first the case where the subspaces have dimension 1 less than the space.]

16. (Another version of the Steinitz Exchange Lemma.) Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_r\}$ and $\{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_s\}$ be linearly independent subsets of a vector space V , and suppose $r \leq s$. Show that it is possible to choose distinct indices i_1, i_2, \dots, i_r from $\{1, 2, \dots, s\}$ such that, if we delete each \mathbf{y}_{i_j} from Y and replace it by \mathbf{x}_j , the resulting set is still linearly independent. Deduce that any two maximal linearly independent subsets of a finite dimensional vector space have the same size.

17. Let \mathbb{F}_p be the field of integers modulo p , where p is a prime number. Let V be a vector space of dimension n over \mathbb{F}_p . How many vectors are there in V ? How many (ordered) bases? How many automorphisms does V have? How many k -dimensional subspaces are there in V ?