

**Linear Algebra: Example Sheet 2 of 4**

1. Write down the three types of elementary matrices and find their inverses. Use elementary matrices to find the inverse of

$$\begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 3 & -1 \end{pmatrix}.$$

2. (Another proof of the row rank column rank equality.) Let  $A$  be an  $m \times n$  matrix of (column) rank  $r$ . Show that  $r$  is the least integer for which  $A$  factorises as  $A = BC$  with  $B \in \text{Mat}_{m,r}(\mathbb{F})$  and  $C \in \text{Mat}_{r,n}(\mathbb{F})$ . Using the fact that  $(BC)^T = C^T B^T$ , deduce that the (column) rank of  $A^T$  equals  $r$ .
3. Let  $V$  be a 4-dimensional vector space over  $\mathbb{R}$ , and let  $\{\xi_1, \xi_2, \xi_3, \xi_4\}$  be the basis of  $V^*$  dual to the basis  $\{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4\}$  for  $V$ . Determine, in terms of the  $\xi_i$ , the bases dual to each of the following:
- (a)  $\{\mathbf{x}_2, \mathbf{x}_1, \mathbf{x}_4, \mathbf{x}_3\}$  ;
  - (b)  $\{\mathbf{x}_1, 2\mathbf{x}_2, \frac{1}{2}\mathbf{x}_3, \mathbf{x}_4\}$  ;
  - (c)  $\{\mathbf{x}_1 + \mathbf{x}_2, \mathbf{x}_2 + \mathbf{x}_3, \mathbf{x}_3 + \mathbf{x}_4, \mathbf{x}_4\}$  ;
  - (d)  $\{\mathbf{x}_1, \mathbf{x}_2 - \mathbf{x}_1, \mathbf{x}_3 - \mathbf{x}_2 + \mathbf{x}_1, \mathbf{x}_4 - \mathbf{x}_3 + \mathbf{x}_2 - \mathbf{x}_1\}$  .
4. For  $A \in \text{Mat}_{n,m}(\mathbb{F})$  and  $B \in \text{Mat}_{m,n}(\mathbb{F})$ , let  $\tau_A(B)$  denote  $\text{tr}AB$ . Show that, for each fixed  $A$ ,  $\tau_A: \text{Mat}_{m,n}(\mathbb{F}) \rightarrow \mathbb{F}$  is linear. Show moreover that the mapping  $A \mapsto \tau_A$  defines a linear isomorphism  $\text{Mat}_{n,m}(\mathbb{F}) \rightarrow \text{Mat}_{m,n}(\mathbb{F})^*$ .
5. (a) Let  $V$  be a non-zero finite dimensional real vector space. Show that there are no endomorphisms  $\alpha, \beta$  of  $V$  with  $\alpha\beta - \beta\alpha = \text{id}_V$ .
- (b) Let  $V$  be the space of infinitely differentiable functions  $\mathbb{R} \rightarrow \mathbb{R}$ . Find endomorphisms  $\alpha$  and  $\beta$  of  $V$  such that  $\alpha\beta - \beta\alpha = \text{id}_V$ .
6. Suppose that  $\psi: U \times V \rightarrow \mathbb{F}$  is a bilinear form of rank  $r$  on finite dimensional vector spaces  $U$  and  $V$  over  $\mathbb{F}$ . Show that there exist bases  $e_1, \dots, e_m$  for  $U$  and  $f_1, \dots, f_n$  for  $V$  such that

$$\psi \left( \sum_{i=1}^m x_i e_i, \sum_{j=1}^n y_j f_j \right) = \sum_{k=1}^r x_k y_k$$

for all  $x_1, \dots, x_m, y_1, \dots, y_n \in \mathbb{F}$ . What are the dimensions of the left and right kernels of  $\psi$ ?

7. (a) Let  $a_0, \dots, a_n$  be distinct real numbers, and let

$$A = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ a_0 & a_1 & \cdots & a_n \\ a_0^2 & a_1^2 & \cdots & a_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ a_0^n & a_1^n & \cdots & a_n^n \end{pmatrix}.$$

Show that  $\det(A) \neq 0$ .

- (b) Let  $P_n$  be the space of real polynomials of degree at most  $n$ . For  $x \in \mathbf{R}$  define  $e_x \in P_n^*$  by  $e_x(p) = p(x)$ . By considering the standard basis  $(1, t, \dots, t^n)$  for  $P_n$ , use (a) to show that  $\{e_0, \dots, e_n\}$  is linearly independent and hence forms a basis for  $P_n^*$ .
- (c) Identify the basis of  $P_n$  to which  $(e_0, \dots, e_n)$  is dual.
8. Let  $A, B$  be  $n \times n$  matrices, where  $n \geq 2$ . Show that, if  $A$  and  $B$  are non-singular, then

$$(i) \text{adj}(AB) = \text{adj}(B)\text{adj}(A), \quad (ii) \det(\text{adj} A) = (\det A)^{n-1}, \quad (iii) \text{adj}(\text{adj} A) = (\det A)^{n-2}A.$$

Show that the rank of the adjugate matrix is  $\text{r}(\text{adj } A) = \begin{cases} n & \text{if } \text{r}(A) = n \\ 1 & \text{if } \text{r}(A) = n - 1 \\ 0 & \text{if } \text{r}(A) \leq n - 2. \end{cases}$

Do (i)-(iii) hold if  $A$  is singular? [Hint: for (i) consider  $A + \lambda I$  for  $\lambda \in \mathbb{F}$ .]

9. Show that the dual of the space  $P$  of real polynomials is isomorphic to the space  $\mathbb{R}^{\mathbb{N}}$  of all sequences of real numbers, via the mapping which sends a linear form  $\xi : P \rightarrow \mathbb{R}$  to the sequence  $(\xi(1), \xi(t), \xi(t^2), \dots)$ . In terms of this identification, describe the effect on a sequence  $(a_0, a_1, a_2, \dots)$  of the linear maps dual to each of the following linear maps  $P \rightarrow P$ :
- The map  $D$  defined by  $D(p)(t) = p'(t)$ .
  - The map  $S$  defined by  $S(p)(t) = p(t^2)$ .
  - The composite  $DS$ .
  - The composite  $SD$ .

Verify that  $(DS)^* = S^*D^*$  and  $(SD)^* = D^*S^*$ .

10. Let  $V$  be a vector space. Suppose that  $f_1, \dots, f_n, g \in V^*$ . Show that  $g$  is in the span of  $f_1, \dots, f_n$  if and only if  $\bigcap_{i=1}^n \ker f_i \subset \ker g$ .
11. Let  $\alpha : V \rightarrow V$  be an endomorphism of a real finite dimensional vector space  $V$  with  $\text{tr}(\alpha) = 0$ .
- Show that, if  $\alpha \neq 0$ , there is a vector  $\mathbf{v}$  with  $\mathbf{v}, \alpha(\mathbf{v})$  linearly independent. Deduce that there is a basis for  $V$  relative to which  $\alpha$  is represented by a matrix  $A$  with all of its diagonal entries equal to 0.
  - Show that there are endomorphisms  $\beta, \gamma$  of  $V$  with  $\alpha = \beta\gamma - \gamma\beta$ .