

1. Let  $X = (X_n)_{n \geq 0}$  be a sequence of independent random variables. Show that  $X$  is a Markov chain. Under what condition is this chain homogeneous?

2. Let  $X = (X_n)_{n \geq 0}$  be a sequence of fair coin tosses (with the two possible outcomes interpreted as 0 and 1) and set  $M_n = \max_{k \leq n} X_k$ . Show that  $(M_n)_{n \geq 0}$  is a Markov chain and find the transition probabilities.

3. Let  $X = (X_n)_{n \geq 0}$  be a Markov chain and let  $(n_r)_{r \geq 0}$  be an unbounded increasing sequence of positive integers. Show that  $Y_r = X_{n_r}$  defines a (possibly inhomogeneous) Markov chain. Find the transition probabilities of  $Y$  when  $n_r = 2r$  and  $X$  is a simple random walk.

4. (Harder) Let  $S = (S_n)_{n \geq 0}$  be a simple (possibly asymmetric) random walk on  $\mathbb{Z}$  with  $S_0 = 0$ . Show that  $X_n = |S_n|$  defines a Markov chain and find its transition probabilities. Let  $M_n = \max_{k \leq n} S_k$  and show that  $Y_n = M_n - S_n$  defines a Markov chain.

5. Let  $X = (X_n)_{n \geq 0}$  and  $Y = (Y_n)_{n \geq 0}$  be Markov chains on the integers  $\mathbb{Z}$ . Is  $Z_n = X_n + Y_n$  necessarily a Markov chain. Justify your answer.

6. A flea hops about at random on the vertices of a triangle where each hop is from the currently occupied vertex of one of the other two vertices each with probability  $1/2$ . Find the probability that after  $n$  hops the flea is back where it started.

Now suppose that the flea is twice as likely to jump clockwise as anticlockwise. What is the probability that after  $n$  hops the flea is back where it started now? [Hint:  $1/2 \pm i/(2\sqrt{3}) = (1/\sqrt{3})e^{\pm i\pi/6}$ .]

7. A die is ‘fixed’ so that when it is rolled the score cannot be the same as the previous score, all other scores having probability  $1/5$ . If the first score is 6, what is the probability  $p$  that the  $n$ th score is 6? What is the probability that the  $n$ th score is  $j$ , where  $j \neq 6$ ?

Suppose instead that the die cannot score one greater (mod 6) than the previous score, all other five scores having equal probability. What is the new value of  $p$ ? [Hint: Think about the relationship between the two dice.]

8. Let  $X = (X_n)_{n \geq 0}$  be a Markov chain on  $\{1, 2, 3\}$  with transition matrix

$$P = \begin{bmatrix} 0 & 1 & 0 \\ 0 & \frac{2}{3} & \frac{1}{3} \\ p & 1-p & 0 \end{bmatrix}.$$

Find  $\mathbb{P}[X_n = 1 | X_0 = 1]$  in each of the following cases: (a)  $p = 1/16$ , (b)  $p = 1/6$ , (c)\*  $p = 1/12$ .

9. Identify the communicating classes of the transition matrix

$$P = \begin{bmatrix} \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & 0 & \frac{1}{2} & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{2} & 0 & 0 & 0 & \frac{1}{2} \end{bmatrix}.$$

Which of the classes are closed?

10. Show that every transition matrix on a finite state space has at least one closed communicating class. Find an example of a transition matrix with no closed communicating class.

11. A gambler has £2 and needs to increase it to £10 in a hurry. She can play a game with the following rules: a fair coin is tossed; if a player bets on the side which actually turns up, she wins a sum equal to her stake, and her stake is returned; otherwise she loses her stake. The gambler decides to use a bold strategy in which she stakes all her money if she has £5 or less and otherwise stakes just enough to increase her capital, if she wins, to £10.

Let  $X_0 = 2$  and  $X = (X_n)_{n \geq 0}$  be her capital after  $n$  throws. Prove that the gambler will achieve her aim with probability  $1/5$ . What is the expected number of tosses until she either achieves her aim or loses her capital?

12. Let  $X = (X_n)_{n \geq 0}$  be a Markov chain on  $\{0, 1, \dots\}$  with transition probabilities given by

$$p_{0,1} = 1, \quad p_{i,i+1} + p_{i,i-1} = 1, \quad p_{i,i+1} = \left(\frac{i+1}{i}\right)^2 p_{i,i-1}, \quad (i \geq 1).$$

Show that if  $X_0 = 0$  then the probability that  $X_n \geq 1$  for all  $n \geq 1$  is  $6/\pi^2$ .

13. Let  $Y_1, Y_2, \dots$  be i.i.d. random variables with  $\mathbb{P}[Y_1 = 1] = \mathbb{P}[Y_1 = -1] = 1/2$  and set  $X_0 = 1$ ,  $X_n = X_0 + Y_1 + \dots + Y_n$  for  $n \geq 1$ . Define

$$H_0 = \inf\{n \geq 0 : X_n = 0\}.$$

Find the probability generating function  $\phi(s) = \mathbb{E}[s^{H_0}]$ .

Suppose the common distribution of the  $Y_i$  is changed to  $\mathbb{P}[Y_1 = 2] = \mathbb{P}[Y_1 = -1] = 1/2$ . Show that the probability generating function  $\phi$  now satisfies

$$s\phi^3 - 2\phi + s = 0.$$