

## Part II

## Algebraic Geometry

### Example Sheet IV, 2016

(For all questions, assume  $k$  is algebraically closed. Further, you can assume the characteristic is not equal to 2 if necessary. A \* indicates a more difficult problem. I should note in general that this example sheet is a fair bit harder than the previous ones, so don't despair if you don't get all the problems!)

1. If  $P$  is a smooth point of an irreducible curve  $X$  and  $t \in \mathcal{O}_{X,P}$  is a local parameter at  $P$ , show that  $\dim_k \mathcal{O}_{X,P}/(t^n) = n$  for every  $n \in \mathbb{N}$ .
2. Show that  $X = Z(x_0^8 + x_1^8 + x_2^8)$  and  $Y = Z(y_0^4 + y_1^4 + y_2^4)$  are irreducible smooth curves in  $\mathbb{P}^2$  provided  $\text{char}(k) \neq 2$ , and that  $\phi: (x_i) \mapsto (x_i^2)$  is a morphism from  $X$  to  $Y$ . Determine the degree of  $\phi$ , and compute  $e_P$  for all  $P \in X$ .
3. Show that the plane cubic  $X = Z(f)$ ,  $f = x_0x_2^2 - x_1^3 + 3x_1x_0^2$ , is smooth if  $\text{char}(k) \neq 2, 3$ . Find the degree and ramification degrees (i.e., the  $e_P$ ) for (i) the projection  $\phi = (x_0 : x_1): X \rightarrow \mathbb{P}^1$  (ii) the projection  $\phi = (x_0 : x_2): X \rightarrow \mathbb{P}^1$ .
4. Let  $X$  be a non-singular projective curve. Let  $V \subset K(X)$  be a finite-dimensional  $k$ -vector subspace of  $K(X)$ . Show that there exists a divisor  $D$  on  $X$  for which  $V \subset \mathcal{L}(D)$ .
5. Let  $X$  be a smooth plane cubic. Assume that  $V$  has equation  $x_0x_2^2 = x_1(x_1 - x_0)(x_1 - \lambda x_0)$ , for some  $\lambda \in k \setminus \{0, 1\}$ .  
Let  $P = (0 : 0 : 1)$  be the point at infinity in this equation. Writing  $x = x_1/x_0$ ,  $y = x_2/x_0$ , show that  $x/y$  is a local parameter at  $P$ . [Hint: consider the affine piece  $x_2 \neq 0$ .] Hence compute  $v_P(x)$  and  $v_P(y)$ . Show that for each  $m \geq 1$ , the space  $\mathcal{L}(mP)$  has a basis consisting of functions  $x^i, x^jy$ , for suitable  $i$  and  $j$ , and that  $\ell(mP) = m$ .
6. Let  $f \in k[x]$  a polynomial of degree  $d > 1$  with distinct roots, and  $V \subset \mathbb{P}^2$  the projective closure of the affine curve with equation  $y^{d-1} = f(x)$ . Assume that  $\text{char}(k)$  does not divide  $d - 1$ . Prove that  $V$  is smooth, and has a single point  $P$  at infinity. Calculate  $v_P(x)$  and  $v_P(y)$ .  
\* Deduce (without using Riemann–Roch) that if  $n > d(d-3)$ , then  $\ell((n+1)P) = \ell(nP) + 1$ .
7. A non-singular projective curve  $X$  is covered by two affine pieces (with respect to different embeddings) which are affine plane curves with equations  $y^2 = f(x)$  and  $v^2 = g(u)$  respectively, with  $f$  a square-free polynomial of even degree  $2n$  and  $u = 1/x$ ,  $v = y/x^n$  in  $K(X)$ . Determine the polynomial  $g(u)$  and show that

the canonical class on  $X$  has degree  $2n - 4$ . Why can we not just say that  $X$  is the projective plane curve associated to the affine curve  $y^2 = f(x)$ ?

8. Let  $X_0 \subset \mathbb{A}^2$  be the affine curve with equation  $y^3 = x^4 + 1$ , and let  $X \subset \mathbb{P}^2$  be its projective closure. Show that  $X$  is smooth, and has a unique point  $Q$  at infinity. Let  $\omega$  be the rational differential  $dx/y^2$  on  $X$ . Show that  $v_P(\omega) = 0$  for all  $P \in X_0$ . prove that  $v_Q(\omega) = 4$  and hence that  $\omega$ ,  $x\omega$  and  $y\omega$  are all regular on  $X$ .
9. Let  $X$  be a non-singular projective curve and  $P \in X$  any point. Show that there exists a nonconstant rational function on  $X$  which is regular everywhere except at  $P$ . Show moreover that there exists a projective embedding of  $X$  which has  $P$  as its unique point at infinity.
10. Let  $P_0$  be a point on an elliptic curve (non-singular projective curve of genus 1!) and  $\phi_{3P_0}: X \rightarrow \mathbb{P}^2$  the projective embedding. Show that  $P \in X$  is a point of inflection if and only if  $3P = 0$  in the group law determined by  $P_0$ . Deduce that if  $P$  and  $Q$  are points of inflection then so is the third point of intersection of the line  $PQ$  with  $X$ .
11. Let  $X$  be a projective non-singular curve of genus 2, and let  $K$  be an effective canonical divisor on  $X$ . Consider the divisor  $K + P_1 + P_2$  for points  $P_i$  with  $P_1 + P_2 \not\sim K$ . Show that the linear system associated to this divisor induces a morphism  $\phi$  from  $X$  to a quartic curve in  $\mathbb{P}^2$ . Show furthermore that  $\phi(P) = \phi(Q)$ , with  $P \neq Q$ , if and only if  $\{P, Q\} = \{P_1, P_2\}$ .
12. Let  $\pi: X \rightarrow \mathbb{P}^1$  be a hyperelliptic cover, and  $P \neq Q$  ramification points of  $\pi$ . Show that as elements of  $Cl^0(X)$ ,  $P - Q \neq 0$  but  $2(P - Q) = 0$ .