

**Exercise 2.1.** Let  $X$  be a normed space. Show that  $X'$  equipped with its norm forms a Banach space. If  $\overline{X}$  is the completion of  $X$  with respect to the metric induced by its norm, show that  $X' = \overline{X}'$ .

**Exercise 2.2.** Suppose  $X$  is a Banach space. Show that if  $\Lambda \in X'$  with  $\Lambda \neq 0$  then  $\Lambda$  is an open mapping (i.e.  $\Lambda(U)$  is open whenever  $U \subset X$  is open).

**Exercise 2.3.** Let  $u : L^p(\mathbb{R}^n; \mathbb{R}) \rightarrow \mathbb{R}$  be a bounded, linear functional.

a) For  $f \in L^p(\mathbb{R}^n; \mathbb{R})$ ,  $f \geq 0$ , define

$$\tilde{u}(f) = \sup\{u(g) : g \in L^p(\mathbb{R}^n; \mathbb{R}), 0 \leq g \leq f\}.$$

Show that  $0 \leq \tilde{u}(f)$  and  $u(f) \leq \tilde{u}(f) \leq \|u\|_{L^{p'}} \|f\|_{L^p}$ , and establish

$$\tilde{u}(f + ag) = \tilde{u}(f) + a\tilde{u}(g)$$

for all  $f, g \in L^p(\mathbb{R}^n; \mathbb{R})$  with  $f, g \geq 0$  and  $a \in \mathbb{R}$ ,  $a > 0$ .

b) For  $f \in L^p(\mathbb{R}^n; \mathbb{R})$ , define  $w(f) = \tilde{u}(f^+) - \tilde{u}(f^-)$ , where  $f^+(x) = \max\{0, f(x)\}$ ,  $f^-(x) = \max\{0, -f(x)\}$ . Show that  $w$  is linear and bounded, and that  $w$  and  $w - u$  are positive.

c) Deduce that  $u = u_+ - u_-$ , where  $u_{\pm}$  are bounded, positive, linear functionals.

**Exercise 2.4.** Suppose  $X$  is a normed space, and  $V \subset X$  is a closed proper subspace of  $X$  and let  $0 < \alpha < 1$ . Show that there exists  $x \in X$  with  $\|x\| = 1$  such that  $\|x - y\| \geq \alpha$  for all  $y \in V$ . Deduce that the Bolzano–Weierstrass theorem does not hold if  $X$  is an infinite dimensional Banach space.

[The first result above is known as Riesz' Lemma]

**Exercise 2.5.** Let  $\mathcal{P}$  be a separating family of seminorms on a vector space  $X$ . Show that a sequence  $(x_k)_{k=1}^{\infty}$  with  $x_k \in X$  converges to  $x \in X$  in the topology  $\tau_{\mathcal{P}}$  if and only if  $p(x_k - x) \rightarrow 0$  for all  $p \in \mathcal{P}$ .

**Exercise 2.6.** Suppose that  $X$  is a Banach space, and let  $(\Lambda_k)_{k=1}^{\infty}$  be a sequence with  $\Lambda_k \in X'$ . Show that:

$$\Lambda_k \rightarrow \Lambda \implies \Lambda_k \rightharpoonup \Lambda \implies \Lambda_k \xrightarrow{*} \Lambda.$$

(\*) Show the stronger statement that  $\tau_{w*} \subset \tau_w \subset \tau_s$ , where  $\tau_{w*}, \tau_w, \tau_s$  are the weak-\*, weak and strong topologies on  $X'$  respectively.

**Exercise 2.7.** For a bounded measurable set  $E \subset \mathbb{R}^n$  of positive measure, and any  $f \in L^1_{loc}(\mathbb{R}^n)$ , define the mean of  $f$  on  $E$  to be:

$$\int_E f(x)dx = \frac{1}{|E|} \int_E f(x)dx.$$

Suppose  $1 < p < \infty$  and let  $(f_j)_{j=1}^\infty$  be a bounded sequence in  $L^p(\mathbb{R}^n)$ . Show that  $f_j \rightharpoonup f$  for some  $f \in L^p(\mathbb{R}^n)$  if and only if

$$\int_E f_j(x)dx \rightarrow \int_E f(x)dx$$

for all bounded measurable sets  $E \subset \mathbb{R}^n$  of positive measure.

**Exercise 2.8.** Suppose  $(H, (\cdot, \cdot))$  is an infinite dimensional Hilbert space and let  $(x_i)_{i=1}^\infty$  be a sequence with  $x_i \in H$ .

i) Show that  $x_i \rightharpoonup x$  if and only if  $(y, x_i) \rightarrow (y, x)$  for all  $y \in H$ .

ii) Show there exists a sequence such that  $x_i \rightharpoonup 0$ , but  $x_i \not\rightarrow 0$ .

iii) Suppose  $x_i \rightharpoonup x$ . Show that

$$\|x\| \leq \liminf_{i \rightarrow \infty} \|x_i\|,$$

and  $\|x_i\| \rightarrow \|x\|$  iff  $x_i \rightarrow x$ .

**Exercise 2.9.** Construct a bounded sequence  $(f_i)_{i=1}^\infty$  of functions  $f_i \in L^1(\mathbb{R})$  such that no subsequence is weakly convergent.

**Exercise 2.10.** Let  $X$  be a Banach space and suppose  $A \subset X$  is a convex neighbourhood of 0. For  $x \in X$  define  $\mu_A(x) = \inf\{t > 0 : t^{-1}x \in A\}$ . Show that  $\mu_A$  is sublinear and satisfies  $\mu_A(x) \leq k\|x\|$  for some  $k > 0$ . Show further that  $\mu_A(y) < 1$  for  $y \in A$ .  $\mu_A$  is called the Minkowski functional of  $A$

**Exercise 2.11.** Let  $\{x_1, \dots, x_n\}$  be a set of linearly independent elements of a Banach space  $X$ . Let  $a_1, \dots, a_n \in \mathbb{C}$ . Show that there exists  $\Lambda \in X'$  such that  $\Lambda(x_i) = a_i$ , for  $i = 1, \dots, n$ .

**Exercise 2.12.** Let  $M$  be a vector subspace of the Banach space  $X$ , and suppose that  $K \subset X$  is open, convex and disjoint from  $M$ . Show that there exists a co-dimension one subspace  $N \subset X$  which contains  $M$  and is disjoint from  $K$ .

*This is Mazur's theorem.*

**Exercise 2.13.** let  $X$  be a reflexive Banach space, and suppose  $Y \subset X$  is a closed subspace. Show that  $Y$  is reflexive.