

Probability and Measure 2

3.1. Suppose that a simple function f has two representations

$$f = \sum_{k=1}^m a_k 1_{A_k} = \sum_{j=1}^n b_j 1_{B_j}.$$

For $\varepsilon = (\varepsilon_1, \dots, \varepsilon_m) \in \{0, 1\}^m$, define $A_\varepsilon = A_1^{\varepsilon_1} \cap \dots \cap A_m^{\varepsilon_m}$ where $A_k^0 = A_k^c$ and $A_k^1 = A_k$. Define similarly B_δ for $\delta \in \{0, 1\}^n$. Then set $f_{\varepsilon, \delta} = \sum_{k=1}^m \varepsilon_k a_k$ if $A_\varepsilon \cap B_\delta \neq \emptyset$ and $f_{\varepsilon, \delta} = 0$ otherwise. Show that, for any measure μ ,

$$\sum_{k=1}^m a_k \mu(A_k) = \sum_{\varepsilon, \delta} f_{\varepsilon, \delta} \mu(A_\varepsilon \cap B_\delta)$$

and deduce that

$$\sum_{k=1}^m a_k \mu(A_k) = \sum_{j=1}^n b_j \mu(B_j).$$

3.2. Let μ and ν be finite Borel measures on \mathbb{R} . Let f be a continuous bounded function on \mathbb{R} . Show that f is integrable with respect to μ and ν . Show further that, if $\mu(f) = \nu(f)$ for all such f , then $\mu = \nu$.

3.3. Let f be an integrable function on a measure space (E, \mathcal{E}, μ) . Suppose that, for some π -system \mathcal{A} containing E and generating \mathcal{E} , we have $\mu(f 1_A) = 0$ for all $A \in \mathcal{A}$. Show that $f = 0$ a.e.

3.4. Let X be a non-negative integer-valued random variable. Show that

$$\mathbb{E}(X) = \sum_{n=1}^{\infty} \mathbb{P}(X \geq n).$$

Deduce that, if $\mathbb{E}(X) = \infty$ and X_1, X_2, \dots is a sequence of independent random variables with the same distribution as X , then, almost surely, $\limsup_n (X_n/n) \geq 1$, and moreover $\limsup_n (X_n/n) = \infty$.

Now suppose that Y_1, Y_2, \dots is any sequence of independent identically distributed random variables with $\mathbb{E}|Y_1| = \infty$. Show that, almost surely, $\limsup_n (|Y_n|/n) = \infty$, and moreover $\limsup_n (|Y_1 + \dots + Y_n|/n) = \infty$.

3.5. For $\alpha \in (0, \infty)$ and $x \in (0, \infty)$, define $f_\alpha(x) = x^{-\alpha}$. Show that f_α is integrable with respect to Lebesgue measure on $(0, 1]$ if and only if $\alpha < 1$. Show also that f_α is integrable with respect to Lebesgue measure on $[1, \infty)$ if and only if $\alpha > 1$.

3.6. Show that the function $\sin x/x$ is not Lebesgue integrable over $[1, \infty)$ but that integral $\int_1^N (\sin x/x) dx$ converges as $N \rightarrow \infty$.

3.7. Show that, as $n \rightarrow \infty$,

$$\int_0^\infty \sin(e^x)/(1 + nx^2) dx \rightarrow 0 \quad \text{and} \quad \int_0^1 (n \cos x)/(1 + n^2 x^{\frac{3}{2}}) dx \rightarrow 0.$$

3.8. Let u and v be differentiable functions on \mathbb{R} with continuous derivatives u' and v' . Suppose that uv' and $u'v$ are integrable on \mathbb{R} and $u(x)v(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Show that

$$\int_{\mathbb{R}} u(x)v'(x) dx = - \int_{\mathbb{R}} u'(x)v(x) dx.$$

3.9. Let (E, \mathcal{E}) and (G, \mathcal{G}) be measurable spaces and let $f : E \rightarrow G$ be a measurable function. Given a measure μ on (E, \mathcal{E}) , consider the image measure $\nu = \mu \circ f^{-1}$ on (G, \mathcal{G}) . Show that $\nu(g) = \mu(g \circ f)$ for all non-negative measurable functions g on G .

3.10. The moment generating function ϕ of a real-valued random variable X is defined by $\phi(\theta) = \mathbb{E}(e^{\theta X})$, $\theta \in \mathbb{R}$.

Suppose that ϕ is finite on an open interval containing 0. Show that ϕ has derivatives of all orders at 0 and that X has finite moments of all orders given by

$$\mathbb{E}(X^n) = \left(\frac{d}{d\theta} \right)^n \Big|_{\theta=0} \phi(\theta).$$

3.11. Let X_1, \dots, X_n be random variables with density functions f_1, \dots, f_n respectively. Suppose that the \mathbb{R}^n -valued random variable $X = (X_1, \dots, X_n)$ also has a density function f . Show that X_1, \dots, X_n are independent if and only if

$$f(x_1, \dots, x_n) = f_1(x_1) \dots f_n(x_n) \quad \text{a.e.}$$

3.12. Show that, for all non-negative measurable functions f on $[0, \infty)$, the function $(x, y) \mapsto f(|(x, y)|)$ is measurable on \mathbb{R}^2 and (without using the Jacobian formula)

$$\int_{\mathbb{R}^2} f(|(x, y)|) dx dy = 2\pi \int_0^\infty r f(r) dr.$$

Hence show that $(2\pi)^{-1/2} e^{-x^2/2}$ is a probability density function.

3.13. Let μ and ν be probability measures on (E, \mathcal{E}) and let $f : E \rightarrow [0, R]$ be a measurable function. Suppose that $\nu(A) = \mu(f1_A)$ for all $A \in \mathcal{E}$. Let $(X_n : n \in \mathbb{N})$ be a sequence of independent random variables in E with law μ and let $(U_n : n \in \mathbb{N})$ be a sequence of independent $U[0, 1]$ random variables. Set

$$T = \min\{n \in \mathbb{N} : RU_n \leq f(X_n)\}, \quad Y = X_T.$$

Show that Y has law ν . (This justifies simulation by rejection sampling.)