

Probability and Measure 4

7.8. Let $(X_n : n \in \mathbb{N})$ be a sequence of random variables in \mathbb{R} and let X be another such random variable. Show that $X_n \rightarrow X$ weakly if and only if $X_n \rightarrow X$ in distribution.

7.9. Let μ be a Borel probability measure on \mathbb{R}^d and let $(\mu_n : n \in \mathbb{N})$ be a sequence of such measures. Suppose that $\mu_n(f) \rightarrow \mu(f)$ for all C^∞ functions on \mathbb{R}^d of compact support. Show that μ_n converges weakly to μ on \mathbb{R}^d .

8.1. Let $X = (X_1, \dots, X_n)$ be a Gaussian random variable in \mathbb{R}^n with mean μ and covariance matrix V . Assume that V is invertible write $V^{-1/2}$ for the positive-definite square root of V^{-1} . Set $Y = (Y_1, \dots, Y_n) = V^{-1/2}(X - \mu)$. Show that Y_1, \dots, Y_n are independent $N(0, 1)$ random variables. Show further that we can write X_2 in the form $X_2 = aX_1 + Z$ where Z is independent of X_1 and determine the distribution of Z .

8.2. Let X_1, \dots, X_n be independent $N(0, 1)$ random variables. Show that

$$\left(\bar{X}, \sum_{m=1}^n (X_m - \bar{X})^2 \right) \quad \text{and} \quad \left(\frac{X_n}{\sqrt{n}}, \sum_{m=1}^{n-1} X_m^2 \right)$$

have the same distribution, where $\bar{X} = (X_1 + \dots + X_n)/n$.

9.1. Let (E, \mathcal{E}, μ) be a measure space and $\tau : E \rightarrow E$ a measure-preserving transformation. Show that $\mathcal{E}_\tau := \{A \in \mathcal{E} : \tau^{-1}(A) = A\}$ is a σ -algebra, and that a measurable function f is \mathcal{E}_τ -measurable if and only if it is *invariant*, that is $f \circ \tau = f$.

9.2. Show that, if θ is an ergodic measure-preserving transformation and f is a θ -invariant function, then there exists a constant $c \in \mathbb{R}$ such that $f = c$ a.e..

9.3. For $x \in [0, 1)$, set $\tau(x) = 2x \bmod 1$. Show that τ is a measure-preserving transformation of $([0, 1), \mathcal{B}([0, 1)), dx)$, and that τ is ergodic. Identify the invariant function \bar{f} corresponding to each integrable function f .

9.4. Fix $a \in [0, 1)$ and define, for $x \in [0, 1)$, $\tau(x) = x + a \bmod 1$. Show that τ is also a measure-preserving transformation of $([0, 1), \mathcal{B}([0, 1)), dx)$. Determine for which values of a the transformation τ is ergodic. *Hint: you may use the fact that any integrable function f on $[0, 1)$ whose Fourier coefficients all vanish must itself vanish a.e..* Identify, for all values of a , the invariant function \bar{f} corresponding to an integrable function f .

9.5. Call a sequence of random variables $(X_n : n \in \mathbb{N})$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ *stationary* if for each $n, k \in \mathbb{N}$ the random vectors (X_1, \dots, X_n) and $(X_{k+1}, \dots, X_{k+n})$ have the same distribution: for $A_1, \dots, A_n \in \mathcal{B}$,

$$\mathbb{P}(X_1 \in A_1, \dots, X_n \in A_n) = \mathbb{P}(X_{k+1} \in A_1, \dots, X_{k+n} \in A_n).$$

Show that, if $(X_n : n \in \mathbb{N})$ is a stationary sequence and $X_1 \in L^p$, for some $p \in [1, \infty)$, then

$$\frac{1}{n} \sum_{i=1}^n X_i \rightarrow X \quad \text{a.s. and in } L^p,$$

for some random variable $X \in L^p$ and find $\mathbb{E}(X)$.

10.1. Let $(X_n : n \in \mathbb{N})$ be a sequence of independent random variables, such that $\mathbb{E}(X_n) = \mu$ and $\mathbb{E}(X_n^4) \leq M$ for all n , for some constants $\mu \in \mathbb{R}$ and $M < \infty$. Set $P_n = X_1 X_2 + X_2 X_3 + \dots + X_{n-1} X_n$. Show that P_n/n converges a.s. as $n \rightarrow \infty$ and identify the limit.

10.2. The Cauchy distribution has density function $f(x) = \pi^{-1}(1+x^2)^{-1}$ for $x \in \mathbb{R}$. Show that the corresponding characteristic function is given by $\varphi(u) = e^{-|u|}$. Show also that, if X_1, \dots, X_n are independent Cauchy random variables, then the random variable $(X_1 + \dots + X_n)/n$ is also Cauchy.

10.3. Let f be a bounded continuous function on $(0, \infty)$, having Laplace transform

$$\hat{f}(\lambda) = \int_0^\infty e^{-\lambda x} f(x) dx, \quad \lambda \in (0, \infty).$$

Let $(X_n : n \in \mathbb{N})$ be a sequence of independent exponential random variables, of parameter λ . Show that \hat{f} has derivatives of all orders on $(0, \infty)$ and that, for all $n \in \mathbb{N}$, for some $C(\lambda, n) \neq 0$ independent of f , we have

$$(d/d\lambda)^{n-1} \hat{f}(\lambda) = C(\lambda, n) \mathbb{E}(f(S_n))$$

where $S_n = X_1 + \dots + X_n$. Deduce that if $\hat{f} \equiv 0$ then also $f \equiv 0$.

10.4. For each $n \in \mathbb{N}$, there is a unique probability measure μ_n on the unit sphere $S^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$ such that $\mu_n(A) = \mu_n(UA)$ for all Borel sets A and all orthogonal $n \times n$ matrices U . Fix $k \in \mathbb{N}$ and, for $n \geq k$, let γ_n denote the probability measure on \mathbb{R}^k which is the law of $\sqrt{n}(x^1, \dots, x^k)$ under μ_n . Show

(a) if $X \sim N(0, I_n)$ then $X/|X| \sim \mu_n$,

(b) if $(X_n : n \in \mathbb{N})$ is a sequence of independent $N(0, 1)$ random variables and if $R_n = \sqrt{X_1^2 + \dots + X_n^2}$ then $R_n/\sqrt{n} \rightarrow 1$ a.s.,

(c) γ_n converges weakly to the standard Gaussian distribution on \mathbb{R}^k as $n \rightarrow \infty$.