

## PART II REPRESENTATION THEORY SHEET 1

*Unless otherwise stated, all groups here are finite, and all vector spaces are finite-dimensional over a field  $F$  of characteristic zero, usually  $\mathbb{C}$ .*

- 1 Let  $\rho$  be a representation of the group  $G$ .
  - (a) Show that  $\delta : g \mapsto \det \rho(g)$  is a 1-dimensional representation of  $G$ .
  - (b) Prove that  $G/\ker \delta$  is abelian.
  - (c) Assume that  $\delta(g) = -1$  for some  $g \in G$ . Show that  $G$  has a normal subgroup of index 2.
- 2 Let  $\theta : G \rightarrow F^\times$  be a 1-dimensional representation of the group  $G$ , and let  $\rho : G \rightarrow \mathrm{GL}(V)$  be another representation. Show that  $\theta \otimes \rho : G \rightarrow \mathrm{GL}(V)$  given by  $\theta \otimes \rho : g \mapsto \theta(g) \cdot \rho(g)$  is a representation of  $G$ , and that it is irreducible if and only if  $\rho$  is irreducible.
- 3 Find an example of a representation of some finite group over some field of characteristic  $p$ , which is not completely reducible. Find an example of such a representation in characteristic 0 for an infinite group.
- 4 Let  $N$  be a normal subgroup of the group  $G$ . Given a representation of the quotient  $G/N$ , use it to obtain a representation of  $G$ . Which representations of  $G$  do you get this way?  
Recall that the derived subgroup  $G'$  of  $G$  is the unique smallest normal subgroup of  $G$  such that  $G/G'$  is abelian. Show that the 1-dimensional complex representations of  $G$  are precisely those obtained from  $G/G'$ .
- 5 Describe Weyl's unitary trick.  
Let  $G$  be a finite group acting on a complex vector space  $V$ , and let  $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$  be a skew-symmetric form, i.e.  $\langle y, x \rangle = -\langle x, y \rangle$  for all  $x, y$  in  $V$ .  
Show that the form  $(x, y) = \frac{1}{|G|} \sum \langle gx, gy \rangle$ , where the sum is over all elements  $g \in G$ , is a  $G$ -invariant skew-symmetric form.  
Does this imply that every finite subgroup of  $\mathrm{GL}_{2m}(\mathbb{C})$  is conjugate to a subgroup of the symplectic group<sup>1</sup>  $\mathrm{Sp}_{2m}(\mathbb{C})$ ?
- 6 Let  $G = \langle g \rangle$  be a cyclic group of order  $n$ .
  - (i)  $G$  acts on  $\mathbb{R}^2$  as symmetries of the regular  $n$ -gon. Choose a basis of  $\mathbb{R}^2$ , and write the matrix  $R(g)$  representing the action of a generator  $g$  in this basis. Is this an irreducible representation?
  - (ii) Now regard  $R(g)$  above as a complex matrix, so that we get a representation of  $G$  on  $\mathbb{C}^2$ . Decompose  $\mathbb{C}^2$  into its irreducible summands.
- 7 Let  $G$  be a cyclic group of order  $n$ . Explicitly decompose the complex regular representation of  $G$  as a direct sum of 1-dimensional representations, by giving the matrix of change of coordinates from the natural basis  $\{e_g\}_{g \in G}$  to a basis where the group action is diagonal.

---

<sup>1</sup>the group of all linear transformations of a  $2m$ -dimensional vector space over  $\mathbb{C}$  that preserve a non-degenerate, skew-symmetric, bilinear form.

**8** Let  $G$  be the dihedral group  $D_{10}$  of order 10,

$$D_{10} = \langle x, y : x^5 = 1 = y^2, yxy^{-1} = x^{-1} \rangle.$$

Show that  $G$  has precisely two 1-dimensional representations. By considering the effect of  $y$  on an eigenvector of  $x$  show that any complex irreducible representation of  $G$  of dimension at least 2 is isomorphic to one of two representations of dimension 2. Show that all these representations can be realised over  $\mathbb{R}$ .

**9** Let  $G$  be the quaternion group  $Q_8$  of order 8,

$$Q_8 = \langle x, y \mid x^4 = 1, y^2 = x^2, yxy^{-1} = x^{-1} \rangle.$$

By considering the effect of  $y$  on an eigenvector of  $x$  show that any complex irreducible representation of  $G$  of dimension at least 2 is isomorphic to the standard representation of  $Q_8$  of dimension 2.

Show that this 2-dimensional representation cannot be realised over  $\mathbb{R}$ ; that is,  $Q_8$  is not a subgroup of  $\mathrm{GL}_2(\mathbb{R})$ .

**10** Suppose that  $F$  is algebraically closed. Using Schur's lemma, show that if  $G$  is a finite group with trivial centre and  $H$  is a subgroup of  $G$  with non-trivial centre, then any faithful representation of  $G$  is reducible on restriction to  $H$ . What happens for  $F = \mathbb{R}$ ?

**11** Let  $G$  be a subgroup of order 18 of the symmetric group  $S_6$  given by

$$G = \langle (123), (456), (23)(56) \rangle.$$

Show that  $G$  has a normal subgroup of order 9 and four normal subgroups of order 3. By considering quotients, show that  $G$  has two representations of degree 1 and four inequivalent irreducible representations of degree 2. Deduce that  $G$  has no faithful irreducible representations.

**12** Show that if  $\rho$  is a homomorphism from the finite group  $G$  to  $\mathrm{GL}_n(\mathbb{R})$ , then there is a matrix  $P \in \mathrm{GL}_n(\mathbb{R})$  such that  $P\rho(g)P^{-1}$  is an orthogonal matrix for each  $g \in G$ . (Recall that the real matrix  $A$  is orthogonal if  $A^t A = I$ .)

Determine all finite groups which have a faithful 2-dimensional representation over  $\mathbb{R}$ .