## Part IID RIEMANN SURFACES (2005–2006): Example Sheet 2

(a.g.kovalev@dpmms.cam.ac.uk)

**1.** If f is a meromorphic doubly-periodic (i.e. elliptic) function of degree k > 0 show that f' is an elliptic function whose degree  $\ell$  satisfies  $k + 1 \leq \ell \leq 2k$ . Give examples to show that both bounds are attained.

Recall from example sheet 1:  $\psi(z,\tau) = \sum_{n=-\infty}^{\infty} \mathbf{e}(\frac{1}{2}(n+\frac{1}{2})^2\tau + (n+\frac{1}{2})(z+\frac{1}{2}))$  and satisfies  $\psi(z+1) = -\psi(z), \ \psi(z+\tau) = -\mathbf{e}(-\frac{\tau}{2}-z)\psi(z)$ , where  $\mathbf{e}(z) = \exp(2\pi i z), \ \psi(z) = -\psi(-z)$ , and has unique zero 'modulo the lattice  $\mathbb{Z} + \tau \mathbb{Z}$ '.

**2.** (i) Prove that if  $z, w \in \mathbb{C}$ , then

$$\wp(z) - \wp(w) = -\psi'(0)^2 \frac{\psi(z-w)\psi(z+w)}{\psi(z)^2\psi(w)^2}.$$

[Hint: Regarding one of w, z as parameter, prove that each side is  $\Lambda$ -periodic in the other variable and has same zeros and poles. Get multiplicative constant by considering Laurent expansion at zero.]

(ii) Deduce that  $\wp'(z) = -\psi'(0)^3 \frac{\psi(2z)}{\psi(z)^4}$  and recover from this formula the zeros of  $\wp'$ .

**3.** Let  $\chi(z) = \psi'(z)/\psi(z)$ . Differentiate 2(i) and interchange z and w to obtain:

$$\frac{1}{2} \frac{\wp'(z) - \wp'(w)}{\wp(z) - \wp(w)} = \chi(z+w) - \chi(z) - \chi(w).$$

Remark for readers of Ahlfors or Jones & Singerman: their  $\sigma$  and  $\zeta$  are not quite the same as  $\psi$  and  $\chi$  here, but for some constants  $A, B, \sigma(z) = \exp(Az^2 + B)\psi(z)$ , so  $\zeta(z) = 2Az + \chi(z)$ .

**4.**<sup>\*</sup> (challenging but feasible) Prove the addition formula for  $\wp$ ,

$$\wp(z+w) = -\wp(z) - \wp(w) + \frac{1}{4} \left(\frac{\wp'(z) - \wp'(w)}{\wp(z) - \wp(w)}\right)^2.$$

[You will need to differentiate the formula in Q3 and use the differential equation satisfied by  $\wp$  to eliminate  $\wp''$ . Note also that  $\wp = a - \chi'$ , for some constant  $a \in \mathbb{C}$  (can you see why?).]

5.\* Elliptic functions may be thought of as generalizations of trigonometric functions. To make this more precise, consider  $\psi(z, it)$  for t > 0. Show that for each fixed z,

$$\exp(\pi t/4)\psi(z,it) \to -2\sin(\pi z), \text{ as } t \to \infty.$$

This suggests the replacement

$$\begin{split} \psi(z,it) & \text{by } \psi_{\infty}(z) = -2\sin \pi z, \\ \chi(z,it) & \text{by } \chi_{\infty}(z) = \psi'_{\infty}(z)/\psi_{\infty}(z) = \pi \cot \pi z, \\ \wp(z,it) & \text{by } \wp_{\infty}(z) = \text{const} - \chi'_{\infty}(z) = \text{const} + \pi^2/\sin^2 \pi z. \end{split}$$

Verify that in order that  $\wp_{\infty}(z) = 1/z^2 + z^2 \cdot \text{(holomorphic function near zero)},$ we must have  $\wp_{\infty}(z) = \frac{\pi^2}{\sin^2 \pi z} - \frac{\pi^2}{3}.$ 

Verify also that  $\wp_{\infty}$  satisfies the differential equation for  $\wp$  for suitable values of  $E_4$  and  $E_6$  (find these values!).

6. Show that any holomorphic map f of degree 2 from an elliptic curve  $\mathbb{C}/\Lambda$  to  $S^2$  is given by a 'Möbius transformation of a shifted  $\wp$ -function':

$$f(z) = \frac{a\wp(z-z_0) + b}{c\wp(z-z_0) + d},$$

for some  $a, b, c, d, z_0 \in \mathbb{C}$ .

7. Show, by considering the unit disc  $\Delta$  and the complex plane  $\mathbb{C}$ , that homeomorphic Riemann surfaces need not be conformally equivalent (biholomorphic).

Show that no two of the following domains in  $\mathbb C$  are conformally equivalent

$$\{1 < |z| < 2\}, \qquad \{0 < |z| < 1\}, \qquad \{0 < |z| < \infty\}.$$

8. (i) Let R and S be some Riemann surfaces,  $f : R \to S$  a continuous map, and p a point in R. Show, directly from the definition of holomorphic maps, that if f is holomorphic on  $R \setminus \{p\}$  then f is in fact holomorphic on all of R.

(ii) Suppose that each of  $A = \{\alpha_1, \alpha_2, \alpha_2, \alpha_4\}$  and  $B = \{\beta_1, \beta_2, \beta_3, \beta_4\}$  is a set of four distinct points in  $S^2$  and  $F : S^2 \setminus A \to S^2 \setminus B$  is a biholomorphic map. Show that F extends to a biholomorphic map of  $S^2$  onto itself, hence the  $\beta_4$  is determined by the other  $\beta_i$ 's and  $\alpha_j$ 's.

**9.** Show that if R and S are Riemann surfaces such that both are connected, R is compact and S is **non-compact** then every holomorphic map  $f : R \to S$  is constant.

**10.** (i) Let R and S be compact connected Riemann surfaces and  $g: R \to S$  a non-constant holomorphic map. Show that the genus of R is greater or equal to the genus of S.

(ii) Let R and S be compact connected Riemann surfaces, such that

$$\operatorname{genus}(R) = \operatorname{genus}(S) = g.$$

Show that if  $f : R \to S$  is a non-constant holomorphic map and g > 1 then f is biholomorphic. What does the argument give in the case when (a) g = 0 or (b) g = 1?

(iii) Show that a holomorphic map  $f: S^2 \to S^2$  of degree  $k \ge 2$  must have ramification points (i.e. points  $p \in S^2$  with  $v_f(p) > 1$ ); recover from this the answer to Q7 in ex. sheet 1.

11. (i) Let f and g be two elliptic functions (with the same lattice of periods) and N a positive integer. By considering the poles of f and g, estimate from above the dimension of the complex vector space spanned by  $f(z)^m g(z)^n$ , for  $0 \le m, n \le N$ . Deduce that when N is sufficiently large there must be a non-trivial linear dependence,

$$\sum_{m,n=0}^{N} a_{m,n} f(z)^m g(z)^n \equiv 0, \quad \text{ for some } a_{m,n} \in \mathbb{C}.$$

Hence show that any two meromorphic functions f, g on an elliptic curve  $\mathbb{C}/\Lambda$  are **'algebraically related'**: there is a polynomial Q in two variables, so that Q(f(z), g(z)) = 0 for all z.

(ii)\* Verify that the argument of (i) works for meromorphic functions on **any compact** Riemann surface.

**12.** Recall from the Lectures that  $\vartheta(z,\tau) = \sum_{n=-\infty}^{\infty} \mathbf{e}(\frac{1}{2}n^2\tau + nz)$ , where  $\mathbf{e}(z) = \exp(2\pi i z)$ and  $\operatorname{Im}(\tau) > 0$ . Show that if k is a positive integer then  $\vartheta(0,\tau)^k = \sum_{n=0}^{\infty} r_n(k)e^{\pi n i \tau}$ , where  $r_n(k)$  is the number of ways to express the integer n as a sum of k squares.