

### Topics in Analysis: Example Sheet 3

LENT 2007-08

N. WICKRAMASEKERA

(1) For each positive integer  $n$  and  $k \in \{1, 2, \dots, n\}$ , let the non-negative numbers  $A_k^{(n)}$  and the “nodes”  $x_k^{(n)} \in [a, b]$  be given such that for each polynomial  $P$ , the error

$$\epsilon_n(P) \equiv \left| \int_a^b P(x) dx - \sum_{k=1}^n A_k^{(n)} P(x_k^{(n)}) \right|$$

in approximating  $\int_a^b P(x) dx$  by  $\sum_{k=1}^n A_k^{(n)} P(x_k^{(n)})$  tends to zero as  $n \rightarrow \infty$ . Prove that  $\epsilon_n(f) \rightarrow 0$  for each continuous function  $f$  on  $[a, b]$ .

(2) Let  $T_j$  be the  $j$ th Chebychev polynomial. Suppose  $\gamma_j$  is a sequence of non-negative numbers with  $\sum_{j=1}^{\infty} \gamma_j < \infty$ . Prove that  $\sum_{j=1}^{\infty} \gamma_j T_{3^j}$  defines a continuous function  $f$  on  $[-1, 1]$  with the property that for each  $n$ , there exist points  $-1 \leq x_0 < x_1 < \dots < x_{3^{n+1}} \leq 1$  such that, writing  $P_n$  for the partial sum  $\sum_{j=1}^n \gamma_j T_{3^j}$ ,

$$f(x_k) - P_n(x_k) = (-1)^k \sum_{j=n+1}^{\infty} \gamma_j$$

for each  $k = 0, 1, \dots, 3^{n+1}$ .

(3) For each  $f \in C([-1, 1])$ , let  $E_n(f)$  be the distance from  $f$  to the subspace  $\mathcal{P}_n$  of polynomials of degree at most  $n$ . That is,  $E_n(f) = \inf_{p \in \mathcal{P}_n} \sup_{x \in [-1, 1]} |f(x) - p(x)|$ . We know by the Weierstrass approximation theorem that  $E_n(f) \rightarrow 0$  for each  $f \in C([-1, 1])$ . Using the result of problem (2), construct a function  $f \in C([-1, 1])$  to show that the convergence  $E_n(f) \rightarrow 0$  can be arbitrarily slow in the following sense. For any given decreasing sequence of non-negative numbers  $\delta_n$  converging to zero, there exists  $f \in C([-1, 1])$  such that  $E_n(f) \geq \delta_n$  for all  $n = 1, 2, \dots$ .

(4) Let  $q_n$ ,  $n = 0, 1, 2, \dots$ , be a dense set of distinct points in  $[0, 1]$  with  $q_0 = 0$  and  $q_1 = 1$ . Let  $f \in C([0, 1])$ , and  $f_n$ ,  $n = 0, 1, 2, \dots$  be the piecewise linear function with  $f_n(q_j) = f(q_j)$  for  $j = 0, 1, \dots, n$ . Prove that  $f_n \rightarrow f$  uniformly on  $[0, 1]$ .

(5) Use the result of problem (4) to prove that there exists a sequence of functions  $\varphi_n \in C([0, 1])$ ,  $n = 0, 1, 2, \dots$ , such that for every  $f \in C([0, 1])$ , there exists a unique series  $\sum_{n=0}^{\infty} a_n \varphi_n$  which converges uniformly to  $f$ .

(6) Let  $X$  be a compact metric space and  $\mathcal{A}$  an algebra of real valued continuous functions on  $X$  which is dense in  $C(X)$ . Does it follow that (a)  $\mathcal{A}$  separates points? (b)  $\mathcal{A}$  contains the constants?

(7) Give an example of an algebra of complex, continuous functions of a real variable on  $[0, 1]$  which separates points and contains the (complex) constants but is not a dense subset of the space of all complex continuous functions on  $[0, 1]$ .

(8) Let  $\Omega$  be a subset of the complex plane  $\mathbf{C}$  and for each  $n = 1, 2, 3, \dots$ , let  $f_n : \Omega \rightarrow \mathbf{C}$  be functions such that  $f_n$  converge uniformly on  $\Omega$  to a bounded function  $f : \Omega \rightarrow \mathbf{C}$ . Prove that for any fixed positive integer  $m$ ,  $f_n^m \rightarrow f^m$  uniformly.

(9) Let  $B_r(z)$  denote the open ball in the complex plane with radius  $r$  and centre  $z$  and let  $\Omega = B_2(1) \setminus \overline{B_1(0)}$ . Suppose that  $f$  is analytic in  $\Omega$ .

(a) Prove that there exists a sequence of polynomials which converges to  $f$  uniformly on compact subsets of  $\Omega$ .

(b) Must there be a sequence of polynomials which converges to  $f$  uniformly on  $\Omega$ ?

(c) If additionally we assume that  $f$  is analytic in some open set containing the closure of  $\Omega$ , must there be a sequence of polynomials which converges to  $f$  uniformly on  $\Omega$ ?

(10) Construct a sequence of polynomials which converges uniformly to  $1/z$  on the semicircle  $\{z : |z| = 1, \operatorname{Re}(z) \geq 0\}$ .

(11) Let  $C$  be a closed curve in the complex plane  $\mathbf{C}$  parametrized by a  $C^1$  function  $\gamma : [0, 1] \rightarrow \mathbf{C}$ , and  $K$  a compact subset disjoint from  $C$ . Define a function  $f : K \rightarrow \mathbf{C}$  by  $f(z) = \int_C \frac{dw}{w-z}$ . By splitting up the interval  $[0, 1]$  into subintervals, prove that  $f$  can be uniformly approximated on  $K$  by functions of the form  $f_n(z) = \sum_{j=1}^n \frac{A_j}{(w_j - z)}$  where  $w_j \in C$  and  $A_j \in \mathbf{C}$  for each  $j = 1, 2, \dots, n$ .

(12) Does there exist a sequence of polynomials  $P_n$  such that  $P_n(0) = 1$  for every  $n = 1, 2, 3, \dots$  and  $P_n(z) \rightarrow 0$  for every  $z \in \mathbf{C} \setminus \{0\}$ ?

(13) Prove that a real number is algebraic if and only if it is a zero of a polynomial with rational coefficients.

(14) Prove that  $\sqrt{2} + \sqrt{3}$  and  $\cosh 1$  are irrational.

(15) By considering the numbers  $\sum_{n=0}^{\infty} \frac{b_n}{10^{n!}}$ , where  $b_n \in \{1, 2\}$ , give another proof that the set of transcendental numbers is uncountable.