

Topics in Analysis: Example Sheet 3

LENT 2008-09

N. WICKRAMASEKERA

(1) For each positive integer n and $k \in \{1, 2, \dots, n\}$, let the non-negative numbers $A_k^{(n)}$ and the “nodes” $x_k^{(n)} \in [a, b]$ be given such that for each polynomial P , the error

$$\epsilon_n(P) \equiv \left| \int_a^b P(x) dx - \sum_{k=1}^n A_k^{(n)} P(x_k^{(n)}) \right|$$

in approximating $\int_a^b P(x) dx$ by $\sum_{k=1}^n A_k^{(n)} P(x_k^{(n)})$ tends to zero as $n \rightarrow \infty$. Prove that $\epsilon_n(f) \rightarrow 0$ for each continuous function f on $[a, b]$.

(2) Let T_j be the j th Chebychev polynomial. Suppose γ_j is a sequence of non-negative numbers with $\sum_{j=1}^{\infty} \gamma_j < \infty$. Prove that $\sum_{j=1}^{\infty} \gamma_j T_{3^j}$ defines a continuous function f on $[-1, 1]$ with the property that for each n , there exist points $-1 \leq x_0 < x_1 < \dots < x_{3^{n+1}} \leq 1$ such that, writing P_n for the partial sum $\sum_{j=1}^n \gamma_j T_{3^j}$,

$$f(x_k) - P_n(x_k) = (-1)^k \sum_{j=n+1}^{\infty} \gamma_j$$

for each $k = 0, 1, \dots, 3^{n+1}$.

(3) For each $f \in C([-1, 1])$, let $E_n(f)$ be the distance from f to the subspace \mathcal{P}_n of polynomials of degree at most n . That is, $E_n(f) = \inf_{p \in \mathcal{P}_n} \sup_{x \in [-1, 1]} |f(x) - p(x)|$. We know by the Weierstrass approximation theorem that $E_n(f) \rightarrow 0$ for each $f \in C([-1, 1])$. Using the result of problem (2), construct a function $f \in C([-1, 1])$ to show that the convergence $E_n(f) \rightarrow 0$ can be arbitrarily slow in the following sense. For any given decreasing sequence of non-negative numbers δ_n converging to zero, there exists $f \in C([-1, 1])$ such that $E_n(f) \geq \delta_n$ for all $n = 1, 2, \dots$.

(4) Let q_n , $n = 0, 1, 2, \dots$, be a dense set of distinct points in $[0, 1]$ with $q_0 = 0$ and $q_1 = 1$. Let f_n be the piecewise linear function with $f_n(q_j) = f(q_j)$ for $j = 0, 1, \dots, n$. Prove that $f_n \rightarrow f$ uniformly on $[0, 1]$.

(5) Use the result problem (4) to prove that there exists a sequence of functions $\varphi_n \in C([0, 1])$, $n = 0, 1, 2, \dots$, such that for every $f \in C([0, 1])$, there exists a unique series $\sum_{n=0}^{\infty} a_n \varphi_n$ which converges uniformly to f .

(6) Let Ω be a subset of the complex plane \mathbf{C} and for each $n = 1, 2, 3, \dots$, let $f_n : \Omega \rightarrow \mathbf{C}$ be functions such that f_n converge uniformly on Ω to a bounded function $f : \Omega \rightarrow \mathbf{C}$. Prove that for any fixed positive integer m , $f_n^m \rightarrow f^m$ uniformly.

(7) Let $f : [-1, 1] \rightarrow \mathbf{R}$ be a function and n an integer ≥ 0 . Show that there can be at most one polynomial P of degree $\leq n$ such that $|f(x) - P(x)| \leq M|x|^{n+1}$ for some constant $M > 0$ and all $x \in [-1, 1]$.

(8) Let $B_r(z)$ denote the open ball in the complex plane with radius r and centre z and let $\Omega = B_2(1) \setminus \overline{B_1(0)}$. Suppose that f is analytic in Ω .

(a) Prove that there exists a sequence of polynomials which converges to f uniformly on compact subsets of Ω .

(b) Must there be a sequence of polynomials which converges to f uniformly on Ω ?

(c) If additionally we assume that f is analytic in some open set containing the closure of Ω , must there be a sequence of polynomials which converges to f uniformly on Ω ?

(9) Construct a sequence of polynomials which converges uniformly to $1/z$ on the semicircle $\{z : |z| = 1, \operatorname{Re}(z) \geq 0\}$.

(10) Let C be a closed curve in the complex plane \mathbf{C} parametrized by a C^1 function $\gamma : [0, 1] \rightarrow \mathbf{C}$, and K a compact subset disjoint from C . Define a function $f : K \rightarrow \mathbf{C}$ by $f(z) = \int_C \frac{dw}{w-z}$. By splitting up the interval $[0, 1]$ into subintervals, prove that f can be uniformly approximated on K by functions of the form $f_n(z) = \sum_{j=1}^n \frac{A_j}{(w_j-z)}$ where $w_j \in C$ and $A_j \in \mathbf{C}$ for each $j = 1, 2, \dots, n$.

(11) (Another problem on the Weierstrass approximation theorem). (a) If $f : (0, 1) \rightarrow \mathbf{R}$ is continuous, prove that there exists a sequence of polynomials P_n such that $P_n \rightarrow f$ uniformly on compact subsets of $(0, 1)$.

(b) If $f : (0, 1) \rightarrow \mathbf{R}$ is continuous and bounded, prove that there exists a sequence of polynomials Q_n such that Q_n are uniformly bounded and $Q_n \rightarrow f$ uniformly on compact subsets of $(0, 1)$.

(12) Prove that a real number is algebraic if and only if it is a zero of a polynomial with rational coefficients.

(13) Prove that $\sqrt{2} + \sqrt{3}$ and $\cosh 1$ are irrational.

(14) By considering the numbers $\sum_{n=0}^{\infty} \frac{b_n}{10^{n!}}$, where $b_n \in \{1, 2\}$, give a (second) proof that the set of transcendental numbers is uncountable.