

Topics in Analysis: Example Sheet 2

MICHAELMAS 2010-11

N. WICKRAMASEKERA

(1) Does there exist a function $f : [0, 1] \rightarrow \mathbf{R}$ with a discontinuity which can be approximated uniformly on $[0, 1]$ by polynomials?

(2) Let $f : [0, 1] \rightarrow \mathbf{R}$ be a continuous function which is not a polynomial. If p_n is a sequence of polynomials converging uniformly to f on $[0, 1]$, and $d_n = \text{degree of } p_n$, prove that $d_n \rightarrow \infty$.

(3) Use the mean value theorem to prove that if P is a real polynomial of degree at most n which vanishes at $(n + 1)$ distinct real numbers, then P is identically zero.

(4) Suppose $f : [-1, 1] \rightarrow \mathbf{R}$ is $(n + 1)$ -times continuously differentiable on $[-1, 1]$ and let $J_n = \{x_0, x_1, \dots, x_n\}$ be a set of $(n + 1)$ distinct points in $[-1, 1]$. Let P_{J_n} be the interpolating polynomial of degree $\leq n$ determined by the requirement $P_{J_n}(x_j) = f(x_j)$ for each $j = 0, 1, 2, \dots, n$. Let $\beta_{J_n}(x) = (x - x_0)(x - x_1)(x - x_2) \dots (x - x_n)$. Prove that for each $x \in [-1, 1]$, there exists $\zeta \in (-1, 1)$ such that

$$f(x) - P_{J_n}(x) = \frac{f^{(n+1)}(\zeta)}{(n+1)!} \beta_{J_n}(x).$$

(Hint: If $x = x_j$ this holds trivially. If not, consider $g(y) = f(y) - P_{J_n}(y) - \lambda \beta_{J_n}(y)$ where λ is chosen so that $g(x) = 0$.)

Deduce that if f is infinitely differentiable in $[-1, 1]$ and $\sup_{x \in [-1, 1]} |f^{(n)}(x)| \leq M^n$ for some fixed constant M and all $n = 1, 2, 3, \dots$, then the interpolating polynomials P_{J_n} (for arbitrary choices of sets of interpolation points $J_n = \{x_0^{(n)}, \dots, x_n^{(n)}\} \subset [-1, 1]$) converge uniformly to f on $[-1, 1]$ as $n \rightarrow \infty$.

(5) Fix $n \geq 1$ and let J be any set of n distinct points $\{x_1, \dots, x_n\} \subset [-1, 1]$. Let β_J be the polynomial defined by $\beta_J(x) = (x - x_1)(x - x_2) \dots (x - x_n)$ and set

$$F(x_1, \dots, x_n) = \sup_{x \in [-1, 1]} |\beta_J(x)|.$$

By considering the n th Chebychev polynomial or otherwise, prove that F is minimized when $x_k = \cos\left(\frac{(2k-1)\pi}{2n}\right)$ for $k = 1, 2, \dots, n$.

(6) It can be shown that the converse of the equal ripple criterion holds. That is to say, if $f \in C([0, 1])$ and p is a polynomial of degree at most n which minimizes $\|f - q\|_\infty = \sup_{x \in [0, 1]} |f(x) - q(x)|$ among all polynomials q of degree at most n , then there exist $(n + 2)$ distinct points $0 \leq x_1 < x_2 < \dots < x_{n+2} \leq 1$ such that either $f(x_j) - p(x_j) = (-1)^j \|f - p\|_\infty$ for all $j = 1, 2, \dots, n + 2$ or $f(x_j) - p(x_j) = (-1)^{j+1} \|f - p\|_\infty$ for all $j = 1, 2, \dots, n + 2$. Assuming this, prove that for any given $f \in C([0, 1])$ and each positive integer n , the minimizer of $\|f - q\|_\infty$ among all polynomials q of degree at most n is unique. (Recall that the existence of such a minimizer was proved in lecture.)

(7) Determine all linear operators $L : C([0, 1]) \rightarrow C([0, 1])$ which satisfy (i) $Lf \geq 0$ for all non-negative $f \in C([0, 1])$ and (ii) $Lf = f$ for the three functions $f(x) = 1, x, x^2$.

(8) If $f : \mathbf{R} \rightarrow \mathbf{R}$ is continuous, show that there exist polynomials p_n , $n = 1, 2, \dots$, such that $p_n(x) \rightarrow f(x)$ for every $x \in \mathbf{R}$.

(9) Let $B_n : C([0, 1]) \rightarrow C([0, 1])$ be the Bernstein operator defined by

$$B_n f(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}.$$

Show directly that $B_n f \rightarrow f$ uniformly on $[0, 1]$ for the function $f(x) = x^3$.

(10) Give a proof of the Weierstrass approximation theorem by completing the following argument: Let $0 < a < b < 1$, and $f : [a, b] \rightarrow \mathbf{R}$ be the continuous function we wish to approximate uniformly on $[a, b]$ by polynomials. Fix any continuous extension of f to all of \mathbf{R} such that the extended function is identical to zero outside $[0, 1]$, and denote it again by f .

(a) For each $\delta \in (0, 1/2)$ and each $n = 1, 2, 3, \dots$, set $I_n = \int_0^1 (1-t^2)^n dt$ and $I_{n,\delta} = \int_\delta^1 (1-t^2)^n dt$. Show that $I_n > (1+n)^{-1}$ and $I_{n,\delta} < (1-\delta^2)^n$. Thus, for any fixed $\delta \in (0, 1/2)$, $I_{n,\delta}/I_n \rightarrow 0$ as $n \rightarrow \infty$.

(b) Choose numbers a_1, b_1 such that $0 < a_1 < a < b < b_1 < 1$, and set, for $x \in \mathbf{R}$ and $n = 1, 2, 3, \dots$,

$$\tilde{p}_n(x) = \int_{a_1}^{b_1} f(y)(1-(y-x)^2)^n dy.$$

Given any $\epsilon > 0$, there exists $\delta > 0$ such that $|f(x+t) - f(x)| < \epsilon$ for $|t| < \delta$ and any x . Why? Use this fact and a change of variables in the integral above to show that for $x \in [a, b]$,

$$\tilde{p}_n(x) = 2f(x)(I_n - I_{n,\delta}) + R_n(x)$$

where $|R_n(x)| \leq 2\epsilon I_n + 2MI_{n,\delta}$.

(c) Set $p_n = (2I_n)^{-1}\tilde{p}_n$. Check that p_n is a polynomial of degree $\leq 2n$, and that $\sup_{x \in [a,b]} |p_n(x) - f(x)| < 2\epsilon$ for all sufficiently large n .

(11) Calculate the first five Chebychev polynomials.

(12) Calculate the first four Legendre polynomials. Do it both using the formula and using orthogonality and check that your answers agree.

(13) If $f \in C([0, 1])$ and $\int_0^1 f(x)x^n dx = 0$ for all $n = 0, 1, 2, \dots$, prove that f is the zero function. If we only assume that $f \in C([0, 1])$ and $\int_0^1 f(x)x^n dx = 0$ for all $n = 1, 2, \dots$, does it still follow that f is the zero function?

(14) Let $a, b \in \mathbf{R}$ with $a < b$ and let n be an integer ≥ 1 . Give an explicit expression, in terms of an appropriate Chebychev polynomial, for the polynomial p of degree $\leq n-1$ satisfying

$$\sup_{x \in [a,b]} |x^n - p(x)| \leq \sup_{x \in [a,b]} |x^n - q(x)|$$

for all polynomials q of degree $\leq n-1$.