

## Topics in Analysis: Example Sheet 1

MICHAELMAS 2011-12

N. WICKRAMASEKERA

(1) Let  $X$  be a subset of  $\mathbf{R}^n$ . If every continuous function  $f : X \rightarrow \mathbf{R}$  is bounded, prove that  $X$  is compact.

(2) Consider the metric space  $(\mathbf{Q}, d)$  where  $\mathbf{Q}$  is the set of rational numbers and  $d$  is the usual Euclidean metric. Let  $P$  be the set of all  $p \in \mathbf{Q}$  such that  $2 < p^2 < 3$ . Prove that  $P$  is closed and bounded in  $\mathbf{Q}$ , but not compact. Give an open cover of  $P$  which has no finite subcover.

(3) Find a metric  $d$  on  $X = (0, 1]$  such that  $(X, d)$  is a complete metric space, and such that a subset of  $X$  is open with respect to  $d$  if and only if it is open with respect to the usual Euclidean metric.

(4) Let  $I_n$ ,  $n = 1, 2, 3, \dots$ , be closed, bounded intervals in  $\mathbf{R}$  such that  $I_n \supset I_{n+1}$  for each  $n = 1, 2, 3, \dots$ . Prove that  $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ . State and prove a generalization of this to  $\mathbf{R}^k$ .

(5) A family  $\mathcal{F}$  of real valued functions defined on a subset  $E$  of a metric space  $(X, d)$  is said to be *equicontinuous* on  $E$  if for each  $\epsilon > 0$ , there is a  $\delta > 0$  such that

$$x, y \in E, \quad d(x, y) < \delta, \quad f \in \mathcal{F} \quad \implies \quad |f(x) - f(y)| < \epsilon.$$

Prove that if  $K$  is a compact subset of  $C([0, 1])$  (with the uniform metric), then  $K$  is equicontinuous. (Hint: argue by contradiction.)

(6) Let  $f : B \rightarrow B$  be a continuous map from the *open* disc  $B = \{(x, y) \in \mathbf{R}^2 : x^2 + y^2 < 1\}$  into itself. Does it follow that  $f$  has a fixed point? Does your answer change if  $f$  is a continuous bijection?

(7) Suppose that  $A$  is a  $3 \times 3$  matrix with positive entries. By considering a suitable map from the triangle  $T = \{\mathbf{x} \in \mathbf{R}^3 : x_1, x_2, x_3 \geq 0, x_1 + x_2 + x_3 = 1\}$  into itself, prove that  $A$  has an eigenvector with positive entries.

(8) Use the Brouwer fixed point theorem to prove that there is a complex number  $z$  with  $|z| \leq 1$  such that  $z^4 - z^3 + 8z^2 + 11z + 1 = 0$ .

(9) Suppose  $(X, d)$  is a compact metric space and  $f : X \rightarrow X$  is a continuous function such that  $d(f(x), f(y)) \geq d(x, y)$  for all  $x, y \in X$ . Prove that  $f$  is a surjection. (Hint: if not show that there is a point  $y \in X \setminus f(X)$  and  $r > 0$  such that  $B_r(y) \subset X \setminus f(X)$ , and consider the sequence  $y, f(y), f(f(y)), \dots$ )

(10) Recall that a function  $f : X \rightarrow Y$  between metric spaces is uniformly continuous if for every  $\epsilon > 0$ , there exists a  $\delta > 0$  such that  $x, y \in X$ ,  $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$ . If  $(X, d_X)$  is compact, prove that every continuous function  $f : X \rightarrow Y$  is uniformly continuous. Give in fact two proofs; a direct proof using the definition of compactness involving open covers, and another proof by contradiction using sequential compactness.

(11) Suppose  $f : \mathbf{R} \rightarrow \mathbf{R}$  has the intermediate value property that if  $a, b \in \mathbf{R}$ ,  $f(a) < c < f(b)$ , then  $f(y) = c$  for some  $y$  between  $a$  and  $b$ .

(i) Show that  $f$  need not be continuous.

(ii) If additionally  $f$  has the property that  $f^{-1}(\{a\})$  is closed for every  $a$  in a dense subset of  $\mathbf{R}$ , show that  $f$  is continuous.

(12) Let  $D$  be the closed unit disc in  $\mathbf{R}^2$ ,  $x_0 \in D$  and suppose that  $f : D \rightarrow D$  is a continuous function such that  $f(x_0) \neq x_0$ .

(i) Show that there is  $r > 0$  such that  $f(x) \neq x$  for all  $x \in B_r(x_0)$ .

(ii) For  $x \in B_r(x_0)$ , define  $g(x)$  to be the point where  $\partial D$  meets the ray obtained by extending the line segment connecting  $f(x)$  to  $x$  in the direction from  $f(x)$  to  $x$ . Prove that  $g : B_r(x_0) \rightarrow \partial D$  is continuous at  $x_0$ .

(Hint: note that any point on the ray in question can be written as  $f(x) + \tau(x - f(x))$  for some  $\tau \geq 0$ , and for each  $x \in D$ , the requirement that this point belongs to  $\partial D$  determines uniquely a non-negative  $\tau = \tau(x)$ .)

(13) Let  $q$  be an integer  $\geq 2$  and  $n$  an integer  $\geq 1$ . Denote by  $\mathcal{Q}$  the set of all unordered  $q$ -tuples of points in  $\mathbf{R}^n$ . Thus  $\mathcal{Q} = \{\{x_1, x_2, \dots, x_q\} : x_1, x_2, \dots, x_q \text{ are not necessarily distinct points in } \mathbf{R}^n\}$ . Define a function  $\mathcal{G} : \mathcal{Q} \times \mathcal{Q} \rightarrow \mathbf{R}$  by

$$\mathcal{G}(\{x_1, x_2, \dots, x_q\}, \{y_1, y_2, \dots, y_q\}) = \inf \left\{ \left( \sum_{j=1}^q |x_j - x_{\sigma(j)}|^2 \right)^{1/2} : \sigma \text{ is a permutation of } 1, 2, \dots, q \right\}.$$

(a) Show that  $\mathcal{G}$  is a metric on  $\mathcal{Q}$ .

(b) Assume  $n = 1$ . Note that in this case, we may, for any point  $x = \{x_1, x_2, \dots, x_q\} \in \mathcal{Q}$ , choose the labeling so that  $x_1 \leq x_2 \leq \dots \leq x_q$ . Assuming we have done this for all points in  $\mathcal{Q}$ , is it true that  $\mathcal{G}(\{x_1, x_2, \dots, x_q\}, \{y_1, y_2, \dots, y_q\}) = \left( \sum_{j=1}^q (x_j - y_j)^2 \right)^{1/2}$ ?

(14) (Optional). If  $f_n : [0, 1] \rightarrow \mathbf{R}$ ,  $f_n(0) = 0$  and  $|f_n(x) - f_n(y)| \leq |x - y| + \frac{1}{n}$  for each  $x, y \in [0, 1]$  and  $n = 1, 2, \dots$ , prove that there is a subsequence  $\{f_{n_k}\}$  of  $\{f_n\}$  such that  $\{f_{n_k}\}$  converges uniformly on  $[0, 1]$  to a continuous function.