

## Introduction to non linear Analysis- Weak convergence and Sobolev spaces

Exercises to be done 2-3-4-6-7.

**Exercise 1** (Fundamental solution of Helmholtz). Let  $d \geq 1$ .

1. Let  $f \in \mathcal{S}(\mathbb{R}^d)$  with spherical symmetry, show that its Fourier transform also has spherical symmetry
2. Let  $\lambda > 0$  and  $E : \mathbb{R}^3 \rightarrow \mathbb{R}$ ,  $x \mapsto \frac{e^{-\sqrt{\lambda}\|x\|}}{\|x\|}$ , show that  $E \in \mathcal{S}'(\mathbb{R}^3)$  and compute its Fourier transform.
3. Given  $f \in \mathcal{S}(\mathbb{R}^3)$ ,  $\lambda > 0$ , solve the Helmholtz equation in  $\mathcal{S}'(\mathbb{R}^3)$

$$(-\Delta + \lambda)u = f$$

and give the representation formula both in Fourier and space variables. Show that for all  $s \in \mathbb{R}$ , the map  $f \mapsto u$  is continuous from  $H^s$  into  $H^{s+2}$ .

**Exercise 2** (Uniform approximation of  $L^p$  functions). Let  $d \geq 1$ . Let  $1 \leq p < +\infty$ .

1. Show that

$$\forall h \in \mathbb{R}^d, \quad \forall x \in \mathbb{R}^d, \quad \forall f \in \mathcal{D}(\mathbb{R}^d), \quad |f(x+h) - f(x)| \leq c \left[ |h|^p \int_0^1 |\nabla f(x+th)|^p dt \right]^{\frac{1}{p}}.$$

2. Show that

$$\forall h \in \mathbb{R}^d, \quad \forall f \in \mathcal{D}(\mathbb{R}^d), \quad \int_{\mathbb{R}^d} |f(x+h) - f(x)|^p \leq C|h|^p \|\nabla f\|_{L^p(\mathbb{R}^d)}^p.$$

3. Let  $\zeta \in C_c^\infty(\mathbb{R}^d)$  with

$$\text{Supp}(\zeta) \subset \{|x| \leq 1\}, \quad \int_{\mathbb{R}^d} \zeta(x) dx = 1, \quad \zeta(x) \geq 0.$$

Let  $\varepsilon > 0$  and

$$\zeta_\varepsilon(x) = \frac{1}{\varepsilon^d} \zeta\left(\frac{x}{\varepsilon}\right).$$

Show that

$$\|\zeta_\varepsilon * f - f\|_{L^p(\mathbb{R}^d)} \leq C|\varepsilon| \|\nabla f\|_{L^p(\mathbb{R}^d)}.$$

4. Show that  $\limsup_{\varepsilon \rightarrow 0} \sup_{\|f\|_{L^2} \leq 1} \|\zeta_\varepsilon * f - f\|_{L^2} > 0$ .

5. Show that for all  $s > 0$ ,  $\limsup_{\varepsilon \rightarrow 0} \sup_{\|f\|_{H^s} \leq 1} \|\zeta_\varepsilon * f - f\|_{L^2} = 0$ .

**Exercise 3** (Weak convergence). Let  $H$  be a separable Hilbert space,  $V$  a dense subset of  $H$ . Let  $u \in H$  and  $(u_n)_{n \geq 1}$  be a sequence of elements in  $H$ . Show that

$$u_n \rightharpoonup u \quad \text{in } H$$

iff  $(u_n)_{n \geq 1}$  is bounded in  $H$  and

$$\forall v \in V, \quad \lim_{n \rightarrow +\infty} \langle u_n, v \rangle_H = \langle u, v \rangle_H.$$

**Exercise 4** (Separating bubbles). Let  $d \geq 1$ . Let  $\ell \in \mathbb{N}^*$  and  $(V_j)_{1 \leq j \leq \ell}$  be  $\ell$  functions in  $H^1(\mathbb{R}^d, \mathbb{C})$ . Let  $\mathbf{x}^j = (x_n^j)_{n \geq 1}$ ,  $1 \leq j \leq \ell$  be  $\ell$  sequences with

$$\forall j \neq k, \quad |x_n^j - x_n^k| \rightarrow +\infty \quad \text{as } n \rightarrow +\infty.$$

1. Show that

$$\left\| \sum_{j=1}^{\ell} V^j(\cdot - x_n^j) \right\|_{L^2}^2 = \sum_{j=1}^{\ell} \|V^j\|_{L^2}^2 + o(1) \quad \text{as } n \rightarrow +\infty.$$

2. Show that

$$\left\| \sum_{j=1}^{\ell} \nabla V^j(\cdot - x_n^j) \right\|_{L^2}^2 = \sum_{j=1}^{\ell} \|\nabla V^j\|_{L^2}^2 + o(1) \quad \text{as } n \rightarrow +\infty.$$

3. Let  $1 < p < +\infty$ . Show that there exists a universal constant  $C_{p,\ell} > 0$  such that for all complex numbers  $(z_j)_{1 \leq j \leq \ell}$ ,

$$\left| \left| \sum_{j=1}^{\ell} z_j \right|^p - \sum_{j=1}^{\ell} |z_j|^p \right| \leq C_{p,\ell} \sum_{j \neq k} |z_j| |z_k|^{p-1}.$$

4. Let  $1 < p < +\infty$ . Show that if  $V_j \in L^p(\mathbb{R}^d)$ , then

$$\left\| \sum_{j=1}^{\ell} V^j(\cdot - x_n^j) \right\|_{L^p}^p = \sum_{j=1}^{\ell} \|V^j\|_{L^p}^p + o(1) \quad \text{as } n \rightarrow +\infty.$$

**Exercice 5** (The trace map). . We define the trace map from  $\mathcal{S}(\mathbb{R}^d)$  to  $\mathcal{S}(\mathbb{R}^{d-1})$  by

$$\tau u(x') = u(0, x'), \quad x' = (x_2, \dots, x_d).$$

1. Show that for all  $u \in \mathcal{S}(\mathbb{R}^d)$  and  $\xi' \in \mathbb{R}^{d-1}$ ,

$$\widehat{\tau u}(\xi') = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{u}(\xi_1, \xi') d\xi_1.$$

2. Show that for  $s > 1/2$ ,  $\exists C(s) > 0$  such that  $\forall u \in \mathcal{S}(\mathbb{R}^d)$ ,

$$\|\tau u\|_{H^{s-1/2}(\mathbb{R}^{d-1})} \leq C \|u\|_{H^s(\mathbb{R}^d)}.$$

Hint : use the previous question to derive the estimate

$$|\widehat{\tau u}(\xi')|^2 \leq \frac{1}{4\pi^2} \left( \int_{\mathbb{R}} |\widehat{u}(\xi)|^2 \langle \xi \rangle^{2s} d\xi_1 \right) \left( \int_{\mathbb{R}} \langle \xi \rangle^{-2s} d\xi_1 \right)$$

and express  $\int_{\mathbb{R}} \langle \xi \rangle^{-2s} d\xi_1$  in terms of  $\langle \xi' \rangle$  (where we noted  $\xi = (\xi_1, \xi')$ ).

3. Let  $s > 1/2$ . Show that the trace application extends uniquely as a continuous map from  $H^s(\mathbb{R}^d)$  onto  $H^{s-1/2}(\mathbb{R}^{d-1})$ .

4. Let  $s > 1/2$  and  $g \in H^{s-1/2}(\mathbb{R}^{d-1})$ . Define

$$\widehat{v}(\xi) = \widehat{g}(\xi') \frac{\langle \xi' \rangle^{2(s-1/2)}}{\langle \xi \rangle^{2s}}.$$

Show that  $v \in H^s(\mathbb{R}^d)$  and  $v(0, x') = Cg(x')$  for some constant  $C \neq 0$ . Conclude that the above trace map is surjective.

**Exercice 6** (Space formulation of the homogeneous Sobolev norm). . Let  $0 < s < 1$ . Show that there exists  $0 < c_1 < c_2$  such that for all  $u \in H^s(\mathbb{R}^d)$ , let

$$I_s(u) = \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|u(x+y) - u(x)|^2}{|y|^{d+2s}} dx dy < \infty$$

then

$$c_1 \|u\|_{H^s}^2 \leq I_s(u) \leq c_2 \|u\|_{H^s}^2.$$

Hint : use Plancherel and Fubini.

**Exercice 7** (A commutator estimate). Let  $\chi \in \mathcal{D}(\mathbb{R}^d)$  and  $s \in [0, 1]$ . Let the Fourier multiplier  $\widehat{|D|^s v} \equiv |\xi|^s \widehat{v}$ , and define the commutator

$$A_s v = [|D|^s, \chi] \equiv |D|^s(\chi v) - \chi |D|^s v.$$

1. Let  $v \in \mathcal{D}(\mathbb{R}^d)$ , compute  $\widehat{A_s v}$  in the form of an integral operator ie  $\widehat{A_s v}(\xi) = \int K(\xi, \xi') \widehat{v}(\xi') d\xi'$ .
2. Show that  $A_s$  is bounded on  $L^2(\mathbb{R}^d)$ .