

Introduction to non linear Analysis

Example sheet n° 2 - Weak convergence and linear dispersion

Exercices to be done : 1-2-3.

Exercice 1 (A precised Gagliardo-Nirenberg inequality). Let $d \geq 1$. Let $2 < p < 2^*$ with

$$2^* = \begin{cases} +\infty & \text{for } d = 1, 2 \\ \frac{2d}{d-2} & \text{for } d \geq 3. \end{cases}$$

1. Let a sequence $u_n \in H^1(\mathbb{R}^d)$ with $u_n \rightarrow 0$ in $H^1(\mathbb{R}^d)$, show that $u_n \rightarrow 0$ in $L^p(\mathbb{R}^d)$.
2. Let $\phi \in H^1(\mathbb{R}^d)$ and $x_n \in \mathbb{R}^d$ with $\lim_{n \rightarrow +\infty} |x_n| = +\infty$, let $u_n(x) = \phi(x - x_n)$. Show that $u_n \rightharpoonup 0$ in $H^1(\mathbb{R}^d)$ as $n \rightarrow +\infty$. Do we have $u_n \rightarrow 0$ in $L^p(\mathbb{R}^d)$?
3. Let $\mathbf{u} = (u_n)_{n \geq 1}$ be a bounded sequence in H^1 and $\mathcal{V}(\mathbf{u})$ be the subset of all possible weak H^1 limit of the translates of u_n : $V \in \mathcal{V}(\mathbf{u})$ iff there exists a subsequence $\phi(n)$ and $x_n \in \mathbb{R}^d$ such that

$$u_{\phi(n)}(\cdot - x_n) \rightharpoonup V \text{ in } H^1(\mathbb{R}^d).$$

Show that $\mathcal{V}(\mathbf{u})$ is a bounded subset of $H^1(\mathbb{R}^d)$. We therefore define

$$\eta(\mathbf{u}) = \sup_{V \in \mathcal{V}(\mathbf{u})} \|V\|_{H^1}.$$

4. Let $f \in H^1(\mathbb{R}^d)$ and $u_n = \frac{1}{n^{\frac{d}{2}}} f\left(\frac{x}{n}\right)$. Compute $\eta(\mathbf{u})$. Show that u_n has a limit in L^p for $2 < p < 2^*$. Does u_n converge to 0 in L^2 ?
5. We now assume that

$$\eta(\mathbf{u}) = 0.$$

Our aim is to show the compactness statement

$$u_n \rightarrow 0 \text{ in } L^p \text{ for } 2 < p < 2^*.$$

Let us fix $\chi \in \mathcal{S}(\mathbb{R}^d)$ with

$$\widehat{\chi}(\xi) = \begin{cases} 1 & \text{for } |\xi| \leq 1 \\ 0 & \text{for } |\xi| \geq 2 \end{cases}, \quad |\widehat{\chi}| \leq 1,$$

and given $R > 0$, let

$$\chi_R(x) = R^d \chi(Rx).$$

Let the low-high frequency splitting

$$u_n = u_n^{(1)} + u_n^{(2)}, \quad \begin{cases} \widehat{u_n^{(1)}} = \widehat{u_n} \widehat{\chi_R} \\ \widehat{u_n^{(2)}} = \widehat{u_n} (1 - \widehat{\chi_R}) \end{cases}$$

Let s be given by $-s + \frac{d}{2} = \frac{d}{p}$. Show that $0 < s < 1$.

6. Show that there exists $C_{d,p,\chi} > 0$ such that

$$\forall n \geq 1, \quad \forall R > 0, \quad \|u_n^{(2)}\|_{L^p} \leq \frac{C_{d,p,\chi}}{R^{1-s}}.$$

7. Show that there exists $C_{d,p,\chi} > 0$ such that

$$\forall n \geq 1, \quad \forall R > 0, \quad \|u_n^{(1)}\|_{L^p} \leq C_{d,p,\chi} \|\chi_R \star u_n\|_{L^\infty}^{1-\frac{2}{p}}.$$

8. Show that

$$\forall R > 0, \lim_{n \rightarrow +\infty} \|\chi_R * u_n\|_{L^\infty} = 0.$$

9. Show that $u_n \rightarrow 0$ in $L^p(\mathbb{R}^d)$.

Exercice 2 (Cauchy problem for (NLS) in \mathbb{R}). Let $S(t)$ be the linear Schrödinger semi group on \mathbb{R} . Show that there exists $\alpha_p > 0$ and $C_p > 0$, such that the following holds : for all $u_0 \in H^1(\mathbb{R})$, let $T = \frac{C_p}{\|u_0\|_{H^1}^\alpha}$, then the map

$$\Phi(u)(t, x) = S(t)u_0 + \int_0^t S(t-s)(u|u|^2(s, \cdot))ds$$

is a contraction mapping in a suitable ball of the Banach space $E = L_{[0, T]}^\infty H_x^1$ equipped with the norm $\|u\|_E = \sup_{t \in [0, T]} \|u(t, \cdot)\|_{H_x^1}$.

Exercice 3 (Dispersion for the free transport). Let the transport equation describing the evolution of the microscopic density $f(t, x, v) \in \mathbb{R}^+$ of free particules which are at $x \in \mathbb{R}^d$ with the speed $v \in \mathbb{R}^d$ at time $t \in \mathbb{R}$:

$$(T) \quad \begin{cases} \partial_t f + v \cdot \nabla_x f = 0, \\ f|_{t=0} = f_0. \end{cases}$$

1. Assume $f_0 = f_0(x, v)$ is differentiable, compute the solution to (T).

2. If f_0 is moreover integrable, show that the total density is converved

$$\int_{\mathbb{R}^d \times \mathbb{R}^d} f(t, x, v) dx dv = \int_{\mathbb{R}^d \times \mathbb{R}^d} f_0(x, v) dx dv.$$

3. We define the macroscopic density $\rho(t, x) \stackrel{\text{def}}{=} \int_{\mathbb{R}^d} f(t, x, v) dv$. Show the pointwise decay :

$$\|\rho(t, \cdot)\|_{L^\infty} \leq \frac{1}{|t|^d} \|\sup_v f_0(\cdot, v)\|_{L^1} \quad \text{for all } t \neq 0.$$

Exercice 4 (Wave equation). Let the free wave equation

$$(W) \quad \begin{cases} \square u = 0 \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

where $\square \stackrel{\text{def}}{=} \partial_t^2 - \Delta$ and where $u = u(t, x) \in \mathbb{R}$, $(t, x) \in \mathbb{R} \times \mathbb{R}^d$.

1. For $d = 1$ and $(u_0, u_1) \in C^2 \times C^1$, show that the C^2 solution is given by d'Alembert's formula :

$$u(t, x) = \frac{1}{2} \left(u_0(x+t) + u_0(x-t) + \int_{x-t}^{x+t} u_1(y) dy \right).$$

Do we have pointwise decay in time ?

2. For $d = 3$, we recall that the solution is given by

$$u(t, x) = \frac{1}{4\pi} \left(\frac{1}{t} \int_{S(x, t)} u_1(\sigma) d\sigma + \frac{d}{dt} \left(\frac{1}{t} \int_{S(x, t)} u_0(\sigma) d\sigma \right) \right)$$

wher $S(x, t)$ is the sphere of center x and radius t . Assume for simplicity $u_0 \equiv 0$, then show :

$$\|u(t)\|_{L^\infty} \leq C \frac{\|\nabla u_1\|_{L^1}}{|t|} + \frac{\|u_1\|_{L^1}}{t^2}.$$

Exercice 5 (Oscillatory integrals). Let $a \in \mathcal{D}(\mathbb{R})$ and Φ a C^2 function such that for some $c_0 > 0$:

$$\forall x \in \text{Supp } a, \Phi''(x) \geq c_0.$$

For $t \in \mathbb{R}$, we define the oscillatory integral

$$I(t) \stackrel{\text{def}}{=} \int_{\mathbb{R}} e^{it\Phi(x)} a(x) dx.$$

For $t \neq 0$, we define the differential operator \mathcal{L}_t acting on derivable functions b by

$$\mathcal{L}_t b(x) \stackrel{\text{def}}{=} \frac{1}{1+t(\Phi'(x))^2} (b(x) - i\Phi'(x)b'(x)).$$

1. Using \mathcal{L}_t , show that $I(t) = I_1(t) + I_2(t)$ with

$$\begin{aligned} I_1(t) &\stackrel{\text{def}}{=} \int e^{it\Phi(x)} \frac{i\Phi'(x)}{1+t(\Phi'(x))^2} a'(x) dx \quad \text{and} \\ I_2(t) &\stackrel{\text{def}}{=} \int \frac{e^{it\Phi(x)}}{1+t(\Phi'(x))^2} \left(1 + i\Phi''(x) - 2i \frac{t(\Phi'(x))^2 \Phi''(x)}{1+t(\Phi'(x))^2}\right) a(x) dx. \end{aligned}$$

2. Noticing that for $x \in \text{Supp } a$,

$$\frac{1}{1+t(\Phi'(x))^2} \leq \frac{1}{c_0} \frac{\Phi''(x)}{1+t(\Phi'(x))^2},$$

show that

$$|I_2(t)| \leq \frac{\pi}{2} \left(\frac{1}{c_0} + 3 \right) \frac{1}{|t|^{\frac{1}{2}}} \|a'\|_{L^1(\mathbb{R})}.$$

3. Conclude that there exists $C_0(c_0)$ such that

$$|I(t)| \leq \frac{C_0}{|t|^{\frac{1}{2}}} \|a'\|_{L^1}.$$

4. Application : Consider the Airy equation

$$\partial_t u + \partial_{xxx}^3 u = 0$$

with data u_0 integrable and with Fourier transform supported in

$$[-2, -1/2] \cup [1/2, 2].$$

(a) Show that the L^2 norm is conserved. Write $u(t) = k_t \star u_0$ for a suitable function k_t and conclude

$$\|u(t)\|_{L^\infty} \leq C|t|^{-\frac{1}{2}} \|u_0\|_{L^1}.$$

(b) What kind of L^p - $L^{p'}$ estimate do we obtain if \widehat{u}_0 is supported in the set $[-2\lambda, -\lambda/2] \cup [\lambda/2, 2\lambda]$?

Hint : use the fact that if φ is smooth with support in $\{\frac{1}{3} \leq |\xi| \leq 3\}$ and equal to 1 on $\{\frac{1}{2} \leq |\xi| \leq 2\}$, then $\widehat{u}_0 = \varphi \widehat{u}_0$.