

**SUPPLEMENT TO THE PAPER  
"ON EXISTENCE OF LOG MINIMAL MODELS II"**

CAUCHER BIRKAR

ABSTRACT. We prove a stronger version of a termination theorem appeared in the paper "On existence of log minimal models II" [2]. We essentially just get rid of the redundant assumptions so the proof is almost the same as in [2]. However, we give a detailed proof here for future reference.

1. INTRODUCTION

We work over a fixed algebraically closed field  $k$  of characteristic zero. See section 2 of [2] for notation, terminology, and definition and basic properties of log minimal models. The following theorem was proved in [2, Theorem 1.5] under stronger assumptions. However, the proof works with little change under weaker assumptions.

**Theorem 1.1.** *Let  $(X/Z, B + C)$  be a lc pair of dimension  $d$  such that  $K_X + B + C$  is nef/ $Z$ ,  $B, C \geq 0$  and  $C$  is  $\mathbb{R}$ -Cartier. Assume that we are given an LMMP/ $Z$  on  $K_X + B$  with scaling of  $C$  as in Definition 1.2 with  $\lambda_i$  the corresponding numbers, and  $\lambda := \lim \lambda_i$ . Then, the LMMP terminates in the following cases:*

- (i)  $(X/Z, B)$  is  $\mathbb{Q}$ -factorial dlt,  $B \geq H \geq 0$  for some ample/ $Z$   $\mathbb{R}$ -divisor  $H$ ;
- (ii)  $(X/Z, B)$  is  $\mathbb{Q}$ -factorial dlt,  $C \geq H \geq 0$  for some ample/ $Z$   $\mathbb{R}$ -divisor  $H$ , and  $\lambda > 0$ ;
- (iii)  $(X/Z, B + \lambda C)$  has a log minimal model, and  $\lambda \neq \lambda_j$  for any  $j$ .

We should remark that much of the difficulties in the proof of Theorem 1.1 are caused by the presence of non-klt singularities. It is also worth to mention that the above theorem and the corresponding Theorem 1.5 of [2] follow quite different goals. The arguments of [2] in essence do not rely on [4]. In contrast, the above theorem heavily relies on [4]. So, this paper is not intended to replace [2].

A *sequence of log flips/ $Z$  starting with  $(X/Z, B)$*  is a sequence  $X_i \dashrightarrow X_{i+1}/Z_i$  in which  $X_i \rightarrow Z_i \leftarrow X_{i+1}$  is a  $K_{X_i} + B_i$ -flip/ $Z$ ,  $B_i$  is the birational transform of  $B_1$  on  $X_1$ , and  $(X_1/Z, B_1) = (X/Z, B)$ . As usual, here  $X_i \rightarrow Z_i$  is an extremal flipping contraction.

**Definition 1.2** (LMMP with scaling) Let  $(X_1/Z, B_1 + C_1)$  be a lc pair such that  $K_{X_1} + B_1 + C_1$  is nef/ $Z$ ,  $B_1 \geq 0$ , and  $C_1 \geq 0$  is  $\mathbb{R}$ -Cartier. Suppose that either  $K_{X_1} + B_1$  is nef/ $Z$  or there is an extremal ray  $R_1/Z$  such that  $(K_{X_1} + B_1) \cdot R_1 < 0$

---

*Date:* April 26, 2011.

2000 Mathematics Subject Classification: 14E30.

and  $(K_{X_1} + B_1 + \lambda_1 C_1) \cdot R_1 = 0$  where

$$\lambda_1 := \inf\{t \geq 0 \mid K_{X_1} + B_1 + tC_1 \text{ is nef}/Z\}$$

When  $(X_1/Z, B_1)$  is  $\mathbb{Q}$ -factorial dlt, the last sentence follows from [3, 3.1] (the same is true in general for lc pairs by the results of Ambro [1] and Fujino [5, Theorem 1.1 (6)]); however, we do not need this stronger version). If  $R_1$  defines a Mori fibre structure, we stop. Otherwise assume that  $R_1$  gives a divisorial contraction or a log flip  $X_1 \dashrightarrow X_2$ . We can now consider  $(X_2/Z, B_2 + \lambda_1 C_2)$  where  $B_2 + \lambda_1 C_2$  is the birational transform of  $B_1 + \lambda_1 C_1$  and continue. That is, suppose that either  $K_{X_2} + B_2$  is nef/ $Z$  or there is an extremal ray  $R_2/Z$  such that  $(K_{X_2} + B_2) \cdot R_2 < 0$  and  $(K_{X_2} + B_2 + \lambda_2 C_2) \cdot R_2 = 0$  where

$$\lambda_2 := \inf\{t \geq 0 \mid K_{X_2} + B_2 + tC_2 \text{ is nef}/Z\}$$

By continuing this process, we obtain a sequence of numbers  $\lambda_i$  and a special kind of LMMP/ $Z$  which is called the *LMMP/ $Z$  on  $K_{X_1} + B_1$  with scaling of  $C_1$* ; note that it is not unique. This kind of LMMP was first used by Shokurov [7]. When we refer to *termination with scaling* we mean termination of such an LMMP. We usually put  $\lambda = \lim \lambda_i$ .

When we have a lc pair  $(X/Z, B)$ , we can always find an ample/ $Z$   $\mathbb{R}$ -Cartier divisor  $C \geq 0$  such that  $K_X + B + C$  is lc and nef/ $Z$ , so we can run the LMMP/ $Z$  with scaling assuming that all the necessary ingredients exist, e.g. extremal rays, log flips.

## 2. PROOF OF THE THEOREM

**Proof of Theorem 1.1.** *Step 1.* Proof of (i): Since  $H$  is ample/ $Z$ , we can perturb the coefficients of  $B$  hence assume that  $(X/Z, B)$  is klt. If  $\lambda_i < 1$  for some  $i$ , then  $(X/Z, B + \lambda_i C)$  is klt. If  $\lambda_i = 1$  for every  $i$ , then we can perturb  $C$  hence in any case we could assume that  $(X/Z, B + C)$  is klt and then use [4].

Proof of (ii): The LMMP is also an LMMP/ $Z$  on  $K_X + B + \frac{\lambda}{2}C$  with scaling of  $(1 - \frac{\lambda}{2})C$ . We can replace  $B$  with  $B + \frac{\lambda}{2}C$ ,  $C$  with  $(1 - \frac{\lambda}{2})C$ , and  $\lambda_i$  with  $\frac{\lambda_i - \frac{\lambda}{2}}{1 - \frac{\lambda}{2}}$ . After this change, we can assume that  $B \geq \frac{H}{2}$ . Now use (i).

Proof of (iii): Note that if

(\*)  $(X/Z, B)$  is  $\mathbb{Q}$ -factorial dlt, and  $C \geq H \geq 0$  for some ample/ $Z$   $\mathbb{R}$ -divisor  $H$ ,

then, for each  $i$ , there is a klt  $K_X + \Delta \sim_{\mathbb{R}} K_X + B + \lambda_i C/Z$ . We continue the proof *without* assuming (\*) (but we will come back to (\*) in Step 7).

*Step 2.* We can replace  $B$  with  $B + \lambda C$  hence assume that  $\lambda = 0$ . Moreover, we may assume that the LMMP consists of only a sequence  $X_i \dashrightarrow X_{i+1}/Z_i$  of log flips starting with  $(X_1/Z, B_1) = (X/Z, B)$ . Pick  $i$  so that  $\lambda_i > \lambda_{i+1}$ . Thus,  $\text{Supp } C_{i+1}$  does not contain any lc centre of  $(X_{i+1}/Z, B_{i+1} + \lambda_{i+1} C_{i+1})$  because  $(X_{i+1}/Z, B_{i+1} + \lambda_i C_{i+1})$  is lc. Then, by replacing  $(X/Z, B)$  with  $(X_{i+1}/Z, B_{i+1})$  and  $C$  with  $\lambda_{i+1} C_{i+1}$  we may assume that no lc centre of  $(X/Z, B + C)$  is inside  $\text{Supp } C$ . Moreover, since there are

finitely many lc centres of  $(X/Z, B)$ , perhaps after truncating the sequence, we can assume that no lc centre is contracted in the sequence.

By assumptions there is a log minimal model  $(Y/Z, B_Y)$  for  $(X/Z, B)$ . Let  $\phi: X \dashrightarrow Y/Z$  be the corresponding birational map. Since  $K_{X_i} + B_i + \lambda_i C_i$  is nef/ $Z$ , we may add an ample/ $Z$   $\mathbb{R}$ -divisor  $G^i$  so that  $K_{X_i} + B_i + \lambda_i C_i + G^i$  becomes ample/ $Z$ , in particular, it is movable/ $Z$ . We can choose the  $G^i$  so that  $\lim_{i \rightarrow \infty} G^i = 0$  in  $N^1(X_1/Z)$  where  $G^i_1$  is the birational transform of  $G^i$  on  $X_1 = X$ . Therefore,

$$K_X + B \equiv \varinjlim_{i \rightarrow \infty} (K_{X_1} + B_1 + \lambda_i C_1 + G^i_1)/Z$$

which implies that  $K_X + B$  is a limit of movable/ $Z$   $\mathbb{R}$ -divisors.

Let  $f: W \rightarrow X$  and  $g: W \rightarrow Y$  be a common log resolution of  $(X/Z, B + C)$  and  $(Y/Z, B_Y + C_Y)$  where  $C_Y$  is the birational transform of  $C$ . By applying the negativity lemma to  $f$ , we see that

$$E := f^*(K_X + B) - g^*(K_Y + B_Y) = \sum_D a(D, Y, B_Y)D - a(D, X, B)D$$

is effective (cf. [3, Remark 2.6]) where  $D$  runs over the prime divisors on  $W$ . Assume that  $D$  is a component of  $E$ . If  $D$  is not exceptional/ $Y$ , then it must be exceptional/ $X$  otherwise  $a(D, X, B) = a(D, Y, B_Y)$  and  $D$  cannot be a component of  $E$ . On the other hand, if  $D$  is not exceptional/ $Y$  but exceptional/ $X$ , then by definition of log minimal models,  $a(D, X, B) \leq a(D, Y, B_Y) = 0$  hence  $a(D, X, B) = 0$  which again shows that  $D$  cannot be a component of  $E$ . Therefore,  $E$  is exceptional/ $Y$ .

*Step 3.* Let  $B_W$  be the birational transform of  $B$  plus the reduced exceptional divisor of  $f$ , and let  $C_W$  be the birational transform of  $C$  on  $W$ . Pick a sufficiently small  $\delta \geq 0$ . Take a general ample/ $Z$  divisor  $L$  so that  $K_W + B_W + \delta C_W + L$  is dlt and nef/ $Z$ . Since  $(X/Z, B)$  is lc,

$$E' := K_W + B_W - f^*(K_X + B) = \sum_D a(D, X, B)D \geq 0$$

where  $D$  runs over the prime exceptional/ $X$  divisors on  $W$ . So,

$$K_W + B_W + \delta C_W = f^*(K_X + B) + E' + \delta C_W = g^*(K_Y + B_Y) + E + E' + \delta C_W$$

Moreover,  $E'$  is also exceptional/ $Y$  because for any prime divisor  $D$  on  $Y$  which is exceptional/ $X$ ,  $a(D, Y, B_Y) = a(D, X, B) = 0$  hence  $D$  cannot be a component of  $E'$ .

On the other hand, since  $Y$  is  $\mathbb{Q}$ -factorial, there are exceptional/ $Y$   $\mathbb{R}$ -divisors  $F, F'$  on  $W$  such that  $C_W + F \equiv 0/Y$  and  $L + F' \equiv 0/Y$ . Now run the LMMP/ $Y$  on  $K_W + B_W + \delta C_W$  with scaling of  $L$  which is the same as the LMMP/ $Y$  on  $E + E' + \delta C_W$  with scaling of  $L$ . Let  $\lambda'_i$  and  $\lambda' = \lim_{i \rightarrow \infty} \lambda'_i$  be the corresponding numbers. If  $\lambda' > 0$ , then by Step 1 the LMMP terminates since  $L$  is ample/ $Z$ . Since  $W \rightarrow Y$  is birational, the LMMP terminates only when  $\lambda'_i = 0$  for some  $i$  which implies that  $\lambda' = 0$ , a contradiction. Thus,  $\lambda' = 0$ . On some model  $V$  in the process of the LMMP, the

pushdown of  $K_W + B_W + \delta C_W + \lambda'_i L$ , say

$$\begin{aligned} K_V + B_V + \delta C_V + \lambda'_i L_V & \\ & \equiv E_V + E'_V + \delta C_V + \lambda'_i L_V \\ & \equiv E_V + E'_V - \delta F_V - \lambda'_i F'_V / Y \end{aligned}$$

is nef/ $Y$ . Applying the negativity lemma over  $Y$  shows that  $E_V + E'_V - \delta F_V - \lambda'_i F'_V \leq 0$ . But if  $i \gg 0$ , then  $E_V + E'_V \leq 0$  because  $\lambda'_i$  and  $\delta$  are sufficiently small. Therefore,  $E_V = E'_V = 0$  as  $E$  and  $E'$  are effective.

*Step 4.* We prove that  $\phi: X \dashrightarrow Y$  does not contract any divisors. Assume otherwise and let  $D$  be a prime divisor on  $X$  contracted by  $\phi$ . Then  $D^\sim$  the birational transform of  $D$  on  $W$  is a component of  $E$  because by definition of log minimal models  $a(D, X, B) < a(D, Y, B_Y)$ . Now, in step 3 take  $\delta = 0$ . The LMMP contracts  $D^\sim$  since  $D^\sim$  is a component of  $E$  and  $E$  is contracted. But this is not possible because  $K_X + B$  is a limit of movable/ $Z$   $\mathbb{R}$ -divisors and  $D^\sim$  is not a component of  $E'$  so the pushdown of  $K_W + B_W = f^*(K_X + B) + E'$  cannot negatively intersect a general curve on  $D^\sim/Y$ . Thus  $\phi$  does not contract divisors, in particular, any prime divisor on  $W$  which is exceptional/ $Y$  is also exceptional/ $X$ . Though  $\phi$  does not contract divisors but  $\phi^{-1}$  might contract divisors. The prime divisors contracted by  $\phi^{-1}$  appear on  $W$ .

*Step 5.* Now take  $\delta > 0$  in step 3 which is sufficiently small by assumptions. As mentioned, we arrive at a model  $Y' := V$  on which  $E_{Y'} + E'_{Y'} = 0$  where  $E_{Y'}$  and  $E'_{Y'}$  are the birational transforms of  $E$  and  $E'$  on  $Y'$ , respectively. In view of

$$K_{Y'} + B_{Y'} \equiv E_{Y'} + E'_{Y'} = 0/Y$$

we deduce that  $(Y'/Z, B_{Y'})$  is a  $\mathbb{Q}$ -factorial dlt blowup of  $(Y/Z, B_Y)$ . Moreover, by construction  $(Y'/Z, B_{Y'} + \delta C_{Y'})$  is also dlt.

*Step 6.* As in step 3,

$$E'' := K_W + B_W + C_W - f^*(K_X + B + C) = \sum_D a(D, X, B + C)D \geq 0$$

is exceptional/ $X$  where  $D$  runs over the prime exceptional/ $X$  divisors on  $W$ . Since  $K_W + B_W + C_W \equiv E''/X$ , any LMMP/ $X$  with scaling of a suitable ample/ $X$  divisor  $L$  terminates for the same reasons as in step 3. Indeed, if  $\lambda'_i$  and  $\lambda' = \lim_{i \rightarrow \infty} \lambda'_i$  are the corresponding numbers in the LMMP, then we may assume  $\lambda' = 0$  by Step 1; on some model  $V$  in the process of the LMMP, the pushdown of  $K_W + B_W + C_W + \lambda'_i L$ , say  $K_V + B_V + C_V + \lambda'_i L_V$  is nef/ $X$ . But since  $W \rightarrow X$  is birational, there is some  $L' \geq 0$  such that  $L \sim_{\mathbb{R}} -L'/X$  hence  $E''_V - \lambda'_i L'_V$  is nef/ $X$ . Now if  $i \gg 0$ , the negativity lemma implies that  $E''_V = 0$  hence the LMMP terminates.

So, we get a  $\mathbb{Q}$ -factorial dlt blowup  $(X'/Z, B' + C')$  of  $(X/Z, B + C)$  where  $K_{X'} + B'$  is the pullback of  $K_X + B$  and  $C'$  is the pullback of  $C$ . In fact,  $X'$  and  $X$  are isomorphic outside the lc centres of  $(X/Z, B + C)$  because the prime exceptional/ $X$  divisors on  $X'$  are exactly the pushdown of the prime exceptional/ $X$  divisors  $D$  on  $W$  with  $a(D, X, B + C) = 0$ , that is, those which are not components of  $E''$ . Since  $\text{Supp } C$

does not contain any lc centre of  $(X/Z, B + C)$  by step 2,  $(X'/Z, B')$  is a  $\mathbb{Q}$ -factorial dlt blowup of  $(X/Z, B)$  and  $C'$  is just the birational transform of  $C$ . Note that the prime exceptional divisors of  $\phi^{-1}$  are not contracted/ $X'$  since their log discrepancy with respect to  $(X/Z, B)$  are all 0, and so their birational transforms are not components of  $E''$ .

*Step 7.* Remember that  $X_1 = X$ ,  $B_1 = B$ , and  $C_1 = C$ . Similarly, put  $X'_1 := X'$ ,  $B'_1 := B'$ , and  $C'_1 := C'$ . Since  $K_{X_1} + B_1 + \lambda_1 C_1 \equiv 0/Z_1$ ,  $K_{X'_1} + B'_1 + \lambda_1 C'_1 \equiv 0/Z_1$ . Run the LMMP/ $Z_1$  on  $K_{X'_1} + B'_1$  with scaling of some ample/ $Z_1$  divisor (which is automatically also an LMMP with scaling of  $\lambda_1 C'_1$ ). Assume that this LMMP terminates with a log minimal model  $(X'_2/Z_1, B'_2)$ . Since  $(X_2/Z_1, B_2)$  is the lc model of  $(X_1/Z_1, B_1)$  and of  $(X'_1/Z_1, B'_1)$ ,  $X'_2$  maps to  $X_2$  and  $K_{X'_2} + B'_2$  is the pullback of  $K_{X_2} + B_2$ . Thus,  $(X'_2/Z, B'_2)$  is a  $\mathbb{Q}$ -factorial dlt blowup of  $(X_2/Z, B_2)$ . Since  $K_{X'_1} + B'_1 + \lambda_1 C'_1 \equiv 0/Z_1$ ,  $K_{X'_2} + B'_2 + \lambda_1 C'_2 \equiv 0/Z_1$  where  $C'_2$  is the birational transform of  $C'_1$  and actually the pullback of  $C_2$ . We can continue this process: that is use the fact that  $K_{X_2} + B_2 + \lambda_2 C_2 \equiv 0/Z_2$  and  $K_{X'_2} + B'_2 + \lambda_2 C'_2 \equiv 0/Z_2$  to run an LMMP/ $Z_2$  on  $K_{X'_2} + B'_2$ , etc. Therefore, we can lift the original sequence to a sequence in the  $\mathbb{Q}$ -factorial dlt case assuming that the following statement holds for each  $i$ :

(\*\*) some LMMP/ $Z_i$  on  $K_{X'_i} + B'_i$  with scaling of some ample divisor terminates with a log minimal model  $(X'_{i+1}/Z_i, B'_{i+1})$ .

In this paragraph, we show that we can assume that (\*\*) holds. First, note that if  $(X/Z, B + C)$  is klt, then  $X' \rightarrow X$  is a small birational morphism and  $(X'/Z, B' + C')$  is also klt hence (\*\*) holds by [4]. Now assume that (\*) in Step 1 holds. Then, there is a klt  $K_{X_i} + \Delta_i \sim_{\mathbb{R}} K_{X_i} + B_i + \lambda_i C_i/Z$ . Thus,

$$\begin{aligned} K_{X'_i} + B'_i &\sim_{\mathbb{R}} (K_{X'_i} + B'_i) + \epsilon(K_{X'_i} + \Delta'_i) = \\ &(1 + \epsilon)(K_{X'_i} + \frac{1}{1 + \epsilon}B'_i + \frac{\epsilon}{1 + \epsilon}\Delta'_i)/Z_i \end{aligned}$$

where  $K_{X'_i} + \Delta'_i$  is the pullback of  $K_{X_i} + \Delta_i$ . If  $\epsilon > 0$  is sufficiently small, then  $(X'_i/Z, \frac{1}{1 + \epsilon}B'_i + \frac{\epsilon}{1 + \epsilon}\Delta'_i)$  is klt hence (\*\*) again follows from [4] under (\*). Now note that (\*\*) itself is an LMMP under the assumption (\*) bearing in mind that some  $\mathbb{Q}$ -factorial dlt blowup of  $(X_{i+1}/Z_i, B_{i+1})$  (which can be constructed as in Step 6) is a log minimal model of  $(X'_i/Z_i, B'_i)$ . So, we can assume that (\*\*) holds; otherwise we can replace the original sequence with the one in (\*\*) and repeat Steps 2-7 again.

Note that  $Y' \dashrightarrow X'$  does not contract divisors: if  $D$  is a prime divisor on  $Y'$  which is exceptional/ $X'$ , then it is exceptional/ $X$  and so it is exceptional/ $Y$  by step 6; but then  $a(D, Y, B_Y) = 0 = a(D, X, B)$  and again by step 6 such divisors are not contracted/ $X'$ , a contradiction. Thus,  $(Y'/Z, B_{Y'})$  of step 5 is a log birational model of  $(X'/Z, B')$  because  $B_{Y'}$  is the birational transform of  $B'$ . On the other hand, assume that  $D$  is a prime divisor on  $X'$  which is exceptional/ $Y'$ . Since  $X \dashrightarrow Y$  does not contract divisors by step 4,  $D$  is exceptional/ $X$ . In particular,

$a(D, X', B') = a(D, X, B) = 0$ ; in this case  $a(D, Y, B_Y) = a(D, Y', B_{Y'}) > 0$  otherwise  $D$  could not be contracted/ $Y'$  by the LMMP of step 5 which started on  $W$  because the birational transform of  $D$  would not be a component of  $E + E' + \delta C_W$ . So,  $(Y'/Z, B_{Y'})$  is actually a log minimal model of  $(X'/Z, B')$ . Therefore, as in step 4,  $X' \dashrightarrow Y'$  does not contract divisors which implies that  $X'$  and  $Y'$  are isomorphic in codimension one. Now replace the old sequence  $X_i \dashrightarrow X_{i+1}/Z_i$  with the new one constructed in the last paragraph and replace  $(Y/Z, B_Y)$  with  $(Y'/Z, B_{Y'})$ . So, from now on we can assume that  $X, X_i$  and  $Y$  are all isomorphic in codimension one, and that  $(X/Z, B + C)$  is  $\mathbb{Q}$ -factorial dlt. In addition, by step 5, we can also assume that  $(Y/Z, B_Y + \delta C_Y)$  is  $\mathbb{Q}$ -factorial dlt for some  $\delta > 0$ .

*Step 8.* Let  $A \geq 0$  be a reduced divisor on  $W$  whose components are general ample/ $Z$  divisors such that they generate  $N^1(W/Z)$ . By step 6,  $(X_1/Z, B_1 + C_1)$  is obtained by running a specific LMMP on  $K_W + B_W + C_W$ . Every step of this LMMP is also a step of an LMMP on  $K_W + B_W + C_W + \varepsilon A$  for any sufficiently small  $\varepsilon > 0$ , in particular,  $(X_1/Z, B_1 + C_1 + \varepsilon A_1)$  is dlt where  $A_1$  is the birational transform of  $A$ . For similar reasons, we can choose  $\varepsilon$  so that  $(Y/Z, B_Y + \delta C_Y + \varepsilon A_Y)$  is also dlt. On the other hand, by [2, Proposition 3.2], perhaps after replacing  $\delta$  and  $\varepsilon$  with smaller positive numbers, we may assume that if  $0 \leq \delta' \leq \delta$  and  $0 \leq A'_Y \leq \varepsilon A_Y$ , then any LMMP/ $Z$  on  $K_Y + B_Y + \delta' C_Y + A'_Y$ , consists of only a sequence of log flips which are flops with respect to  $(Y/Z, B_Y)$ . Note that  $K_Y + B_Y + \delta' C_Y + A'_Y$  is a limit of movable/ $Z$   $\mathbb{R}$ -divisors for reasons similar to those used in Step 2, so no divisor is contracted by such an LMMP.

*Step 9.* Fix some  $i \gg 0$  so that  $\lambda_i < \delta$ . Then, by [2, Proposition 3.2], there is  $0 < \tau \ll \varepsilon$  such that  $(X_i/Z, B_i + \lambda_i C_i + \tau A_i)$  is dlt and such that if we run the LMMP/ $Z$  on  $K_{X_i} + B_i + \lambda_i C_i + \tau A_i$  with scaling of some ample/ $Z$  divisor, then it will be a sequence of log flips which would be a sequence of flops with respect to  $(X_i/Z, B_i + \lambda_i C_i)$ . Moreover, since the components of  $A_i$  generate  $N^1(X_i/Z)$ , we can assume that there is an ample/ $Z$   $\mathbb{R}$ -divisor  $H \geq 0$  such that  $\tau A_i \equiv H + H'/Z$  where  $H' \geq 0$  and  $(X_i/Z, B_i + \lambda_i C_i + H + H')$  is dlt. Hence the LMMP terminates by (i) and we get a model  $T$  on which both  $K_T + B_T + \lambda_i C_T$  and  $K_T + B_T + \lambda_i C_T + \tau A_T$  are nef/ $Z$ . Again since the components of  $A_T$  generate  $N^1(T/Z)$ , there is  $0 \leq A'_T \leq \tau A_T$  so that  $K_T + B_T + \lambda_i C_T + A'_T$  is ample/ $Z$  and  $\text{Supp } A'_T = \text{Supp } A_T$ . Now run the LMMP/ $Z$  on  $K_Y + B_Y + \lambda_i C_Y + A'_Y$  with scaling of some ample/ $Z$  divisor where  $A'_Y$  is the birational transform of  $A'_T$ . The LMMP terminates for reasons similar to the above and we end up with  $T$  since  $K_T + B_T + \lambda_i C_T + A'_T$  is ample/ $Z$ . Moreover, the LMMP consists of only log flips which are flops with respect to  $(Y/Z, B_Y)$  by Step 8 hence  $K_T + B_T$  will also be nef/ $Z$ . So, by replacing  $Y$  with  $T$  we could assume that  $K_Y + B_Y + \lambda_i C_Y$  is nef/ $Z$ . In particular,  $K_Y + B_Y + \lambda_j C_Y$  is nef/ $Z$  for any  $j \geq i$  since  $\lambda_j \leq \lambda_i$ .

*Step 10.* Pick  $j > i$  so that  $\lambda_j < \lambda_{j-1} \leq \lambda_i$  and let  $r: U \rightarrow X_j$  and  $s: U \rightarrow Y$  be a common resolution. Then, we have

$$\begin{aligned} r^*(K_{X_j} + B_j + \lambda_j C_j) &= s^*(K_Y + B_Y + \lambda_j C_Y) \\ r^*(K_{X_j} + B_j) &\not\geq s^*(K_Y + B_Y) \\ r^*C_j &\not\leq s^*C_Y \end{aligned}$$

where the first equality holds because both  $K_{X_j} + B_j + \lambda_j C_j$  and  $K_Y + B_Y + \lambda_j C_Y$  are nef/ $Z$  and  $X_j$  and  $Y$  are isomorphic in codimension one, the second inequality holds because  $K_Y + B_Y$  is nef/ $Z$  but  $K_{X_j} + B_j$  is not nef/ $Z$ , and the third follows from the other two. Now

$$\begin{aligned} r^*(K_{X_j} + B_j + \lambda_{j-1} C_j) & \\ &= r^*(K_{X_j} + B_j + \lambda_j C_j) + r^*(\lambda_{j-1} - \lambda_j) C_j \\ &\leq s^*(K_Y + B_Y + \lambda_j C_Y) + s^*(\lambda_{j-1} - \lambda_j) C_Y \\ &= s^*(K_Y + B_Y + \lambda_{j-1} C_Y) \end{aligned}$$

However, since  $K_{X_j} + B_j + \lambda_{j-1} C_j$  and  $K_Y + B_Y + \lambda_{j-1} C_Y$  are both nef/ $Z$ , we have

$$r^*(K_{X_j} + B_j + \lambda_{j-1} C_j) = s^*(K_Y + B_Y + \lambda_{j-1} C_Y)$$

This is a contradiction and the sequence of log flips terminates as claimed.  $\square$

#### REFERENCES

- [1] F. Ambro; *Quasi-log varieties*. Tr. Mat. Inst. Steklova 240 (2003), Biratsion. Geom. Linein. Sist. Konechno Porozhden- nye Algebr, 220-239; translation in Proc. Steklov Inst. Math. 2003, no. 1 (240), 214-233.
- [2] C. Birkar; *On existence of log minimal models II*. To appear in "J. Reine Angew Math." arXiv:0907.4170v1.
- [3] C. Birkar; *On existence of log minimal models*. Compositio Math. volume 145 (2009), 1442-1446.
- [4] C. Birkar, P. Cascini, C. Hacon, J. McKernan; *Existence of minimal models for varieties of log general type*. J. Amer. Math. Soc. 23 (2010), 405-468.
- [5] O. Fujino; *Fundamental theorems for the log minimal model program*. arXiv:0909.4445v2.
- [6] V.V. Shokurov; *Prelimiting flips*. Proc. Steklov Inst. Math. 240 (2003), 75-213.
- [7] V.V. Shokurov; *3-fold log flips*. With an appendix in English by Yujiro Kawamata. Russian Acad. Sci. Izv. Math. 40 (1993), no. 1, 95-202.

DPMMS, Centre for Mathematical Sciences,  
 Cambridge University,  
 Wilberforce Road,  
 Cambridge, CB3 0WB,  
 UK  
 email: c.birkar@dpmms.cam.ac.uk

Fondation Sciences Mathématiques de Paris,

IHP, 11 rue Pierre et Marie Curie,  
75005 Paris,  
France