

DIFFERENTIAL GEOMETRY EXAMPLES 3

G.P. Paternain Lent 2005

Comments on and/or corrections to the questions on this sheet are always welcome, and may be e-mailed to me at g.p.paternain@dpms.cam.ac.uk.

1. Let $\alpha : I \rightarrow S$ be a geodesic. Show that if α is a plane curve and $\ddot{\alpha}(t) \neq 0$ for some $t \in I$, then $\dot{\alpha}(t)$ is an eigenvector of the differential of the Gauss map at $\alpha(t)$.

2. Show that if all geodesics of a connected surface are plane curves, then the surface is contained in a plane or a sphere (Hint: use Problem 15 of Example sheet 2).

3. Let $f : S_1 \rightarrow S_2$ be an isometry between two surfaces.

(i) Let $\alpha : I \rightarrow S_1$ be a curve and V a parallel vector field along α . Show that $df_{\alpha(t)}(V(t))$ is a parallel vector field along $f \circ \alpha$.

(ii) Show that f maps geodesics to geodesics.

4. Consider the surface of revolution from Problem 9, Example sheet 2.

(i) Find the equations of the geodesics;

(ii) Establish *Clairaut's relation*: $f^2 \dot{u}$ is constant along geodesics. Show that if θ is the angle that a geodesic makes with a parallel and r is the radius of the parallel at the intersection point, then Clairaut's relation says that $r \cos \theta$ is constant along geodesics.

(iii) Show that meridians are geodesics; when is a parallel a geodesic?

5. Show that any geodesic of the paraboloid of revolution $z = x^2 + y^2$ which is not a meridian intersects itself an infinite number of times (Hint: use Clairaut's relation. You may assume that no geodesic of a surface of revolution can be asymptotic to a parallel which is not itself a geodesic.)

6. Let S be a connected surface. Assume that the parallel transport between any two points does not depend on the curve joining the points. Prove that the Gaussian curvature of S is zero. (You may find this problem a little hard, but try to spot the main geometrical idea and worry about technicalities later and you may interpret "later" as mexicans do.)

7. The intrinsic distance of a surface S is defined as follows. Given p and q in S let $d(p, q) = \inf_{\alpha \in \Omega(p, q)} \ell(\alpha)$. It can be shown that d is a distance (can you see which property requires some care?) which is compatible with the topology of S . If S is complete the Hopf-Rinow theorem asserts that given two points p and q there exists a geodesic γ joining the points such that $d(p, q) = \ell(\gamma)$.

(i) Show that if $f : S_1 \rightarrow S_2$ is an isometry, then $d_2(f(p), f(q)) = d_1(p, q)$ for all p and q in S_1 .

(ii) A geodesic $\gamma : [0, \infty) \rightarrow S$ is called a *ray leaving from p* if it realizes the distance between $\gamma(0)$ and $\gamma(s)$ for all $s \in [0, \infty)$. Let p be a point in a complete, noncompact surface S . Prove that S contains a ray leaving from p .

8. Show that there are no compact minimal surfaces in \mathbb{R}^3 .

9. The existence of isothermal coordinates is a hard theorem. However for the case of minimal surfaces without planar points it is possible to give an easy proof along the following lines.

(i) Let S be a regular surface without umbilical points. Prove that S is a minimal surface if and only if the Gauss map $N : S \rightarrow S^2$ satisfies

$$\langle dN_p(v_1), dN_p(v_2) \rangle = \lambda(p) \langle v_1, v_2 \rangle$$

for all $p \in S$ and all $v_1, v_2 \in T_p S$, where $\lambda(p) \neq 0$ is a number which depends only on p .

(ii) By considering stereographic projection and (i) show that isothermal coordinates exist around a non planar point in a minimal surface.

For the remaining questions we consider the Weierstrass representation of a minimal surface determined by functions f and g on a simply connected domain $D \subseteq \mathbb{C}$ as we saw in lectures.

10. Show that if ϕ is the parametrization defined by the Weierstrass representation, then ϕ is an immersion if and only if f vanishes only at the poles of g and the order of its zero at such a point is *exactly* twice the order of the pole of g .

11. Find D , f and g representing the catenoid and the helicoid.

12. Show that the Gaussian curvature of the minimal surface determined by the Weierstrass representation is given by

$$K = - \left(\frac{4 |g'|}{|f| (1 + |g|^2)^2} \right)^2.$$

Show that either $K \equiv 0$ or its zeros are isolated.

13. The Weierstrass representation is not unique: if $\phi_{(f,g)} : D \rightarrow \mathbb{R}^3$ is the associated parametrization and $\alpha : W \rightarrow D$ is a bijective holomorphic map, then $\phi_{(f,g)} \circ \alpha$ is another representation of the same minimal surface and it must have the same form with different f and g . Show that by choosing $\alpha(z) = g^{-1}(z)$ we can assume that our pair (f, g) is of the form (F, id) (g can be inverted near a point which is not a pole and for which g' is not zero). We denote such a representation by ϕ_F .

14. Show that the minimal surfaces given by $\phi_{e^{-i\theta} F}$ for θ real are all locally isometric. With an appropriate choice of F , show that the catenoid and the helicoid are locally isometric.