

Yablo's Paradox and the Omitting Types Theorem for Propositional Languages

Thomas Forster

April 2, 2012

This on-line version contains a proof of the extended omitting types theorem which is omitted (Ha!) from the published version

We start by recapitulating Yablo's paradox from [1].

We have infinitely many assertions $\{p_i : i \in \mathbb{N}\}$ and each p_i is equivalent to the assertion that all subsequent p_j are false. A contradiction follows.

There is a wealth of literature on this delightful puzzle, and I have been guilty of a minor contribution to it myself. This literature places Yablo's paradox in the *semantical* column of Ramsey's division of the paradoxes into *semantical* versus *logical* paradoxes. However—as I hope to show below—there is merit to be gained by regarding it as a purely logical puzzle.

Yablo's Paradox in Propositional Logic

If we are to treat Yablo's paradox as a purely logical puzzle we should try to capture it entirely within a first-order language with no special predicates. In fact we can even make progress while using nothing more than a *propositional* language; the obvious language \mathcal{L} to use has infinitely many propositional letters $\{p_i : i \in \mathbb{N}\}$. Next we want a propositional theory with axioms

$$p_i \leftrightarrow \bigwedge_{j>i} \neg p_j \tag{1}$$

for each $i \in \mathbb{N}$,

...but of course we cannot do this in a finitary language. However, one thing we can do in a finitary language is capture the left-to-right direction of these biconditionals, and we do that with the simple scheme

$$p_i \rightarrow \neg p_j \tag{2}$$

for all $i < j \in \mathbb{N}$.

It can be seen that this is equivalent to the even simpler scheme

$$\neg p_i \vee \neg p_j \tag{3}$$

for all $i \neq j \in \mathbb{N}$.

Let us call this theory Y . Y says that at most one p_i can be true.

It is the right-to-left direction of the biconditionals that gives us trouble . . .

$$\left(\bigwedge_{j>i} \neg p_j\right) \rightarrow p_i \tag{4}$$

for each $i \in \mathbb{N}$.

For each i the right-to-left direction of the i th biconditional (4) asserts that at least one of the formulæ in the set $\Sigma(i)$ is false:

$$\{\neg p_j : j \geq i\} \tag{\Sigma(i)}$$

$\Sigma(i)$ is an example of what model theorists call a 0-type, a *type* being nothing more than a set of formulæ¹. The ‘0’ means that the formulæ in the type have no free variables. Our desire that at least one thing in a type should be false is—in the terminology of model theory—a desire to *omit* that type. What we need is a theorem that tells us that a theory can have models that omit a specified type. There is such a theorem, and it is known as the *Omitting Types Theorem*. We say a theory T in a language \mathcal{L} *locally omits* a type Σ if, whenever $\phi \in \mathcal{L}$ is a formula such that T proves $\phi \rightarrow \sigma$ for every $\sigma \in \Sigma$, then $T \vdash \neg\phi$. The omitting types theorem for propositional languages now says:

Theorem 1:

Let T be a consistent theory in a propositional language \mathcal{L} . If T locally omits a type Σ then there is an \mathcal{L} -valuation v that satisfies every theorem of T but falsifies at least one σ in Σ .

We say in these circumstances that v **omits** Σ .

However, what we need here is the slightly stronger:

Theorem 2: (Extended Omitting Types Theorem)

Let T be a consistent theory in a propositional language \mathcal{L} . If T locally omits each type Σ in a countable class \mathfrak{S} of types then there is an \mathcal{L} -valuation that satisfies every theorem of T but, for each $\Sigma \in \mathfrak{S}$, falsifies at least one σ in Σ .

I will omit a proof of this result, since it is standard in the model-theoretic literature.

¹A countably infinite set unless otherwise specified.

In asserting the right-to-left directions (4) of the biconditionals we are restricting ourselves to \mathcal{L} -valuations that omit all the types $\Sigma(i)$. There are countably many of these types so it would be natural to reach for the extended omitting types theorem, theorem 2. Now if we are to exploit theorem 2 we want our theory Y to locally omit each $\Sigma(i)$. But it doesn't. The formula p_0 , in conjunction with the axioms of Y , implies $\neg p_i$ for every $i > 0$ and thereby implies everything in $\Sigma(1)$. If Y were to locally omit $\Sigma(1)$ as we desire then we would have to have $Y \vdash \neg p_0$. But Y clearly does not prove $\neg p_0$. If we were to add $\neg p_0$ as part of a project of adding axioms to Y to obtain a theory that did omit $\Sigma(1)$ we would find by the same token that we would have to add $\neg p_i$ for all other $i \in \mathbb{N}$ as well, and then we end up realising all the $\Sigma(i)$.

Thus Y does not locally omit even one of the $\Sigma(i)$, let alone all of them. So we cannot invoke theorem 2. However, for each i the valuation that makes p_i true and everything else false satisfies Y all right, and it omits all $\Sigma(j)$ for all $j < i$. This illustrates how a theory T can sometimes have a model that omits a type Σ even though T does not locally omit Σ .

Very well: for each i there is an \mathcal{L} -valuation that satisfies Y and omits $\Sigma(j)$ for all $j < i$. Can we find a \mathcal{L} -valuation that satisfies Y and omits all the $\Sigma(i)$? No! Such a valuation would satisfy all the right-to-left directions of the biconditionals in (1), namely the conditionals in (4) and thereby manifest Yablo's paradox!

Conclusion

Yablo's paradox provides us with an illustration of a setting where there is a theory Y and an infinite family $\{\Sigma(i) : i \in \mathbb{N}\}$ of types where, although Y does not locally omit any of the $\Sigma(i)$, it nevertheless has valuations that omit any finite set of them. Further, it has no valuation that omits them all. That last fact illustrates how the condition in theorem 2—namely that T locally omit every $\Sigma \in \mathfrak{S}$ —really is necessary, so the extended omitting types theorem for propositional logic really is best possible.

For T to have a model omitting all the Σ_i it is not sufficient for it to have models omitting any given finite family of them; we really do need the stronger condition that T should locally omit every finite subset of Σ_i .

It illustrates that for T to have a model that omits a type Σ is sufficient but not necessary for T to locally omit Σ .

This is pædagogically quite instructive!

Appendix: A Proof of the Extended Omitting Types theorem for Propositional Logic

I supply a proof of this fact in this on-line version of the paper because—despite the cheerful observation above that the result is standard in the literature—I cannot actually find a proof anywhere!

A *0-type* (Hereafter merely ‘type’: the ‘0’ means that the formulæ in the type have no free variables) in a propositional language \mathcal{L} is a set of formulæ (a countably infinite set unless otherwise specified).

For T an \mathcal{L} -theory a *T-valuation* is an \mathcal{L} -valuation that satisfies T . A valuation v *realises* a type Σ if $v(\sigma) = \mathbf{true}$ for every $\sigma \in \Sigma$. Otherwise v *omits* Σ . We say a theory T *locally omits* a type Σ if, whenever ϕ is a formula such that T proves $\phi \rightarrow \sigma$ for every $\sigma \in \Sigma$, then $T \vdash \neg\phi$.

THEOREM 1 *The Omitting Types Theorem for Propositional Logic*

Let T be a propositional theory, and $\Sigma \subseteq \mathcal{L}(T)$ a type. If T locally omits Σ then there is a T -valuation omitting Σ

Proof:

By contraposition. Suppose there is no T -valuation omitting Σ . Then every formula in Σ is a theorem of T so there is an expression ϕ (namely ‘ \top ’) such that $T \vdash \phi \rightarrow \sigma$ for every $\sigma \in \Sigma$ but $T \not\vdash \neg\phi$. Contraposing, we infer that if $T \vdash \neg\phi$ for every ϕ such that $T \vdash \phi \rightarrow \sigma$ for every $\sigma \in \Sigma$ then there is a T -valuation omitting Σ . ■

However, we can prove something stronger.

THEOREM 2 *The Extended Omitting Types Theorem for Propositional Logic*

Let T be a propositional theory and, for each $i \in \mathbb{N}$, let $\Sigma_i \subseteq \mathcal{L}(T)$ be a type. If T locally omits every Σ_i then there is a T -valuation omitting all of the Σ_i .

Proof:

We will show that whenever $T \cup \{\neg\phi_1, \dots, \neg\phi_i\}$ is consistent, where $\phi_n \in \Sigma_n$ for each $n \leq i$, then we can find $\phi_{i+1} \in \Sigma_{i+1}$ such that $T \cup \{\neg\phi_1, \dots, \neg\phi_i, \neg\phi_{i+1}\}$ is consistent.

Suppose not, then $T \vdash (\bigwedge_{1 \leq j \leq i} \neg\phi_j) \rightarrow \phi_{i+1}$ for every $\phi_{i+1} \in \Sigma_{i+1}$. But, by assumption, T locally omits Σ_{i+1} , so we would have $T \vdash \neg \bigwedge_{1 \leq j \leq i} \neg\phi_j$ contradicting the assumption that $T \cup \{\neg\phi_1, \dots, \neg\phi_i\}$ is consistent.

Now, as long as there is an enumeration of the formulæ in $\mathcal{L}(T)$, we can run an iterative process where at each stage we pick for ϕ_{i+1} the first formula in Σ_{i+1} such that $T \cup \{\neg\phi_1, \dots, \neg\phi_i, \neg\phi_{i+1}\}$ is consistent. This gives us a theory $T \cup \{\neg\phi_i : i \in \mathbb{N}\}$ which is consistent by compactness. Any model of $T \cup \{\neg\phi_i : i \in \mathbb{N}\}$ is a model of T that omits each Σ_i . ■

References

- [1] Steve Yablo, Paradox without self-reference. **Analysis** **53.4** (1993) pp 251–52.

Department of Pure Mathematics and Mathematical Statistics
Centre for Mathematical Sciences
Wilberforce Road
Cambridge CB3 0WB
United Kingdom