

Example sheet 4

1. (a) Let  $p$  be a prime and let  $e$  be a positive integer. Show that  $|\mathrm{SL}_2(\mathbb{Z}/p^e\mathbb{Z})| = p^{3e}(1 - 1/p^2)$ .  
 (b) Show that  $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma(N)] = |\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})| = N^3 \prod_{p|N} (1 - 1/p^2)$ .
2. For an integer  $h$  put  $q_h = e^{2\pi i\tau/h}$ . If  $f \in S_k(\Gamma)$ ,  $s \in \mathbf{P}^1(\mathbb{Q})$  a cusp, then  $s = \sigma(\infty)$  for some  $\sigma \in \mathrm{SL}_2(\mathbb{Z})$  and  $f|[\sigma]_k$  has an expansion  $F(q_h) = \sum_{n>0} a_n q_h^n$  for  $h > 0$  the smallest integer such that  $\begin{pmatrix} 1 & h \\ 0 & 1 \end{pmatrix} \in (\sigma^{-1}\Gamma\sigma)_\infty$  (call this the *Fourier expansion of  $f$  at  $s$  with respect to  $\sigma$* ). Prove that there exists  $\kappa$  such that  $|a_n| \leq \kappa n^{k/2}$ .
3. Let  $\Gamma \subset \mathrm{SL}_2(\mathbb{Z})$  be a congruence subgroup containing  $-1$ . Let  $\Gamma_\infty \subset \Gamma$  be the stabiliser of the cusp  $\infty$ . Show that if  $k > 2$  is even, then the series

$$E_{\Gamma,k}(\tau) = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \frac{1}{(c\tau + d)^k} \quad \text{where } \gamma = \begin{pmatrix} * & * \\ c & d \end{pmatrix}$$

converges and defines an element of  $M_k(\Gamma)$  which is not a cusp form. Identify  $E_{k,\Gamma}(\tau)$  in the case  $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ .

4. Prove that for  $k \geq 1$  the theta series  $\theta(\tau, k)$  converge absolutely and uniformly on compact subsets.
5. Show that  $\Gamma_0(4)$  is generated by  $\pm \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\pm \begin{pmatrix} 1 & 0 \\ 4 & 1 \end{pmatrix}$ . (Hint: Multiply an element of  $\Gamma_0(4)$  on the right alternately by  $\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ 4n & 1 \end{pmatrix}$  for appropriate values of  $n$  to reduce the absolute value of the entries in the bottom row until one of them becomes 0.)
6. Let  $f \in A_{2k}(\Gamma)$  be a non-zero automorphic form for a congruence subgroup  $\Gamma$  of weight  $2k \geq 0$ . Prove that

$$\sum_{\tau_x} (\mathrm{ord}_{\tau_x}(f)/e_{\tau_x} - k(1 - 1/e_{\tau_x})) = k(2g - 2) + k\nu_\infty,$$

where  $\nu_\infty$  is the number of cusps of  $X(\Gamma)$  and the sum is over  $\tau_x \in \mathbb{H}$  giving a set of representatives for the points in  $\Gamma \backslash \mathbb{H}$ .

[Comments and corrections on this Example sheet to T.Berger@dpmms.cam.ac.uk]