

EXAMPLE SHEET 3

1. Suppose $A \subset X$, and let $\partial : H_n(X, A) \rightarrow H_{n-1}(A)$ and $\partial^* : H^{n-1}(A) \rightarrow H^n(X, A)$ be the boundary maps in the long exact sequence of a pair. If $\alpha \in H^{n-1}(A)$, $a \in H_n(X, A)$, show that $\langle \partial^* \alpha, a \rangle = \langle \alpha, \partial a \rangle$.
2. Let C_* be a free finitely generated chain complex defined over \mathbb{Z} . Show that $H_*(C) = 0$ if and only if $H_*(C \otimes \mathbb{Z}/p) = 0$ for every prime p . Now suppose $f : C_* \rightarrow C'_*$ is a chain map. Using problem 5 on Example Sheet 1, show that f is a chain homotopy equivalence if and only if the map $f_* : H_*(C \otimes \mathbb{Z}/p) \rightarrow H_*(C' \otimes \mathbb{Z}/p)$ is an isomorphism for every prime p .
3. Let $R = \mathbb{C}[x]/(x^3)$, and for $i = 1, 2$, let M_i be the R -module $\mathbb{C}[x]/(x^i)$. Find a free resolution of M_1 and use it to compute $\text{Tor}_*^R(M_1, M_1)$ and $\text{Tor}_*^R(M_1, M_2)$.
4. Compute $H_*(L^3(12, 1) \times \mathbb{R}P^3)$ with coefficients in \mathbb{Z} , $\mathbb{Z}/2$, and $\mathbb{Z}/4$.
5. If X is a space, let $\Delta_X : X \rightarrow X \times X$ be the *diagonal map* given by $\Delta(x) = (x, x)$. Compute $\Delta_{S^2_*}([S^2]) \in H_2(S^2 \times S^2)$ and $\Delta_{T^2_*}([T^2]) \in H_2(T^2 \times T^2)$.
6. * Let G be a topological group (*i.e.* G is a group and a space, and the actions of multiplication and taking inverse are continuous.) Show there is a map $\Delta : H^*(G) \rightarrow H^*(G) \otimes H^*(G)$ satisfying the following properties:
 - (a) $\Delta(a \cup b) = \Delta(a) \cup \Delta(b)$, where $(a_1 \otimes a_2) \cup (b_1 \otimes b_2) = (-1)^{|a_2||b_1|} (a_1 \cup b_1) \otimes (a_2 \cup b_2)$.
 - (b) $\Delta(a) = a \otimes 1 + 1 \otimes a + \sum_i a'_i \otimes a''_i$, where $|a'_i| < |a|$ for all i .
 Deduce that $S^1 \times S^{2n}$ cannot be given the structure of a topological group.
7. Let $U, V \subset X$ be open sets. If $x \in H_*(X, U)$ and $y \in H_*(X, V)$, show that $x \cup y \in H_*(X, U \cup V)$. Deduce that $\mathbb{C}P^n$ cannot be covered by n contractible open sets.
8. Let Σ_g be the surface of genus g . Show that if $g < h$, any map $\Sigma_g \rightarrow \Sigma_h$ has degree 0.
9. Let $X_1 = \mathbb{C}P^2 \# \mathbb{C}P^2$, $X_2 = \mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$, and $X_3 = S^2 \times S^2$. Show that $H_*(X_1) \simeq H_*(X_2) \simeq H_*(X_3)$, but that no two of the X_i are homotopy equivalent.
10. Given that $f : S^2 \times S^2 \times S^2 \rightarrow \mathbb{C}P^3$, what are the possible values of the degree of f ?
11. Let M be the Mobius bundle over S^1 . Show that $M \oplus M$ is the trivial bundle.
12. Let $E = TS^2$ be the tangent bundle of S^2 . Show that the unit sphere bundle $S(E)$ is homeomorphic to $SO(3)$, which is also homeomorphic to $\mathbb{R}P^3$.

13. Identify $S^3 - 0$ with \mathbb{R}^3 by stereographic projection. Describe what the fibres of the Hopf fibration look like under this identification. Sketch three distinct fibres.
14. Let $E \rightarrow B$ be a real vector bundle equipped with a Riemannian metric, and let $F \subset E$ be a subbundle. Show that F^\perp is a vector bundle, and that $F \oplus F^\perp \cong E$.
15. Let M be a smooth manifold, and let $\Delta \subset M \times M$ be the image of the diagonal embedding $M \rightarrow M \times M$ which sends x to (x, x) . Show that $\nu_\Delta \simeq TM$.

J.Rasmussen@dpmms.cam.ac.uk