

## EXAMPLE SHEET 1

1. Let  $a : S^n \rightarrow S^n$  be the antipodal map,  $a(x) = -x$ . Show that  $a$  is homotopic to the identity map when  $n$  is odd. [Try  $n = 1$  first.]
2. Let  $f : S^1 \rightarrow S^1$  be a map which is not homotopic to the identity map. Show that there exists an  $x \in S^1$  such that  $f(x) = x$ , and a  $y \in S^1$  so that  $f(y) = -y$ .
3. Suppose that  $f : X \rightarrow Y$  is a map for which there exist maps  $g, h : Y \rightarrow X$  such that  $g \circ f \sim \text{id}_X$  and  $f \circ h \sim \text{id}_Y$ . Show that  $f$ ,  $g$ , and  $h$  are homotopy equivalences.
4. Show that a retract of a contractible space is contractible.
5. Construct a space which contains both the annulus  $S^1 \times I$  and the Möbius band as strong deformation retracts.
6. For  $m < n$ , consider  $S^m$  as a subspace of  $S^n$  given by

$$\{(x_1, x_2, \dots, x_{m+1}, 0, \dots, 0) \mid \sum x_i^2 = 1\}.$$


Show that the complement  $S^n - S^m$  is homotopy equivalent to  $S^{n-m-1}$ .

7. For a map  $f : S^1 \rightarrow X$  we define the *space obtained by attaching a 2-cell to  $X$  along  $f$*  to be the quotient space

$$X \cup_f D^2 := (X \amalg D^2) / \sim$$

where  $\sim$  is the smallest equivalence relation containing  $b \sim f(b)$  for every  $b \in S^1 \subset D^2$ . Show that if  $f, f' : S^1 \rightarrow X$  are homotopic maps then  $X \cup_f D^2 \sim X \cup_{f'} D^2$ .

8. The *dunce cap* is the space obtained from a solid triangle by gluing the edges together as shown.



Show that this space is contractible. [Hint: use the previous question.]

9. Show that the Möbius band does not retract onto its boundary.

10. For based spaces  $(X, x_0)$  and  $(Y, y_0)$  show there is an isomorphism

$$\pi_1(X \times Y, (x_0, y_0)) \cong \pi_1(X, x_0) \times \pi_1(Y, y_0).$$

11. Given a homomorphism  $\varphi : \pi_1(T^2, x_0) \rightarrow \pi_1(T^2, x_0)$ , construct a continuous map  $f : (T^2, x_0) \rightarrow (T^2, x_0)$  with  $f_* = \varphi$ . [Use  $T^2 \cong \mathbb{R}^2/\mathbb{Z}^2$ .] Which  $\varphi$  can be realized as  $f_*$  where  $f$  is a homeomorphism?

12. Show that every homeomorphism  $f : S^1 \rightarrow S^1$  extends to a homeomorphism  $F : D^2 \rightarrow D^2$ . Which of the homeomorphisms  $f : T^2 \rightarrow T^2$  that you constructed in question 11 extend to homeomorphisms  $F : S^1 \times D^2 \rightarrow S^1 \times D^2$ ?

13. Construct a covering map  $\pi : \mathbb{R}^2 \rightarrow K$  of the Klein bottle, and hence show that  $\pi_1(K, k_0)$  is isomorphic to the group  $G$  with elements  $(m, n) \in \mathbb{Z}^2$  and group operation

$$(m, n) * (p, q) = (m + (-1)^n \cdot p, n + q).$$

Show that  $K$  has a covering space homeomorphic to the torus  $T^2$ , but that the torus does not have a covering space homeomorphic to  $K$ .

14.\* A *topological group* consists of a set  $G$  equipped with both a topology and a group structure, so that the inversion map  $i : G \rightarrow G$  (that sends  $g \mapsto g^{-1}$ ) and the multiplication map  $m : G \times G \rightarrow G$  (that sends  $(g, h) \mapsto gh$ ) are continuous. (Here,  $G \times G$  is equipped with the product topology.)

Let  $G$  be a path-connected, locally-path-connected topological group, and  $p : \widehat{G} \rightarrow G$  be a path-connected covering space. Let  $e$  be the identity of  $G$  and  $\epsilon \in p^{-1}(e)$ .

- (i) Show that  $\widehat{G}$  has a unique structure of a topological group with unit  $\epsilon$  so that  $p$  is a continuous homomorphism.
- (ii) Show that  $\text{Ker}(p) \subset \widehat{G}$  lies in the centre of  $\widehat{G}$ .
- (iii) Show that  $SO(3)$ , the group of rotations of  $\mathbb{R}^3$  (or equivalently of orthogonal  $3 \times 3$  matrices of determinant 1), is homeomorphic to the projective space  $\mathbb{RP}^3$ .
- (iv) Together, (i) and (iii) give a covering space  $\widehat{SO(3)}$  homeomorphic to  $S^3$ . Identify this group with a well-known matrix group.