

**Part III Algebraic Topology // Example Sheet 2**

**1.** Construct a natural map  $H^n(X) \rightarrow \text{Hom}(H_n(X), \mathbb{Z})$ , and similarly for relative (co)homology, and prove that these maps commute with the  $\partial$ -maps in the long exact sequence for a pair. Show that your map is a surjection, but that it is not always an isomorphism.

**2.\*** If  $f : X \rightarrow X$  is a homeomorphism, let  $T_f$  be the quotient space of  $X \times [0, 1]$  by  $(x, 0) \sim (f(x), 1)$ . By choosing an appropriate open cover, construct a long exact sequence

$$\cdots \rightarrow H_{n+1}(T_f) \rightarrow H_n(X) \xrightarrow{1-f_*} H_n(X) \rightarrow H_n(T_f) \rightarrow \cdots$$

Calculate  $H_*(T_f)$  when (a)  $f : S^n \rightarrow S^n$  is the antipodal map, (b)  $f : \mathbb{R}^2/\mathbb{Z}^2 \rightarrow \mathbb{R}^2/\mathbb{Z}^2$  is induced by the matrix  $\begin{pmatrix} 3 & 4 \\ 2 & 3 \end{pmatrix}$ .

**3.** Say a map  $f : X \rightarrow Y$  between cell complexes is *cellular* if  $f(X^n) \subset Y^n$  for every  $n$ . Show how to associate to such an  $f$  a chain map  $f_{\#}^{cell} : C_{\bullet}^{cell}(X) \rightarrow C_{\bullet}^{cell}(Y)$  and show that the induced map  $f_*^{cell} : H_*^{cell}(X) \rightarrow H_*^{cell}(Y)$  agrees with  $f_* : H_*(X) \rightarrow H_*(Y)$  under a suitable identification of the homology groups.

**4.\*** If  $f : S^n \rightarrow X$  is a map, let  $X \cup_f D^{n+1}$  be the space obtained by gluing  $D^{n+1}$  to  $X$  along the map  $f$ .

(i) If  $f \simeq f' : S^n \rightarrow X$ , show that  $X \cup_f D^{n+1} \simeq X \cup_{f'} D^{n+1}$ .

(ii) Let  $Y = S^n \cup_f D^{n+1}$  be constructed using a map  $f : S^n \rightarrow S^n$  of degree  $m > 1$ . Show that the natural quotient map  $Y \rightarrow Y/S^n \cong S^{n+1}$  is trivial on homology  $H_{* > 0}$ , but is non-trivial on cohomology  $H^{* > 0}$ . What happens if we instead consider the inclusion  $S^n \hookrightarrow Y$ ?

**5.**

(i) Let  $X$  be a cell complex and  $A \subset X$  be a subcomplex. Prove that the pair  $(X, A)$  is good.

(ii) Let  $X$  be a cell complex and  $K \subset X$  a compact subspace. Prove that  $K$  intersects only finitely many open cells in  $X$ . Hence show that any element of  $H_i(X)$  lies in the image of  $H_i(X^m) \rightarrow H_i(X)$  for some  $m \gg 0$ .

**6.** If  $X$  and  $Y$  are finite cell complexes with cells  $\{e_{\alpha}\}_{\alpha \in I}$  and  $\{f_{\beta}\}_{\beta \in J}$ , construct a cell structure on  $X \times Y$  with cells  $\{e_{\alpha} \times f_{\beta}\}_{(\alpha, \beta) \in I \times J}$ . Hence show that there is an isomorphism of chain complexes  $C_{\bullet}^{cell}(X \times Y) \cong C_{\bullet}^{cell}(X) \otimes C_{\bullet}^{cell}(Y)$ , where the latter has differential  $d(e_{\alpha} \otimes f_{\beta}) = d(e_{\alpha}) \otimes f_{\beta} + (-1)^{\dim(e_{\alpha})} e_{\alpha} \otimes d(f_{\beta})$ . [Hint: to understand this sign, it may help to think about the cellular chain complex of  $D^p \times D^q$ .]

Use this to calculate  $H_*(\mathbb{R}P^2 \times \mathbb{R}P^2)$ .

**7.** Show that for  $m, n \in \mathbb{N}$  and any space  $X$  there are short exact sequences of chain complexes

$$0 \rightarrow C^{\bullet}(X) \rightarrow C^{\bullet}(X) \rightarrow C^{\bullet}(X; \mathbb{Z}/m) \rightarrow 0$$

$$0 \rightarrow C^{\bullet}(X; \mathbb{Z}/n) \rightarrow C^{\bullet}(X; \mathbb{Z}/n \cdot m) \rightarrow C^{\bullet}(X; \mathbb{Z}/m) \rightarrow 0$$

and hence describe ‘‘Bockstein operations’’

$$\tilde{\beta} : H^i(X; \mathbb{Z}/m) \rightarrow H^{i+1}(X) \quad \text{and} \quad \beta : H^i(X; \mathbb{Z}/m) \rightarrow H^{i+1}(X; \mathbb{Z}/n).$$

How are these two operations related? Compute the effect of  $\beta$  and  $\tilde{\beta}$  for  $m = 2$ ,  $n = 2^r$ , and  $X = \mathbb{RP}^k$ .

Show that  $\beta(x \smile y) = \beta(x) \smile y + (-1)^{|x|} x \smile \beta(y)$ .

**8.** A map  $\pi : E \rightarrow B$  is called a *covering map* if there is an open cover  $\{U_\alpha\}$  of  $B$  such that  $\pi^{-1}(U_\alpha)$  is a disjoint union  $\coprod V_{\alpha,\beta}$  with each  $\pi|_{V_{\alpha,\beta}} : V_{\alpha,\beta} \rightarrow U_\alpha$  a homeomorphism.

(i) If  $\pi : E \rightarrow B$  is a covering map with finite fibres of cardinality  $N$ , show how to construct a map  $\pi^! : H_*(B) \rightarrow H_*(E)$  such that  $\pi_* \circ \pi^!$  is multiplication by  $N$ .

(ii) In the same situation, if  $B$  is a finite cell complex show that  $\chi(E) = N \cdot \chi(B)$ .

(iii) Show there is a covering map  $\Sigma_g \rightarrow \Sigma_h$  if and only if  $g = kh - k + 1$  for some  $k \in \mathbb{N}$ .

**9.** Show that there is a *relative cup product*

$$\smile : H^i(X, A) \times H^j(X, B) \longrightarrow H^{i+j}(X, A \cup B)$$

[Hint: it may be helpful to consider a cochain complex  $C_{A+B}^*(X)$  of cochains vanishing on simplices lying wholly in  $A$  or  $B$ , and use the Small Simplices Theorem.] Using this, show that if  $X$  has a cover by  $n$  contractible (i.e. homotopy equivalent to a point) open sets, then the *cup-length*

$$\max \{k \mid \exists a_1, \dots, a_k \in H^{*>0}(X), a_1 \smile \dots \smile a_k \neq 0\}$$

is strictly smaller than  $n$ . What does this say about the ring  $H^*(\Sigma X)$ , where  $\Sigma$  is the suspension operation?

**10.**

(i) Let  $e : [0, 1]^k \rightarrow S^n$  be a map which is a homeomorphism onto its image  $D \subset S^n$ . By considering the open sets

$$A = S^n \setminus e([0, 1]^{k-1} \times [0, 1/2]) \quad B = S^n \setminus e([0, 1]^{k-1} \times [1/2, 1])$$

in  $S^n$ , show by induction on  $k$  that  $\tilde{H}_i(S^n \setminus D) = 0$ .

(ii) If  $e : S^k \rightarrow S^n$  is a map which is a homeomorphism onto its image  $S \subset S^n$ , compute  $\tilde{H}_i(S^n \setminus S)$ . Think about the consequence of this in the case  $(n, k) = (2, 1)$ .

Comments or corrections to [or257@cam.ac.uk](mailto:or257@cam.ac.uk)