Representation Theory — Examples Sheet 3

On this sheet all groups are finite and all representations are complex representations

- 1. Calculate S^2V and Λ^2V for the two-dimensional irreducible representations of D_8 and of Q_8 . Which has the trivial representation as a subrepresentation in each case?
- 2. Let $G = S_n$ act naturally on the set $X = \{1, ..., n\}$. For each non-negative integer r, let X_r be the set of all r-element subsets of X equipped with the natural action of G, and π_r be the character of the corresponding permutation represention. If $0 \le l \le k \le n/2$, show that

$$\langle \pi_k, \pi_l \rangle_G = l + 1.$$

Deduce that $\pi_r - \pi_{r-1}$ is a character of an irreducible representation for each $1 \le r \le n/2$. What happens for r > n/2?

3. Suppose $\rho: G \to GL(V)$ is an irreducible representation of G with character χ . By considering $V \otimes V$, S^2V and Λ^2V show that

$$\frac{1}{|G|} \sum_{g \in G} \chi(g^2) = \begin{cases} 0 & \text{if } \chi \text{ is not real-valued} \\ \pm 1 & \text{if } \chi \text{ is real valued}. \end{cases}$$

Deduce that if |G| is odd then G has only one real-valued irreducible character.

- 4. Let $\rho: G \to GL(V)$ be a representation of G of dimension d.
 - (a) Compute dim S^nV and dim Λ^nV for all n.
 - (b) Let $g \in G$ and $\lambda_1, \ldots, \lambda_d$ be the eigenvalues of $\rho(g)$. What are the eigenvalues of g on S^nV and Λ^nV ?
 - (c) Let $f(t) = \det(tI \rho(g))$ be the characteristic polynomial of $\rho(g)$. What is the relationship between the coefficients of f and $\chi_{\Lambda^n V}$?
 - (d) What is the relationship between $\chi_{S^nV}(g)$ and f? (Hint: start with case d=1).
- 5. Recall the character table of D_{10} from sheet 2. Explain how to view D_{10} as a subgroup of A_5 and then use induction from D_{10} to A_5 to reconstruct the character table of A_5 .
- 6. Obtain the character table of the dihedral group D_{2m} by using induction from the cyclic group C_m ; you will want to split into two cases according as m is odd or even.
- 7. Find all the characters of S_5 obtained by inducing irreducible representations of S_4 . Use these to reconstruct the character table of S_5 . Then repeat, replacing S_4 by the subgroup $\langle (12345), (2354) \rangle$ of S_5 of order 20.
- 8. Prove that if H is a subgroup of a group G, and K is a subgroup of H, and W is a representation of K then $\operatorname{Ind}_K^G W \cong \operatorname{Ind}_H^G \operatorname{Ind}_K^H W$.
- 9. Let H be a subgroup of a group G. Show that for every irreducible representation (ρ, V) of G there is an irreducible representation (ρ', W) of H such that ρ is an irreducible component of $\operatorname{Ind}_H^G W$.
 - Deduce that if A is an abelian subgroup of G then every irreducible representation of G has dimension at most |G/A|.
- 10. Suppose that G is a Frobenius group with Frobenius kernel K. Show that if V is a non-trivial irreducible representation of K then $\operatorname{Ind}_K^G V$ is also irreducible. Hence, explain how to construct the character table of G given the character tables of K and G/K.
- 11. Suppose that V is a faithful representation of a group G such that χ_V takes r distinct values. Show that each irreducible representation of G is a summand of $V^{\otimes n}$ for some n < r.
- 12. Suppose G is a finite group of odd order and with k conjugacy classes. Show that $|G| \equiv k \mod 16$.

Comments and corrections to S.J.Wadsley@dpmms.cam.ac.uk.