

(1) Show the tangent space to S^2 at a point $P = (x, y, z) \in S^2$ is the plane normal to the vector \overrightarrow{OP} , where O denotes the origin.

(2) Let V be the open subset $\{0 < u < \pi, 0 < v < 2\pi\}$, and $\sigma : V \rightarrow S^2$ be given by

$$\sigma(u, v) = (\sin u \cos v, \sin u \sin v, \cos u).$$

Prove that σ defines a smooth parametrization on a certain open subset of S^2 . [You may assume that \cos^{-1} is continuous on $(-1, 1)$, and that \tan^{-1}, \cot^{-1} are continuous on $(-\infty, \infty)$.]

(3) Show the stereographic projection map $\pi : S \setminus \{N\} \rightarrow \mathbf{C}$, where N denotes the north pole, defines a chart. Check that the spherical metric on $S \setminus \{N\}$ corresponds under π to the Riemannian metric on \mathbf{C} given by

$$4(dx^2 + dy^2)/(1 + x^2 + y^2)^2.$$

(4) For an embedded circular cylinder S in \mathbf{R}^3 , show that the first fundamental form corresponds to a locally Euclidean Riemannian metric on S . Identify the geodesics on S .

(5) Given a smooth curve $\Gamma : [0, 1] \rightarrow S$ on an abstract surface S with a Riemannian metric, show that the length l is unchanged under reparametrizations of the form $f : [0, 1] \rightarrow [0, 1]$, with $f'(t) > 0$ for all $t \in [0, 1]$. Prove that there exists such a reparametrization $\tilde{\Gamma} = \Gamma \circ f$ for which $\|d\tilde{\Gamma}/dt\|$ is constant, namely $1/l$.

(6) Let T denote the embedded torus in \mathbf{R}^3 obtained by revolving around the z -axis the circle $(x - 2)^2 + z^2 = 1$ in the xz -plane. Using the formal definition of area in terms of a parametrization, calculate the surface area of T .

(8) If one places S^2 inside a (vertical) circular cylinder of radius one, prove that the radial (horizontal) projection map from S^2 to the cylinder preserves areas (this is usually known as *Archimedes Theorem*). Deduce the existence of an atlas on S^2 , for which the charts all preserve areas and the transition functions have derivatives with determinant one.

(9) Show that a 2-holed torus may be obtained topologically by suitably identifying the sides of a regular octagon. Indicate briefly how to extend your argument to show that a g -holed torus may be obtained topologically by suitably identifying the sides of a regular $4g$ -gon?

(10) For a surface of revolution S , corresponding to a curve $\eta : (a, b) \rightarrow \mathbf{R}^3$ given by $\eta(u) = (f(u), 0, g(u))$, where η is parametrized in such a way that $\|\eta'\| = 1$, prove that the second fundamental form at a given point is given by

$$(f'g'' - f''g')du^2 + fg'dv^2.$$

Deduce that the Gaussian curvature K is given by the formula $K = -f''/f$.

(11) Using the results from the previous question, calculate the Gaussian curvature K of the unit sphere and the hyperboloid of one sheet. For the embedded torus, as defined in Question 6, identify those points at which $K = 0$, $K > 0$ and $K < 0$. Verify the global Gauss–Bonnet theorem on the embedded torus.

(12) Let S be an embedded surface in \mathbf{R}^3 which is closed and bounded. By considering the smallest closed ball centred on the origin which contains S , or otherwise, show that the Gaussian curvature must be strictly positive at some point of S .

(13) Show that Mercator’s parametrization of the sphere (minus poles)

$$\sigma(u, v) = (\operatorname{sech} u \cos v, \operatorname{sech} u \sin v, \tanh u)$$

determines a chart (on the complement of a longitude) which preserves angles and sends meridians and parallels on the sphere to straight lines in the plane.

(14) For a smooth curve Γ on an embedded surface S in \mathbf{R}^3 , prove that the geodesic equations are equivalent to the statement that $d^2\Gamma/dt^2$ is always normal to S . Deduce that any such geodesic Γ has $\|d\Gamma/dt\|$ constant.

(15) Let $f(u) = e^u$, $g(u) = (1 - e^{2u})^{\frac{1}{2}} - \cosh^{-1}(e^{-u})$, where $u < 0$, and S be the surface of revolution corresponding to the curve $\eta : (-\infty, 0) \rightarrow \mathbf{R}^3$ given by $\eta(u) = (f(u), 0, g(u))$. Show that S has constant Gauss curvature -1 ; S is called the *pseudosphere*. By considering coordinates v and $w = e^{-u}$ on S , show that the pseudosphere is isometric to the open subset of the upper half-plane model of the hyperbolic plane given by $\operatorname{Im}(z) > 1$. Identify the geodesics in this model corresponding to the meridians in S . Show that any other geodesic in S has two distinct points of the circle $x^2 + y^2 = 1, z = 0$ as limit points.

[It is a theorem that there are no complete embedded surfaces in \mathbf{R}^3 with constant negative Gauss curvature, and so in particular we cannot realise all of the hyperbolic plane as an embedded surface.]

Note to the reader : You should look at all the questions up to Question 12, and then any further questions you have time for.