

Groups Rings and Modules: Example Sheet 3 of 4

All rings in this course are commutative with a 1.

1. Show that $\mathbb{Z}[\sqrt{-2}]$ and $\mathbb{Z}[\omega]$ are Euclidean domains, where $\omega = \frac{1}{2}(-1 + \sqrt{-3})$. Show also that the usual Euclidean function $\phi(r) = N(r)$ does not make $\mathbb{Z}[\sqrt{-3}]$ into a Euclidean domain. Could there be some other Euclidean function ϕ making $\mathbb{Z}[\sqrt{-3}]$ into a Euclidean domain?
2. Show that the ideal $(2, 1 + \sqrt{-7})$ in $\mathbb{Z}[\sqrt{-7}]$ is not principal.
3. Find an element of $\mathbb{Z}[\sqrt{-17}]$ that is a product of two irreducibles and also a product of three irreducibles.
4. Determine whether or not the following rings are fields, PIDs, UFDs, integral domains:
 $\mathbb{Z}[X]$, $\mathbb{Z}[X]/(X^2+1)$, $\mathbb{Z}[X]/(2, X^2+1)$, $\mathbb{Z}[X]/(2, X^2+X+1)$, $\mathbb{Z}[X]/(3, X^3-X+1)$.
5. Determine which of the following polynomials are irreducible in $\mathbb{Q}[X]$:
 $X^4 + 2X + 2$, $X^4 + 18X^2 + 24$, $X^3 - 9$, $X^3 + X^2 + X + 1$, $X^4 + 1$, $X^4 + 4$.
6. Let R be an integral domain. The *greatest common divisor* (gcd) of non-zero elements a and b in R is an element d in R such that d divides both a and b , and if c divides both a and b then c divides d .
 - (i) Show that the gcd of a and b , if it exists, is unique up to multiplication by a unit.
 - (ii) In lectures we have seen that, if R is a UFD, the gcd of two elements exists. Give an example to show that this is not always the case in an integral domain.
 - (iii) Show that if R is a PID, the gcd of elements a and b exists and can be written as $ra + sb$ for some $r, s \in R$. Give an example to show that this is not always the case in a UFD.
 - (iv) Explain briefly how, if R is a Euclidean domain, the Euclidean algorithm can be used to find the gcd of any two non-zero elements. Use the algorithm to find the gcd of $11 + 7i$ and $18 - i$ in $\mathbb{Z}[i]$.
7. Find all ways of writing the following integers as sums of two squares: 221 , 209×221 , 121×221 , 5×221 .
8. By considering factorisations in $\mathbb{Z}[\sqrt{-2}]$, show that the only integer solutions to the equation $x^2 + 2 = y^3$ are $x = \pm 5$, $y = 3$.
9. Let R be any ring. Show that the ring $R[X]$ is a principal ideal domain if and only if R is a field. If I and J are ideals in a ring R then must the set $\{ab : a \in I, b \in J\}$ also be an ideal in R ?
10. Exhibit an integral domain R and a (non-zero, non-unit) element of R that is not a product of irreducibles.

11. Let \mathbb{F}_q be a finite field with q elements.
- Show that the prime subfield K (that is, the smallest subfield) of \mathbb{F}_q has p elements for some prime number p . Show that \mathbb{F}_q is a vector space over K and deduce that $q = p^k$, for some k .
 - Show that the multiplicative group of the non-zero elements of \mathbb{F}_q is cyclic. Deduce that if $q = p^2$ then $\mathrm{SL}_2(\mathbb{F}_p)$ contains an element of order $p + 1$.

Further Questions

12. (i) Consider the polynomial $f = X^3Y + X^2Y^2 + Y^3 - Y^2 - X - Y + 1$ in $\mathbb{C}[X, Y]$. Write it as an element of $(\mathbb{C}[X])[Y]$, that is collect together terms in powers of Y , and then use Eisenstein's criterion to show that f is prime in $\mathbb{C}[X, Y]$.
- (ii) Let F be any field. Show that the polynomial $f = X^2 + Y^2 - 1$ is irreducible in $F[X, Y]$, unless F has characteristic 2. What happens in that case?
13. Show that the subring $\mathbb{Z}[\sqrt{2}]$ of \mathbb{R} is a Euclidean domain. Show that the units are $\pm(1 \pm \sqrt{2})^n$ for $n \geq 0$.
14. If a UFD has at least one irreducible, must it have infinitely many (pairwise non-associate) irreducibles?
15. Let V be a 2-dimensional vector space over a field \mathbb{F}_q of q elements, let Ω be the set of its 1-dimensional subspaces.
- Show that Ω has size $q + 1$ and $\mathrm{GL}_2(\mathbb{F}_q)$ acts on it. Show that the kernel Z of this action consists of scalar matrices and the group $\mathrm{PGL}_2(\mathbb{F}_q) = \mathrm{GL}_2(\mathbb{F}_q)/Z$ has order $q(q^2 - 1)$. Show that the group $\mathrm{PSL}_2(\mathbb{F}_q)$ obtained similarly from $\mathrm{SL}_2(\mathbb{F}_q)$ has order $q(q^2 - 1)/d$ with $d = \gcd(q - 1, 2)$.
 - Show that Ω may be identified with the set $\mathbb{F}_q \cup \{\infty\}$ in such a way that $\mathrm{GL}_2(\mathbb{F}_q)$ acts on Ω as the group of Möbius transformations $z \mapsto \frac{az+b}{cz+d}$.
16. Show that the groups $\mathrm{SL}_2(\mathbb{F}_4)$ and $\mathrm{PSL}_2(\mathbb{F}_5)$ defined above both have order 60. Use this and some questions from Example Sheet 1 to show that they are both isomorphic to the alternating group A_5 . Show that $\mathrm{SL}_2(\mathbb{F}_5)$ and $\mathrm{PGL}_2(\mathbb{F}_5)$ both have order 120, that $\mathrm{SL}_2(\mathbb{F}_5)$ is not isomorphic to S_5 , but $\mathrm{PGL}_2(\mathbb{F}_5)$ is.
- [You may find it helpful to show, using the Cayley-Hamilton theorem or otherwise, that the order of an element $I \neq A \in \mathrm{SL}_2(\mathbb{F}_4)$ is uniquely determined by its trace.]