

1. Let  $X$  be a Markov chain containing an absorbing state  $s$  to which all other states lead (i.e.,  $j \rightarrow s$  for all  $j$ ). Show that all states other than  $s$  are transient.

2. Compute  $p_{11}^{(n)}$  and classify the states of the Markov chain with state space  $I = \{1, 2, 3\}$  and transition matrix

$$\begin{pmatrix} 1-2p & 2p & 0 \\ p & 1-2p & p \\ 0 & 2p & 1-2p \end{pmatrix}.$$

3. A particle performs a random walk on the vertices of a cube. At each step it remains where it is with probability  $\frac{1}{4}$ , and moves to each of its neighbouring vertices with probability  $\frac{1}{4}$ . Let  $v$  and  $w$  be two diametrically opposite vertices. If the walk starts at  $v$ , find (a) the mean number of steps until its first return to  $v$ , (b) the mean number of steps until its first visit to  $w$ , (c) the mean number of visits to  $w$  before its first return to  $v$ .

4. (Harder) Let  $X$  be a Markov chain on  $\{0, 1, 2, \dots\}$  with transition matrix given by  $p_{0,j} = a_j$  for  $j \geq 0$ ,  $p_{i,i} = r$  and  $p_{i,i-1} = 1 - r$  for  $i \geq 1$ . Assume that  $0 < r < 1$ . Classify the states of the chain, and find their mean recurrence times. [You may find it useful to define  $J = \sup\{j : a_j > 0\}$ .]

5. In Exercise 1.10, which states are recurrent and which are transient?

6. What can be said about the number of visits to each state in the case where (a) a Markov chain is transient, and (b) a Markov chain is recurrent?

Consider the Markov chain  $(X_n)_{n \geq 0}$  of Exercise 1.13. Show for this chain that  $\mathbb{P}[X_n \rightarrow \infty \text{ as } n \rightarrow \infty] = \mathbb{P}[\forall m, \exists n \text{ such that } X_N \geq m \text{ for all } N \geq n] = 1$ .

Suppose the transition probabilities satisfy instead

$$p_{i,i+1} = \left(\frac{i+1}{i}\right)^\alpha p_{i,i-1}.$$

For each  $\alpha \in (0, \infty)$  find the value of  $\mathbb{P}[X_n \rightarrow \infty \text{ as } n \rightarrow \infty]$ .

7. The rooted binary tree is an infinite graph  $T$  with one distinguished vertex  $R$  from which comes a single edge; at every other vertex there are three edges and there are no closed loops. The random walk on  $T$  jumps from a vertex along each available edge with equal probability. Show that the random walk is transient.

8. Show (by projection onto  $\mathbb{Z}^3$  or otherwise) that the simple symmetric random walk in  $\mathbb{Z}^4$  is transient.

9. Find all invariant distributions of the transition matrix in Exercise 9 of Example Sheet 1.

10. Two containers A and B are placed adjacently to one another, and gas is allowed to pass through a small aperture joining them. There are  $N$  molecules in all, and we assume that, at each epoch of time, one molecule (chosen at random) passes through the aperture. Show that the number of molecules in A evolves as a Markov chain. What are the transition probabilities? What is the invariant distribution of this chain? [This is the ‘Ehrenfest urn model’, first introduced by Ehrenfest under the name ‘dog–flea model’.]

11. A fair die is thrown repeatedly. Let  $X_n$  denote the sum of the first  $n$  throws. Find

$$\lim_{n \rightarrow \infty} \mathbb{P}[X_n \text{ is a multiple of } 13]$$

quoting carefully any general theorems that you use.

12. Find the invariant distributions of the transition matrices in Exercise 8 of Example Sheet 1, parts (a), (b) and (c), and compare them with your answers to that exercise.

13. Each morning a student takes one of the three books (labelled 1, 2, 3) he owns from his shelf. The probability that he chooses the book with label  $i$  is  $\alpha_i$  (where  $0 < \alpha_i < 1$ ,  $i = 1, 2, 3$ ), and choices on successive days are independent. In the evening he replaces the book at the left-hand end of the shelf. If  $p_n$  denotes the probability that on day  $n$  the student finds the books in the order 1,2,3, from left to right, show that, irrespective of the initial arrangement of the books,  $p_n$  converges as  $n \rightarrow \infty$ , and determine the limit.

14. In each of the following cases determine whether the stochastic matrix  $P$  corresponds to a chain which is reversible in equilibrium:

(a)  $\begin{pmatrix} p & 1-p \\ q & 1-q \end{pmatrix}$ ;

(b)  $\begin{pmatrix} 0 & p & 1-p \\ 1-p & 0 & p \\ p & 1-p & 0 \end{pmatrix}$ ;

(c)  $I = \{0, 1, 2, \dots\}$  and  $p_{01} = 1$ ,  $p_{i,i+1} = p$ ,  $p_{i,i-1} = 1-p$  for  $i \geq 1$ .

(d)  $p_{ij} = p_{ji}$  for all  $i, j \in I$ .

15. A random walk on the set  $\{0, 1, 2, \dots, b\}$  has transition matrix given by  $p_{00} = 1 - \lambda_0$ ,  $p_{bb} = 1 - \mu_b$ ,  $p_{i,i+1} = \lambda_i$  and  $p_{i+1,i} = \mu_{i+1}$  for  $0 \leq i < b$ , where  $0 < \lambda_i, \mu_i < 1$  for all  $i$ , and  $\lambda_i + \mu_i = 1$  for  $1 \leq i < b$ . Show that this process is time-reversible in equilibrium.

16. Let  $X$  be an irreducible non-null recurrent aperiodic Markov chain. Show that  $X$  is time-reversible in equilibrium if and only if

$$p_{j_1 j_2} p_{j_2 j_3} \cdots p_{j_{n-1} j_n} p_{j_n j_1} = p_{j_1 j_n} p_{j_n j_{n-1}} \cdots p_{j_2 j_1}$$

for all  $n$  and all finite sequences  $j_1, j_2, \dots, j_n$  of states.