

Part II

Algebraic Geometry

Example Sheet IV, 2016

(For all questions, assume k is algebraically closed. Further, you can assume the characteristic is not equal to 2 if necessary. A * indicates a more difficult problem.)

1. If P is a smooth point of an irreducible curve X and $t \in \mathcal{O}_{X,P}$ is a local parameter at P , show that $\dim_k \mathcal{O}_{X,P}/(t^n) = n$ for every $n \in \mathbb{N}$.
2. Show that $X = Z(x_0^8 + x_1^8 + x_2^8)$ and $Y = Z(y_0^4 + y_1^4 + y_2^4)$ are irreducible smooth curves in \mathbb{P}^2 provided $\text{char}(k) \neq 2$, and that $\phi: (x_i) \mapsto (x_i^2)$ is a morphism from X to Y . Determine the degree of ϕ , and compute e_P for all $P \in X$.
3. Show that the plane cubic $X = Z(f)$, $f = x_0x_2^2 - x_1^3 + 3x_1x_0^2$, is smooth if $\text{char}(k) \neq 2, 3$. Find the degree and ramification degrees (i.e., the e_P) for (i) the projection $\phi = (x_0 : x_1): X \rightarrow \mathbb{P}^1$ (ii) the projection $\phi = (x_0 : x_2): X \rightarrow \mathbb{P}^1$.
4. (i) Let $\phi = (1 : f): \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a morphism given by a nonconstant polynomial $f \in k[t] \subset K(\mathbb{P}^1)$. Show that $\deg(\phi) = \deg f$, and determine the ramification points of ϕ — that is, the points $P \in \mathbb{P}^1$ for which $e_P > 1$. Do the same for a rational function $f \in k(t)$.
(ii) Assume the characteristic of k is 0. Let $\phi = (t^2 - 7 : t^3 - 10): \mathbb{P}^1 \rightarrow \mathbb{P}^1$. Compute $\deg(\phi)$ and e_P for all $P \in \mathbb{P}^1$.
(iii) Let $f, g \in k[t]$ be coprime polynomials with $\deg(f) > \deg(g)$, and $\text{char}(k) = 0$. Assume that every root of $f'g - g'f$ is a root of fg . Show that g is constant and f is a power of a linear polynomial.
(iv) Let $\phi: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be a finite morphism in characteristic zero. Suppose that every ramification point $P \in \mathbb{P}^1$ satisfies $\phi(P) \in \{0, \infty\}$. Show that $\phi = (F_0^n : F_1^n)$ for some linear forms F_i . [Hint: choose coordinates so that $\phi(0) = 0$ and $\phi(\infty) = \infty$.]
(v) Suppose $\text{char}(k) = p \neq 0$, and let $\phi: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be given by $t^p - t \in k(t)$. Show that ϕ has degree p and that it is only ramified at ∞ .
5. Let X be a non-singular projective curve. Let $V \subset K(X)$ be a finite-dimensional k -vector subspace of $K(X)$. Show that there exists a divisor D on V for which $V \subset \mathcal{L}(D)$.
6. Let X be a smooth plane cubic. Assume that V has equation $x_0x_2^2 = x_1(x_1 - x_0)(x_1 - \lambda x_0)$, for some $\lambda \in k \setminus \{0, 1\}$.
Let $P = (0 : 0 : 1)$ be the point at infinity in this equation. Writing $x = x_1/x_0$, $y = x_2/x_0$, show that x/y is a local parameter at P . [Hint: consider the affine

piece $x_2 \neq 0$.] Hence compute $v_P(x)$ and $v_P(y)$. Show that for each $m \geq 1$, the space $\mathcal{L}(mP)$ has a basis consisting of functions $x^i, x^j y$, for suitable i and j , and that $\ell(mP) = m$.

7. Let $f \in k[x]$ a polynomial of degree $d > 1$ with distinct roots, and $V \subset \mathbb{P}^2$ the projective closure of the affine curve with equation $y^{d-1} = f(x)$. Assume that $\text{char}(k)$ does not divide $d - 1$. Prove that V is smooth, and has a single point P at infinity. Calculate $v_P(x)$ and $v_P(y)$.

* Deduce (without using Riemann–Roch) that if $n > d(d - 3)$, then $\ell((n + 1)P) = \ell(nP) + 1$.

8. A non-singular projective curve X is covered by two affine pieces (with respect to different embeddings) which are affine plane curves with equations $y^2 = f(x)$ and $v^2 = g(u)$ respectively, with f a square-free polynomial of even degree $2n$ and $u = 1/x, v = y/x^n$ in $K(X)$. Determine the polynomial $g(u)$ and show that the canonical class on X has degree $2n - 4$. Why can we not just say that X is the projective plane curve associated to the affine curve $y^2 = f(x)$?

9. Let $X_0 \subset \mathbb{A}^2$ be the affine curve with equation $y^3 = x^4 + 1$, and let $X \subset \mathbb{P}^2$ be its projective closure. Show that X is smooth, and has a unique point Q at infinity. Let ω be the rational differential dx/y^2 on X . Show that $v_P(\omega) = 0$ for all $P \in X_0$. prove that $v_Q(\omega) = 4$ and hence that $\omega, x\omega$ and $y\omega$ are all regular on X .

10. Let X be a non-singular projective curve and $P \in X$ any point. Show that there exists a nonconstant rational function on X which is regular everywhere except at P . Show moreover that there exists a projective embedding of X which has P as its unique point at infinity. If X has genus g , show that there exists a nonconstant morphism $X \rightarrow \mathbb{P}^1$ of degree g .

11. Let P_0 be a point on an elliptic curve (non-singular projective curve of genus 1!) and $\phi_{3P_0}: X \rightarrow \mathbb{P}^2$ the projective embedding. Show that $P \in X$ is a point of inflection if and only if $3P = 0$ in the group law determined by P_0 . Deduce that if P and Q are points of inflection then so is the third point of intersection of the line PQ with X .

12. Let $\pi: X \rightarrow \mathbb{P}^1$ be a hyperelliptic cover, and $P \neq Q$ ramification points of π . Show that as elements of $Cl^0(X)$, $P - Q \neq 0$ but $2(P - Q) = 0$.