

## Example Sheet 3

1. Compute the average busy period for a  $M/M/\infty$  and a  $M/M/1$  queue. (The busy period  $B$  is the length of time between the arrival of the first customer and the first time afterwards that all servers are free).
2. Consider the  $M/M/n$  queue, where the arrival rate is  $\lambda$  and the service rate in each queue is  $\mu$ . For which values of the parameters is the queue length transient, positive recurrent and null recurrent? Compute the invariant distribution when there exists one.
3. *Queue with impatient customers.* Customers arrive at a single server at rate  $\lambda$  and require an exponential amount of service with rate  $\mu$ . Customers waiting in line are impatient and if they are not in service they will leave at rate  $\delta$ , independent of their position in the queue. (a) Show that for any  $\delta > 0$  the system has an invariant distribution. (b) Find the invariant distribution when  $\delta = \mu$ .
4. Is the time-reversal of a tandem of  $M/M/1$  queues reversible at equilibrium? If not can you describe the time-reversal? Find the distribution of the process  $D_t$  which counts the number of customers whose service is completed by time  $t$ .
5. Consider the following queue. Customers arrive at rate  $\lambda > 0$  and are served by one server at rate  $\mu$ . After service, each customer returns to the beginning of the queue with probability  $p \in (0, 1)$ . Let  $(L_t)_{t \geq 0}$  denote the queue length. Show that  $L$  has the same distribution as a  $M/M/1$  queue with modified rates. For which parameters is  $L$  transient, and for which is it recurrent?
6. Let  $(L_t)_{t \geq 0}$  denotes the length of an  $M/M/1$  queue with rates  $\lambda < \mu$ . Let  $\pi$  denote the equilibrium distribution. Let  $(D_t)_{t \geq 0}$  denotes the departure process from the queue. By considering all possibilities leading to the events below, show directly that as  $h \rightarrow 0$ ,

$$\mathbb{P}_\pi(D_h - D_0 = 0) = 1 - \lambda h + o(h)$$

and that

$$\mathbb{P}_\pi(D_h - D_0 \geq 1) = \lambda h + o(h).$$

What have you proved?

7. Prove that the traffic equations for a Jackson network have a unique solution.
8. Let  $X_t = (X_t^1, \dots, X_t^N, t \geq 0)$  denote a Jackson network of  $N$  queues, with arrival rate  $\lambda_i$  and service rate  $\mu_i$  in queue  $i$ , and each customer moves to queue  $j \neq i$  with probability  $p_{ij}$  after service from queue  $i$ . We assume  $\sum_j p_{ij} < 1$  for each  $i = 1, \dots, N$  and that the traffic equations have a solution such that  $\bar{\lambda}_i < \mu_i$ .

Describe the time-reversal of  $X$  at equilibrium.

Let  $D_i(t)$  be the process of (final) departures from queue  $i$ . Show that, at equilibrium,  $(D_i(t), t \geq 0)_{1 \leq i \leq N}$  are independent Poisson processes and specify the rates. Show further that  $X_t$  is independent  $(D_i(s), 1 \leq i \leq N, 0 \leq s \leq t)$ .

9. Consider a system of  $N$  queues serving a finite number  $K$  of customers. The system evolves as follows. At station  $1 \leq i \leq N$ , customers are served one at a time at rate  $\mu_i$ . After

service, each customer moves to queue  $j$  with probability  $p_{ij} > 0$ . We assume here that the system is closed, ie,  $\sum_j p_{ij} = 1$  for all  $1 \leq i \leq N$ .

Let  $S = \{(n_1, \dots, n_N) : n_i \in \mathbb{N}, \sum_{i=1}^N n_i = K\}$  be the state space of the Markov chain. Write down its  $Q$ -matrix. Also write down the  $Q$ -matrix  $R$  corresponding to the position in the network of one customer (that is, when  $K = 1$ ). Show that there is a unique distribution  $(\lambda_i)_{1 \leq i \leq N}$  such that  $\lambda R = 0$ . Show that

$$\pi(n) = C_N \prod_{i=1}^N \lambda_i^{n_i}, n \in S$$

defines an invariant measure for the chain. Are the queue lengths independent at equilibrium?

**10.** Kafkaian Insurances Inc. has a peculiar way of processing claims. Claims arrive at a rate of 10 per day, and are initially randomly assigned to one of two departments, respectively  $D_1$  and  $D_2$ . The service rates in  $D_1$  and  $D_2$  are  $\mu_1 = 15$  and  $\mu_2 = 20$  per day, respectively. After looking at each claim, the relevant department settles the claim with probability 1/2, and otherwise finds a pretext to hand it over to the other department to process it. This goes on until the claim is finally settled by one of the two departments.

- (a) What proportion of claims is finally settled by  $D_1$ ?
- (b) How many claims are settled on average every month by Kafkaian Insurances Inc.?
- (c) The manager of the company wants to reward the work of his employees based on the number of claims that their department settles. Is that a good idea?

**11.** Consider a G/M/1 queueing system: the  $n$ th client arrives at time  $A_n = \sum_{i=1}^n \xi_i$ , where  $(\xi_i)$  is a sequence of nonnegative i.i.d. random variables, and the service times are i.i.d. exponential with rate  $\mu$ . Let  $X_n = L(A_n)$  be the size of the queue just before the  $n$ th arrival.

- (i) Show that  $(X_n)$  is a discrete-time Markov chain, and specify its transition matrix.
- (ii) Show that if  $\rho := (\mu \mathbb{E}A)^{-1} < 1$  then the chain  $(X_n)$  has a unique equilibrium distribution  $\pi = (\pi_i)$  and hence is positive recurrent. Here

$$\pi_i = (1 - \eta)\eta^i, \quad i = 0, 1, \dots$$

and  $\eta \in (0, 1)$  is a solution to  $\eta = \phi(\mu(\eta - 1))$ , where for  $\theta \in \mathbb{R}$ ,  $\phi(\theta) = \mathbb{E}(e^{\theta\xi})$ .

**12.** Consider the square lattice  $\mathbb{Z}^2$ , and endow each site  $x \in \mathbb{Z}^2$  with a weight  $W_x$ , which is an independent exponential random variable of rate  $\mu$ . An oriented path  $\pi$  between  $(1, 1)$  and a point  $(M, N)$ , with  $M, N \geq 1$ , is called increasing if it only ever goes in the North and East directions. Define the weight of an increasing path  $\pi$  to be  $W(\pi) = \sum_{x \in \pi} W_x$ , and the passage time from  $(1, 1)$  to  $(M, N)$  to be

$$T(M, N) = \max_{\pi} W(\pi)$$

where the max is over increasing  $\pi$ 's from  $(1, 1)$  to  $(M, N)$ . This model is called *Last Passage Percolation*. [Simulations showing optimal paths from  $(0,0)$  are interesting.]

The goal of this question is to relate this model to a sequence of  $N$  queues operating under the following protocol. At time 0 there are  $M$  customers in the first queue, and none at any other queue. Customers are served one at a time at rate  $\mu$  in each queue, and after service at queue  $i$ , a customer moves on to queue  $i + 1$ . Customers leave the system for good after being served at queue  $N$ . Let  $\tau(M, N)$  denote the time at which the  $M$ th customer completes service in queue  $N$ . Show that  $\tau(M, N)$  and  $T(M, N)$  have the same distribution.