

Number Fields: Example Sheet 2 of 3

1. Let K be a number field, and let $\mathfrak{a}$ and $\mathfrak{b}$ be non-zero ideals in $\mathcal{O}_K$. Determine the factorisations into prime ideals of $\mathfrak{a} + \mathfrak{b}$ and $\mathfrak{a} \cap \mathfrak{b}$ in terms of those for $\mathfrak{a}$ and $\mathfrak{b}$. Show that if $\mathfrak{a} + \mathfrak{b} = \mathcal{O}_K$ then $\mathfrak{a} \cap \mathfrak{b} = \mathfrak{a}\mathfrak{b}$ and there is an isomorphism of rings

$$\mathcal{O}_K/\mathfrak{a}\mathfrak{b} \cong \mathcal{O}_K/\mathfrak{a} \times \mathcal{O}_K/\mathfrak{b}.$$

2. Let $K = \mathbb{Q}(\sqrt{-5})$. Show by computing norms, or otherwise, that $\mathfrak{p} = (2, 1 + \sqrt{-5})$, $\mathfrak{q}_1 = (7, 3 + \sqrt{-5})$ and $\mathfrak{q}_2 = (7, 3 - \sqrt{-5})$ are prime ideals in $\mathcal{O}_K$. Which (if any) of the ideals $\mathfrak{p}$, $\mathfrak{q}_1$, $\mathfrak{q}_2$, $\mathfrak{p}^2$, $\mathfrak{p}\mathfrak{q}_1$, $\mathfrak{p}\mathfrak{q}_2$ and $\mathfrak{q}_1\mathfrak{q}_2$ are principal? Factor the principal ideal $(9 + 11\sqrt{-5})$ as a product of prime ideals.
3. Let K be a number field, and $\mathfrak{a} \subset \mathcal{O}_K$ a non-zero ideal. Let m be the least positive integer in $\mathfrak{a}$. Prove that m and $N\mathfrak{a}$ have the same prime factors.
4. Let $K = \mathbb{Q}(\sqrt{35})$ and $\omega = 5 + \sqrt{35}$. Verify the ideal equations $(2) = (2, \omega)^2$, $(5) = (5, \omega)^2$ and $(\omega) = (2, \omega)(5, \omega)$. Show that the class group of K contains an element of order 2. Find all ideals of norm dividing 100 and determine which are principal.
5. Let K be a number field, and $\mathfrak{a} = (x_1, \dots, x_k)$ an ideal in $\mathcal{O}_K$. Show that $N\mathfrak{a}$ divides $\gcd(N_{K/\mathbb{Q}}(x_1), \dots, N_{K/\mathbb{Q}}(x_k))$. Are these numbers always equal?
6. Let p be an odd prime and $K = \mathbb{Q}(\zeta_p)$ where ζ_p is a primitive p th root of unity. Determine $[K : \mathbb{Q}]$. Calculate $N_{K/\mathbb{Q}}(\pi)$ and $\text{Tr}_{K/\mathbb{Q}}(\pi)$ where $\pi = 1 - \zeta_p$.
- (i) By considering traces $\text{Tr}_{K/\mathbb{Q}}(\zeta_p^j \alpha)$ show that $\mathbb{Z}[\zeta_p] \subset \mathcal{O}_K \subset \frac{1}{p}\mathbb{Z}[\zeta_p]$.
- (ii) Show that $(1 - \zeta_p^r)/(1 - \zeta_p^s)$ is a unit for all $r, s \in \mathbb{Z}$ coprime to p , and that $\pi^{p-1} = u\pi$ where u is a unit.
- (iii) Prove that the natural map $\mathbb{Z} \rightarrow \mathcal{O}_K/(\pi)$ is surjective. Deduce that for any $\alpha \in \mathcal{O}_K$ and $m \geq 1$ there exist $a_0, \dots, a_{m-1} \in \mathbb{Z}$ such that

$$\alpha \equiv a_0 + a_1\pi + \dots + a_{m-1}\pi^{m-1} \pmod{\pi^m \mathcal{O}_K}.$$

- (iv) Deduce that $\mathcal{O}_K = \mathbb{Z}[\zeta_p]$.
7. Let $K = \mathbb{Q}(\sqrt{-d})$ where d is a positive square-free integer. Establish the following facts about the factorisation of principal ideals in $\mathcal{O}_K$.
- (i) If d is composite and p is an odd prime divisor of d then $(p) = \mathfrak{p}^2$ where $\mathfrak{p}$ is not principal.
- (ii) If $d \equiv 1$ or $2 \pmod{4}$ then $(2) = \mathfrak{p}^2$ where $\mathfrak{p}$ is not principal unless $d = 1$ or 2 .
- (iii) If $d \equiv 7 \pmod{8}$ then $(2) = \mathfrak{p}\bar{\mathfrak{p}}$ where $\mathfrak{p}$ is not principal unless $d = 7$.

Deduce that if K has class number 1 then either $d = 1, 2$ or 7 , or d is prime and $d \equiv 3 \pmod{8}$.

8. Let $K = \mathbb{Q}(\sqrt{-d})$ where $d > 1$ is the product of distinct primes $p_1, \dots, p_k$. Show that $(p_i) = \mathfrak{p}_i^2$ where $\mathfrak{p}_i = (p_i, \sqrt{-d})$. Show that just two of the ideals $\prod \mathfrak{p}_i^{r_i}$ with $r_i \in \{0, 1\}$ are principal. Deduce that the class group Cl_K contains a subgroup isomorphic to $(\mathbb{Z}/2\mathbb{Z})^{k-1}$. [If you like, just do the case $d \not\equiv 3 \pmod{4}$.]
9. Let $K = \mathbb{Q}(\theta)$ where θ is a root of $X^3 - 4X + 7$. Determine the ring of integers and discriminant of K . Determine the factorisation into prime ideals of $p\mathcal{O}_K$ for $p = 2, 3, 5, 7, 11$. Find all non-zero ideals $\mathfrak{a}$ of $\mathcal{O}_K$ with $N\mathfrak{a} \leq 11$.
10. Let $K = \mathbb{Q}(\alpha)$ where α is a root of $f(X) = X^3 + X^2 - 2X + 8$. [This polynomial is irreducible over $\mathbb{Q}$ and has discriminant -4×503 .]
 - (i) Show that $\beta = 4/\alpha \in \mathcal{O}_K$ and $\beta \notin \mathbb{Z}[\alpha]$. Deduce that $\mathcal{O}_K = \mathbb{Z}[\alpha, \beta]$.
 - (ii) Show that there is an isomorphism of rings $\mathcal{O}_K/2\mathcal{O}_K \cong \mathbb{F}_2 \times \mathbb{F}_2 \times \mathbb{F}_2$. Deduce that 2 splits completely in K .
 - (iii) Use Dedekind's criterion to show that $\mathcal{O}_K \neq \mathbb{Z}[\theta]$ for any θ .

The following extra questions may or may not be harder than the earlier questions.

11. Let $f(X) \in \mathbb{Z}[X]$ be a monic, irreducible polynomial, and let $K = \mathbb{Q}(\theta)$, where θ is a root of f .
 - (i) Show that if p is a prime not dividing $\text{Disc}(f)$ and $r \in \mathbb{Z}$ with $f(r) \equiv 0 \pmod{p}$, then there is a ring homomorphism $\mathcal{O}_K \rightarrow \mathbb{F}_p$ that sends θ to $r \pmod{p}$.
 - (ii) Suppose that $f(X) = X^3 - X - 1$. Show that θ is not a square in K .
 - (iii) Suppose instead that $f(X) = X^5 + 2X - 2$. Show that the equation $x^4 + y^4 + z^4 = \theta$ has no solutions with $x, y, z \in \mathcal{O}_K$.
12. Let K be a quadratic field and $\mathfrak{a} \subset \mathcal{O}_K$ an ideal. Show that $\mathfrak{a} = (\alpha, \beta)$ for some $\alpha \in \mathbb{Z}$ and $\beta \in \mathcal{O}_K$. Let $c = \gcd(\alpha^2, \alpha \text{Tr}\beta, N\beta)$. By computing the norm and trace show that $\frac{\alpha\beta}{c} \in \mathcal{O}_K$. Deduce that $(\alpha, \beta)(\alpha, \bar{\beta})$ is principal where $\bar{\beta}$ is the conjugate of β .
13. (Some applications of the Chinese Remainder Theorem, as proved in Question 1.) Let K be a number field and $\mathfrak{a} \subset \mathcal{O}_K$ a non-zero ideal. Let $\phi(\mathfrak{a}) = |(\mathcal{O}_K/\mathfrak{a})^*|$.
 - (i) Show that every ideal in the ring $\mathcal{O}_K/\mathfrak{a}$ is principal.
 - (ii) Deduce that every ideal in $\mathcal{O}_K$ can be generated by 2 elements.
 - (iii) Show that $\phi(\mathfrak{a}) = N(\mathfrak{a}) \prod_{\mathfrak{p}|\mathfrak{a}} (1 - \frac{1}{N\mathfrak{p}})$.
14. Let K be a number field, and let $p \in \mathbb{Z}$ be a prime. Write $(p) = \mathfrak{p}_1^{e_1} \dots \mathfrak{p}_r^{e_r}$ where $\mathfrak{p}_1, \dots, \mathfrak{p}_r$ are distinct prime ideals with $N\mathfrak{p}_i = p^{f_i}$.
 - (i) Let $\alpha \in \mathfrak{a} = \mathfrak{p}_1 \dots \mathfrak{p}_r$. Show that $\text{Tr}_{K/\mathbb{Q}}(\alpha) \equiv 0 \pmod{p}$.
 - (ii) Let $\theta_1, \dots, \theta_n$ be an integral basis for K , and $\alpha_1, \dots, \alpha_n$ a $\mathbb{Z}$ -basis for $\mathfrak{a}$. By considering the matrix with entries $\text{Tr}_{K/\mathbb{Q}}(\alpha_i \theta_j)$, show that D_K is divisible by $\prod_{i=1}^r p^{(e_i-1)f_i}$.