

**PART II REPRESENTATION THEORY  
SHEET 4**

Unless otherwise stated, all vector spaces are finite-dimensional over  $\mathbb{C}$ . In the first six questions we let  $G = \text{SU}(2)$ .

**1** Let  $V_n$  be the vector space of complex homogeneous polynomials of degree  $n$  in the variables  $x$  and  $y$ .

(a) Describe a representation  $\rho_n$  of  $G$  on  $V_n$  and show that it is irreducible. What is its character?

(b) Show that  $V_n$  is isomorphic to its dual  $V_n^*$ .

**2** (a) Decompose the representations  $V_4 \otimes V_3$  into irreducible  $G$ -spaces (that is, find a direct sum of irreducible representations which is isomorphic to  $V_4 \otimes V_3$ . In this and the following questions, you are not being asked to find such an isomorphism explicitly.)

(b) Repeat for  $V_3^{\otimes 2}$ ,  $\Lambda^2 V_3$  and  $S^2 V_3$ .

(c) How do  $V_1^{\otimes n}$ ,  $S^n V_1$ ,  $S^2 V_n$  and  $\Lambda^2 V_n$  decompose into irreducibles for  $n \geq 1$ ? What about  $S^3 V_2$ ?

**3** Let  $G$  act on the space  $M_3(\mathbb{C})$  of  $3 \times 3$  complex matrices, by conjugation:

$$A : X \mapsto A_1 X A_1^{-1},$$

where  $A_1$  is the  $3 \times 3$  block diagonal matrix with block diagonal entries  $A, 1, 1$ . Show that this gives a representation of  $G$  and decompose it into irreducibles.

**4** Let  $\chi_n$  be the character of the irreducible representation  $\rho_n$  of  $G$  on  $V_n$  of dimension  $n + 1$ .

Show that

$$\frac{1}{2\pi} \int_0^{2\pi} K(z) \chi_n \overline{\chi_m} d\theta = \delta_{nm},$$

where  $z = e^{i\theta}$  and  $K(z) = \frac{1}{2}(z - z^{-1})(z^{-1} - z)$ .

[ Note that all you need to know about integrating on the circle is orthogonality of characters:  $\frac{1}{2\pi} \int_0^{2\pi} z^n d\theta = \delta_{n,0}$ . This is really a question about Laurent polynomials. ]

**5** Check that the usual formula for integrating functions defined on  $S^3 \subseteq \mathbf{R}^4$  defines an  $G$ -invariant inner product on

$$G = \text{SU}(2) = \left\{ \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} : a\bar{a} + b\bar{b} = 1 \right\},$$

and normalize it so that the integral over the group is one.

**6** Compute the character of the representation  $S^n V_2$  of  $G$  for any  $n \geq 0$ . Calculate  $\dim_{\mathbb{C}}(S^n V_2)^G$  (by which we mean the subspace of  $S^n V_2$  where  $G$  acts trivially).

Deduce that the ring of complex polynomials in three variables  $x, y, z$  which are invariant under the action of  $\text{SO}(3)$  is a polynomial ring. Find a generator for this polynomial ring.

**7** (a) Let  $G$  be a compact group. Show that there is a continuous group homomorphism  $\rho : G \rightarrow O(n)$  if and only if  $G$  has an  $n$ -dimensional representation over  $\mathbb{R}$ . Here  $O(n)$  denotes the subgroup of  $GL_n(\mathbb{R})$  preserving the standard (positive definite) symmetric bilinear form. (b) Explicitly construct such a representation  $\rho : SU(2) \rightarrow SO(3)$  by showing that  $SU(2)$  acts on the vector space of matrices of the form

$$\left\{ A = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \in M_2(\mathbb{C}) : A + \overline{A}^t = 0 \right\}$$

by conjugation. Show that this subspace is isomorphic to  $\mathbb{R}^3$ , that  $(A, B) \mapsto -\text{tr}(AB)$  is a positive definite non-degenerate invariant bilinear form, and that  $\rho$  is surjective with kernel  $\{\pm I\}$ .

**8** The *Heisenberg group* of order  $p^3$  is the (non-abelian) subgroup

$$G = \left\{ \begin{pmatrix} 1 & a & x \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} : a, b, x \in \mathbb{F}_p \right\}.$$

of matrices over the finite field  $\mathbb{F}_p$  ( $p$  prime). Let  $H$  be the subgroup of  $G$  comprising matrices with  $a = 0$  and  $Z$  be the subgroup of  $G$  of matrices with  $a = b = 0$ .

(a) Show that  $Z = Z(G)$ , the centre of  $G$ , and that  $G/Z = \mathbb{F}_p^2$ . Note that this implies that the derived subgroup  $G'$  is contained in  $Z$ . [You can check by explicit computation that it equals  $Z$ , or you can deduce this from the list of irreducible representations found in (d) below.]

(b) Find all 1-dimensional representations of  $G$ .

(c) Let  $\psi : \mathbb{F}_p \rightarrow \mathbb{C}^\times$  be a non-trivial 1-dimensional representation of the cyclic group  $\mathbb{F}_p = \mathbb{Z}/p$ , and define a 1-dimensional representation  $\rho_\psi$  of  $H$  by

$$\rho_\psi \begin{pmatrix} 1 & 0 & x \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} = \psi(x).$$

Show that  $\text{Ind}_H^G \rho_\psi$  is an irreducible representation of  $G$ .

(d) Prove that the collection of representations constructed in (b) and (c) gives a complete list of all irreducible representations.

(e) Determine the character of the irreducible representation  $\text{Ind}_H^G \rho_\psi$ .

**9** Recall Sheet 3, q.8 where we used inner products to construct some irreducible characters  $\chi^{(n-r,r)}$  for  $S_n$ . Let  $n \in \mathbb{N}$ , and let  $\Omega$  be the set of all ordered pairs  $(i, j)$  with  $i, j \in \{1, 2, \dots, n\}$  and  $i \neq j$ . Let  $G = S_n$  act on  $\Omega$  in the obvious manner (namely,  $\sigma(i, j) = (\sigma i, \sigma j)$  for  $\sigma \in S_n$ ). Let's write  $\pi^{(n-2,1,1)}$  for the permutation character of  $S_n$  in this action.

Prove that

$$\pi^{(n-2,1,1)} = 1 + 2\chi^{(n-1,1)} + \chi^{(n-2,2)} + \psi,$$

where  $\psi$  is an irreducible character. Writing  $\psi = \chi^{(n-2,1,1)}$ , calculate the degree of  $\chi^{(n-2,1,1)}$ . Find its value on any transposition and on any 3-cycle. Returning to the character table of  $S_6$  calculated on Sheet 3, identify the character  $\chi^{(4,1,1)}$ .

**10** Recall the character table of  $G = \text{PSL}_2(7)$  from Sheet 2, q.9. Identify the columns corresponding to the elements  $x$  and  $y$  where  $x$  is an element of order 7 (eg the unitriangular matrix with 1 above the diagonal) and  $y$  is an element of order 3 (eg the diagonal matrix with entries 4 and 2).

The group  $G$  acts as a permutation group of degree 8 on the set of Sylow 7-subgroups (or the set of 1-dimensional subspaces of the vector space  $(\mathbb{F}_7)^2$ ). Obtain the permutation character of this action and decompose it into irreducible characters.

\*(Harder) Show that the group  $G$  is generated by an element of order 2 and an element of order 3 whose product has order 7.

[Hint: for the last part use the formula that the number of pairs of elements conjugate to  $x$  and  $y$  respectively, whose product is conjugate to  $t$ , equals  $c \sum \chi(x)\chi(y)\chi(t^{-1})/\chi(1)$ , where the sum runs over all the irreducible characters of  $G$ , and  $c = |G|^2(|C_G(x)||C_G(y)||C_G(t)|)^{-1}$ .]

SM, Lent Term 2014

Comments on and corrections to this sheet may be emailed to [sm@dpmms.cam.ac.uk](mailto:sm@dpmms.cam.ac.uk)