

Representation Theory — Examples Sheet 3

On this sheet all groups are finite and all representations are complex representations

1. Calculate S^2V and Λ^2V for the two-dimensional irreducible representations of D_8 and of Q_8 . Which has the trivial representation as a subrepresentation in each case?
2. Show both directly and using characters that if U, V and W are representations of G then

$$V^* \otimes W \cong \text{Hom}_k(V, W) \text{ and } \text{Hom}_k(V \otimes W, U) \cong \text{Hom}_k(V, \text{Hom}_k(W, U))$$

as representations of G . Deduce that if V is self-dual then either S^2V or Λ^2V contains a non-zero subrepresentation with trivial G -action.

3. Suppose $\rho: G \rightarrow GL(V)$ is an irreducible representation of G with character χ . By considering $V \otimes V, S^2V$ and Λ^2V show that

$$\frac{1}{|G|} \sum_{g \in G} \chi(g^2) = \begin{cases} 0 & \text{if } \chi \text{ is not real-valued} \\ \pm 1 & \text{if } \chi \text{ is real valued.} \end{cases}$$

Deduce that if $|G|$ is odd then G has only one real-valued irreducible character.

4. Let $\rho: G \rightarrow GL(V)$ be a representation of G of dimension d .
 - (a) Compute $\dim S^n V$ and $\dim \Lambda^n V$ for all n .
 - (b) Let $g \in G$ and $\lambda_1, \dots, \lambda_d$ be the eigenvalues of $\rho(g)$. What are the eigenvalues of g on $S^n V$ and $\Lambda^n V$?
 - (c) Let $f(t) = \det(tI - \rho(g))$ be the characteristic polynomial of $\rho(g)$. What is the relationship between the coefficients of f and $\chi_{\Lambda^n V}$?
 - (d) What is the relationship between $\chi_{S^n V}(g)$ and f ? (Hint: start with case $d = 1$).
5. Recall the character table of D_{10} from sheet 2. Explain how to view D_{10} as a subgroup of A_5 and then use induction from D_{10} to A_5 to reconstruct the character table of A_5 .
6. Obtain the character table of the dihedral group D_{2m} by using induction from the cyclic group C_m ; you will want to split into two cases according as m is odd or even.
7. Find all the characters of S_5 obtained by inducing irreducible representations of S_4 . Use these to reconstruct the character table of S_5 . Then repeat, replacing S_4 by the subgroup $\langle (12345), (2354) \rangle$ of S_5 of order 20.
8. Prove that if H is a subgroup of a group G , and K is a subgroup of H , and W is a representation of K then $\text{Ind}_K^G W \cong \text{Ind}_H^G \text{Ind}_K^H W$.
9. Let H be a subgroup of a group G . Show that for every irreducible representation (ρ, V) of G there is an irreducible representation (σ, W) of H such that ρ is an irreducible component of $\text{Ind}_H^G W$.
Deduce that if A is an abelian subgroup of G then every irreducible representation of G has dimension at most $|G/A|$.
10. Suppose that G is a Frobenius group with Frobenius kernel K . Show that if V is a non-trivial irreducible representation of K then $\text{Ind}_K^G V$ is also irreducible. Hence, explain how to construct the character table of G given the character tables of K and G/K .
11. Suppose that V is a faithful representation of a group G such that χ_V takes r distinct values. Show that each irreducible representation of G is a summand of $V^{\otimes n}$ for some $n < r$.
12. Suppose G is a finite group of odd order and with k conjugacy classes. Show that $|G| \equiv k \pmod{16}$.