

APPENDIX: ADEQUATE SUBGROUPS

ROBERT GURALNICK, FLORIAN HERZIG, RICHARD TAYLOR,
AND JACK THORNE

Let l be a prime, and let Γ be a finite subgroup of $\mathrm{GL}_n(\overline{\mathbb{F}}_l) = \mathrm{GL}(V)$. With these assumptions we say that *Condition (C) holds* if for every irreducible Γ -submodule $W \subset \mathrm{ad}^0 V$ there exists an element $g \in \Gamma$ with an eigenvalue α such that $\mathrm{tr} e_{g,\alpha} W \neq 0$. Here, $e_{g,\alpha}$ denotes the projection to the generalised α -eigenspace of g . This condition arises in the definition of adequacy in section 2.

Let Γ^{ss} denote the subset of Γ consisting of the elements that are semisimple (i.e. of order prime to l).

Lemma 1. *Suppose that Γ acts irreducibly on V . The following are equivalent.*

- (i) *Condition (C).*
- (ii) *For every irreducible submodule $W \subset \mathrm{ad}^0 V$ there exists $g \in \Gamma^{\mathrm{ss}}$ and $\alpha \in \overline{\mathbb{F}}_l$ such that $\mathrm{tr} e_{g,\alpha} W \neq 0$.*
- (iii) *The set Γ^{ss} spans $\mathrm{ad} V$ as an $\overline{\mathbb{F}}_l$ -vector space.*

Proof. Note that for any $g \in \Gamma$, Γ contains both its semisimple and unipotent parts g_s and g_u , respectively. (They are powers of g , as we work over $\overline{\mathbb{F}}_l$.) Since $e_{g,\alpha} = e_{g_s,\alpha}$ for all $g \in \Gamma$, the first two conditions are equivalent.

To show that the last two conditions are equivalent, let $Z \subset \mathrm{ad} V$ be the span of the semisimple elements in Γ . Let U denote the annihilator of Z under the (non-degenerate, Γ -invariant) trace pairing:

$$\begin{aligned} (1) \quad U &= \{w \in \mathrm{ad} V : \mathrm{tr}(gw) = 0 \quad \forall g \in \Gamma^{\mathrm{ss}}\} \\ (2) \quad &= \{w \in \mathrm{ad} V : \mathrm{tr}(e_{g,\alpha} w) = 0 \quad \forall g \in \Gamma^{\mathrm{ss}}, \alpha \in \overline{\mathbb{F}}_l\}, \end{aligned}$$

where we used that $e_{g,\alpha}$ is a polynomial in g and that $g = \sum \alpha e_{g,\alpha}$ for g semisimple.

The first author was partially supported by NSF grants DMS-0653873 and DMS-1001962. The second author was partially supported by NSF grant DMS-0902044 and agreement DMS-0635607. The third author was partially supported by NSF grant DMS-0600716 and by the IAS Oswald Veblen and Simonyi Funds. The second and third authors are grateful to the IAS for its hospitality during some of the work on this appendix.

Note that $U \subset \text{ad}^0 V$ by taking $g = 1$ in (1). From (2) it thus follows that the second condition is equivalent to $U = 0$. Equivalently, $Z = \text{ad } V$, which is the third condition. \square

Lemma 2.

- (i) *Suppose that Γ acts irreducibly on V . Condition (C) holds whenever Γ has order prime to l .*
- (ii) *Suppose that V, V' are finite-dimensional vector spaces over $\overline{\mathbb{F}}_l$ and that $\Gamma \subset \text{GL}(V), \Gamma' \subset \text{GL}(V')$ are finite subgroups that act irreducibly. If they both satisfy (C), then the image of $\Gamma \times \Gamma'$ in $\text{GL}(V \otimes V')$ also satisfies (C).*

Proof. By Burnside's theorem, Γ spans $\text{ad } V$. If Γ has order prime to l , then every element is semisimple, so the lemma above applies.

The second part of the proposition follows on noting that if g, h are semisimple elements then $g \otimes h$ is semisimple, and appealing to the third characterization of condition (C) in the lemma above. \square

Next we establish some preliminary results to prepare for our main theorem.

Lemma 3. *Suppose that T is a torus over \mathbb{F}_l . Let $X^* = X^*(T/\overline{\mathbb{F}}_l)$ and $X_* = X_*(T/\overline{\mathbb{F}}_l)$. There is a natural action of Frobenius Fr as an automorphism of X^* and X_* . Suppose that $\Delta_* \subset X_*$ is a finite subset that is stable under the action of Fr and spans $X_* \otimes \mathbb{Q}$.*

- (i) *If $\mu \in X^*$ with $|\langle \mu, \delta \rangle| < l - 1$ for all $\delta \in \Delta_*$ then $\mu(T(\mathbb{F}_l))$ is trivial iff $\mu = 0$.*
- (ii) *If V is a $T/\overline{\mathbb{F}}_l$ -module and all the weights μ of $T/\overline{\mathbb{F}}_l$ on V satisfy $|\langle \mu, \delta \rangle| < (l - 1)/2$ for all $\delta \in \Delta_*$ then the $\overline{\mathbb{F}}_l$ -span of $T(\mathbb{F}_l)$ in $\text{ad } V$ equals the $\overline{\mathbb{F}}_l$ -span of $T(\overline{\mathbb{F}}_l)$.*

Proof. We can identify $\text{Hom}(T(\mathbb{F}_l), \overline{\mathbb{F}}_l^\times)$ with $X^*/(l - \text{Fr})X^*$. To prove the first part, suppose that $|\langle \mu, \delta \rangle| < l - 1$ for $\delta \in \Delta_*$ and that $\mu(T(\mathbb{F}_l))$ is trivial, so $\mu = (l - \text{Fr})\lambda$. Choose δ_1 in Δ_* with $|\langle \lambda, \delta_1 \rangle|$ maximal. If $\langle \lambda, \delta_1 \rangle \neq 0$ then

$$l - 1 > |\langle \mu, \delta_1 \rangle| \geq l|\langle \lambda, \delta_1 \rangle| - |\langle \lambda, \text{Fr}^{-1} \delta_1 \rangle| \geq (l - 1)|\langle \lambda, \delta_1 \rangle| \geq l - 1,$$

a contradiction. Therefore $\langle \lambda, \delta_1 \rangle = 0$, so $\lambda = 0$ and $\mu = 0$. In particular we see that if μ_1 and μ_2 are two elements of X^* with $|\langle \mu_i, \delta \rangle| < (l - 1)/2$ for $\delta \in \Delta_*$ and $i = 1, 2$ then $\mu_1|_{T(\mathbb{F}_l)} = \mu_2|_{T(\mathbb{F}_l)}$ iff $\mu_1 = \mu_2$. The second part now follows since both subspaces of $\text{ad } V$ equal the $\overline{\mathbb{F}}_l$ -linear span of the $T/\overline{\mathbb{F}}_l$ -equivariant projectors onto the weight spaces of $T/\overline{\mathbb{F}}_l$ in V . \square

Lemma 4. *Suppose that G is a connected simply connected semisimple algebraic group over $\overline{\mathbb{F}}_l$ and $\phi : G \rightarrow \mathrm{GL}(V)$ a finite-dimensional representation. Let $G \supset B \supset T$ denote a Borel and maximal torus, and suppose that $|\langle \mu_1 - \mu_2, \alpha^\vee \rangle| < l$ for all weights μ_1, μ_2 of T on V and all simple roots α . Then there exist connected simply connected semisimple algebraic subgroups I and J of G such that $G = I \times J$, $\phi(J) = 1$, and ϕ induces a central isogeny of I onto its image \overline{I} , which is a semisimple algebraic group.*

Proof. Let J denote the connected component of the kernel of ϕ with its reduced scheme structure. Then J is smooth ([Mil], Proposition I.5.18). By Theorem 8.1.5 of [Spr09] and its proof, J is semisimple and there is a second semisimple algebraic group $I \subset G$ which commutes with J and such that $I \times J \rightarrow G$ is a central isogeny. It follows from the simply-connectedness of G that it is an isomorphism of $I \times J$ onto G . In particular, I and J are simply connected. Note that $T = T_I \times T_J$ and that $B = B_I \times B_J$ where (B_I, T_I) (resp. (B_J, T_J)) is a Borel and maximal torus in I (resp. J). (This follows from the fact that any smooth connected soluble subgroup of (resp. torus in) G is conjugate to a subgroup of B (resp. T).) Moreover $U = U_I \times U_J$, where U denotes the unipotent radical of B . Let \overline{I} denote the image of I under ϕ . Then \overline{I} is again reduced and connected and hence also smooth. In fact it is semisimple. (See Proposition 14.10(1)(c) of [Bor91].) The map ϕ factors through an isogeny $I \rightarrow \overline{I} \subset \mathrm{GL}(V)$. Let $\overline{B}, \overline{T}, \overline{U}$ denote the images of B_I, T_I, U_I in \overline{I} . Then these are all reduced and hence smooth. Moreover \overline{T} is a torus, \overline{B} is connected and soluble, \overline{U} is connected unipotent and $\overline{B} = \overline{T}\overline{U}$. As $\dim \overline{I} = \dim I = \dim T_I + 2 \dim U_I = \dim \overline{T} + 2 \dim \overline{U}$ we see that \overline{B} must be a Borel subgroup of \overline{I} with unipotent radical \overline{U} and that \overline{T} is a maximal torus in \overline{I} . The isogeny $I \rightarrow \overline{I}$ induces an l -morphism from the root datum of \overline{I} to the root datum of I . (See section 9.6.3 of [Spr09].) Then $I \rightarrow \overline{I}$ is a central isogeny, as otherwise T would have a weight occurring in $\mathrm{Lie} \overline{I} \subset \mathrm{ad} V$ of the form $l\mu$ with μ non-zero and this would contradict our assumption on the weights of T on V . \square

Suppose that we are given $\overline{\mathbb{F}}_l$ -vector spaces W_i with $\dim W_i \leq l$ for $i = 1, \dots, r$. Then the maps

$$\begin{aligned} \exp : X &\mapsto 1 + X + \frac{X^2}{2!} + \cdots + \frac{X^{l-1}}{(l-1)!} \\ \log : 1 + u &\mapsto u - \frac{u^2}{2} + \frac{u^3}{3} \pm \cdots - \frac{u^{l-1}}{l-1} \end{aligned}$$

define inverse bijections between the set of nilpotent elements in $\prod \text{End}(W_i)$ and the set of unipotent elements in $\prod \text{GL}(W_i)$.

Lemma 5. *Suppose that $G \subset \prod \text{GL}(W_i)$ is a connected reductive group over $\overline{\mathbb{F}}_l$ with $\dim W_i \leq l$ for all i . Let T be a maximal torus and U be the unipotent radical of a Borel subgroup of G that contains T . Suppose that $|\langle \mu_1 - \mu_2, \alpha^\vee \rangle| < l$ for all weights μ_1, μ_2 of T on $V = \bigoplus W_i$ and all simple roots α .*

- (i) *The maps \exp and \log induce inverse isomorphisms of varieties between $\text{Lie } U \subset \text{End}(V)$ and $U \subset \text{GL}(V)$.*
- (ii) *For any positive root α we have $\exp(\text{Lie } U_\alpha) = U_\alpha$.*
- (iii) *The map $\exp : \text{Lie } U \rightarrow U$ depends only on G and U , but not on V, W_i , or the representation $G \hookrightarrow \text{GL}(V)$.*
- (iv) *If θ is an automorphism of G that preserves T and U , then we have a commutative diagram:*

$$\begin{array}{ccc} \text{Lie } U & \xrightarrow{d\theta} & \text{Lie } U \\ \exp \downarrow & & \downarrow \exp \\ U & \xrightarrow{\theta} & U \end{array}$$

Proof. By the Lie–Kolchin theorem we may suppose U is contained in the group $U' = \prod U'_i$, where U'_i denotes the unipotent radical of a Borel subgroup of $\text{GL}(W_i)$. The maps \exp and \log provide mutually inverse isomorphisms of varieties between U' and $\text{Lie } U'$. It remains to show that $\exp \text{Lie } U = U$. Note that the product of any l elements of $\text{Lie } U'$ is zero. Thus the Zassenhaus formula (see [Mag54], section IV) tells us that to check that $\exp \text{Lie } U \subset U$ it suffices to check that for any root α we have $\exp(\text{Lie } U_\alpha) \subset U$. Let $x_\alpha : \mathbb{G}_a \rightarrow U_\alpha$ be the root homomorphism corresponding to α and let $X_\alpha = dx_\alpha(1) \in \text{Lie } U_\alpha$. Then formula II.1.19(6) of [Jan03] shows that for $a \in \overline{\mathbb{F}}_l$,

$$(3) \quad x_\alpha(a) = \sum_{n=0}^{l-1} a^n \frac{X_\alpha^n}{n!} = \exp(aX_\alpha)$$

in $\text{GL}(V)$, on noting that for $n < l$ we have $X_{\alpha,n} = X_\alpha^n/n!$ while $X_{\alpha,n}$ acts trivially on V for $n \geq l$. (This latter assertion follows from formula II.1.19(5) of [Jan03] because V_λ and $V_{\lambda+n\alpha}$ cannot both be non-zero.) Now by the Baker–Campbell–Hausdorff formula (see section IV.8 in part I of [Ser92]) and the fact that the product of any l elements of $\text{Lie } U'$ is zero we see that $\exp \text{Lie } U$ is a subgroup of U . As U is connected and smooth and $\dim \text{Lie } U \geq \dim U$ we deduce that $\exp \text{Lie } U = U$. This proves the first two parts.

The third part follows inductively from equation (3) and the Zassenhaus formula: fix a total order $<$ on the set of positive roots such that if $\alpha, \beta, \alpha + \beta$ are positive roots, then $\max(\alpha, \beta) < \alpha + \beta$. We induct on the positive root γ . Suppose that we know that \exp depends only on G and U on the subspace $\bigoplus_{\alpha > \gamma} \text{Lie } U_\alpha$. Then the same is true for $\exp(X + Y)$ for any $X \in \text{Lie } U_\gamma$ and $Y \in \bigoplus_{\alpha > \gamma} \text{Lie } U_\alpha$ by the Zassenhaus formula. (Note that $[\text{Lie } U_\alpha, \text{Lie } U_\beta] \subset \text{Lie } U_{\alpha+\beta}$ whenever α, β are positive roots.) This completes the proof of the third part.

The last part follows from the third part, by considering the representation $G \xrightarrow{\theta} G \hookrightarrow \text{GL}(V)$. \square

Lemma 6. *Suppose that G is a connected simply connected semisimple algebraic group over $\overline{\mathbb{F}}_l$. Suppose that $l > 3$ and that G has no simple factor isomorphic to SL_n with $l|n$. Let \mathfrak{g} denote the Lie algebra of G . Then \mathfrak{g} contains no non-trivial abelian ideal, and the natural map $\text{Aut}(G) \rightarrow \text{Aut}(\mathfrak{g})$ is a bijection. Moreover, a connected normal subgroup of G is preserved by an automorphism $\theta \in \text{Aut}(G)$ if and only if its Lie algebra is preserved by $d\theta \in \text{Aut}(\mathfrak{g})$.*

Here, $\text{Aut}(G)$ (resp., $\text{Aut}(\mathfrak{g})$) denotes the abstract group of automorphisms of the algebraic group G (resp., its Lie algebra \mathfrak{g}). In the proof we use Chevalley groups in the sense of Steinberg's Yale notes [Ste68b].

Proof. The universal Chevalley group over $\overline{\mathbb{F}}_l$ constructed using the complex semisimple Lie algebra \mathcal{L} of the same root system as G is an algebraic group isomorphic to G (see [Ste68b], §5). (In the notation of [Ste68b], we can let V be any representation whose weights span the weight lattice, so that $\mathcal{L}_{\mathbb{Z}} \subset \mathcal{L}$ is the \mathbb{Z} -lattice spanned by the fixed Chevalley basis H_i, X_α ; see Cor. 2 on p. 18 of [Ste68b].) In particular, $\mathfrak{g} \cong \mathcal{L}_{\mathbb{Z}} \otimes \overline{\mathbb{F}}_l$ (by the remark on p. 64 of [Ste68b]). Write $G = \prod G_i$ as a product of almost simple simply connected algebraic groups and correspondingly $\mathfrak{g} = \bigoplus \mathfrak{g}_i$. Then $Z(\mathfrak{g}_i) = 0$ by our assumption on l and G (see Theorem 2.3 in [Hur82]) and hence all \mathfrak{g}_i are simple ([Ste61], 2.6(5)). Moreover $\mathfrak{g}_i \cong \mathfrak{g}_j$ implies $G_i \cong G_j$ ([Ste61], 8.1). The G_i (resp., \mathfrak{g}_i) are uniquely characterised as the minimal non-trivial connected normal subgroups of G (resp., minimal non-trivial ideals of \mathfrak{g}), so they are permuted by automorphisms. Therefore if $\text{Aut}(G_i) \rightarrow \text{Aut}(\mathfrak{g}_i)$ is a bijection for all i , then so is $\text{Aut}(G) \rightarrow \text{Aut}(\mathfrak{g})$, and also the final claim of the proposition follows. (Note that any connected normal subgroup is a product of some of the G_i .) We can thus assume, without loss of generality, that G is almost simple.

Let G^{ad} denote the adjoint form of G . As G is the universal cover of G^{ad} and as $G^{\text{ad}} = G/Z(G)$, we have $\text{Aut}(G) = \text{Aut}(G^{\text{ad}})$. As $Z(\mathfrak{g}) = 0$

we see that the natural map $\mathfrak{g} \rightarrow \text{Lie } G^{\text{ad}}$ is an isomorphism. Thus it suffices to show that $\text{Aut}(G) = \text{Aut}(\mathfrak{g})$ whenever G is simple of *adjoint* type and $\mathfrak{g} = \text{Lie } G$. Thus we write G for G^{ad} from now on.

As an algebraic group G is isomorphic to the adjoint Chevalley group over $\overline{\mathbb{F}}_l$ (again by [Ste68b], §5). (In the notation of [Ste68b], we take V to be the adjoint representation \mathfrak{g} .) Thus we can identify $G(\overline{\mathbb{F}}_l)$ with the subgroup of $\text{GL}(\mathfrak{g})$ generated by the elements $x_\alpha(t) := \exp(\text{ad}(tX_\alpha))$, where $t \in \overline{\mathbb{F}}_l$ and α is any root. As each $\text{ad}(tX_\alpha)$ is a derivation of \mathfrak{g} , the group $G(\overline{\mathbb{F}}_l)$ is actually contained in $\text{Aut}(\mathfrak{g})$. For any $\eta \in \text{Aut}(\mathfrak{g})$, we have $\eta \circ \text{ad } X \circ \eta^{-1} = \text{ad}(\eta X)$ in $\text{GL}(\mathfrak{g})$. It follows that the natural action of $G(\overline{\mathbb{F}}_l) \subset \text{GL}(\mathfrak{g})$ on \mathfrak{g} agrees with the adjoint action of $G(\overline{\mathbb{F}}_l)$ on $\mathfrak{g} \subset \text{End}(\mathfrak{g})$.

The choice of Chevalley basis gives rise to a maximal torus T and a Borel B that contains it ([Ste68b], §5). From Theorem 9.6.2 in [Spr09] we deduce the following, using that G is adjoint. For each symmetry π of the Dynkin diagram \mathcal{D} there is a unique $\pi' \in \text{Aut}(G)$ that preserves (B, T) and that permutes the $x_{\alpha_i}(1) \in B$ according to π (where α_i are the simple roots). Moreover, $\text{Aut}(G)$ is the semidirect product of G (acting by inner automorphisms) and $\text{Aut}(\mathcal{D})$. Also, the elements of $\text{Aut}(\mathcal{D})$ biject with the “graph automorphisms” of \mathfrak{g} ([Ste61], §3).

The result now follows from ([Ste61], 4.2 and 4.5), as the group \mathfrak{H} in [Ste61] is actually contained in $G(\overline{\mathbb{F}}_l)$ since $\overline{\mathbb{F}}_l$ is algebraically closed (see Lemma 19 on p. 27 of [Ste68b]). (Note that the uniqueness statement in ([Ste61], 4.2) is incorrect and seems to be a typo.) \square

The following proposition may be of independent interest. The proof uses the classification of finite simple groups. Without it, the proof still goes through for l sufficiently large (depending on d and ineffective) by appealing to [LP] instead of [Gur99].

Proposition 7. *Suppose that V is a finite-dimensional $\overline{\mathbb{F}}_l$ -vector space and that $\Gamma \subset \text{GL}(V)$ is a finite subgroup that acts semisimply on V . Let $\Gamma^0 \subset \Gamma$ be the subgroup generated by elements of l -power order. Then V is a semisimple Γ^0 -module. Let $d \geq 1$ be the maximal dimension of an irreducible Γ^0 -submodule of V . Suppose that $l \geq 2(d+1)$. Then there exists an algebraic group G over $\overline{\mathbb{F}}_l$ and a semisimple representation $r : G_{/\overline{\mathbb{F}}_l} \rightarrow \text{GL}(V)$ with the following properties:*

- (i) *The connected component G^0 is semisimple, simply connected.*
- (ii) *$G \cong G^0 \rtimes H$, where H is a finite group of order prime to l .*
- (iii) *$r(G(\overline{\mathbb{F}}_l)) = \Gamma$.*

Moreover, if $T \subset G^0$ is a maximal torus and if μ is a weight of $T_{/\overline{\mathbb{F}}_l}$ on V then $\sum |\langle \mu, \alpha^\vee \rangle| < 2d$, where α ranges over the roots of $G_{/\overline{\mathbb{F}}_l}^0$. Also, Γ does not have any composition factor of order l .

Proof. Write $V = \bigoplus_i W_i$ as a direct sum of irreducible Γ^0 -modules. Since $\dim W_i \leq l$ for all i , we see that every element of l -power order in the image of $\Gamma^0 \rightarrow \mathrm{GL}(W_i)$ actually has order dividing l . Since $\Gamma^0 \hookrightarrow \prod \mathrm{GL}(W_i)$, we deduce that every element of Γ^0 of l -power order actually has order dividing l . Note that Γ/Γ^0 has order prime to l .

Step 1. We show that there exists a connected simply connected semisimple algebraic group G^0 over \mathbb{F}_l and a finite central subgroup $Z_0 \subset G^0(\mathbb{F}_l)$ with $G^0(\mathbb{F}_l)/Z_0 \cong \Gamma^0$. Let Γ_i denote the image of Γ^0 in $\mathrm{GL}(W_i)$. Note that Γ_i has no non-trivial normal subgroup of l -power order (since Γ_i acts faithfully on W_i , and an l -group acting on a non-zero $\overline{\mathbb{F}}_l$ -vector space has non-zero fixed points). So by Theorem B of [Gur99], Γ_i is a central product of quasisimple Chevalley groups. (Note that if $l = 11$ then $\dim W_i < 7$.) Now Γ^0 is a subgroup of $\prod \Gamma_i$ that surjects onto each factor, so $Z(\Gamma^0) = \Gamma^0 \cap \prod Z(\Gamma_i)$. Thus $\Gamma^0/Z(\Gamma^0)$ is a subgroup of $\prod \Gamma_i/Z(\Gamma_i)$, a product of simple Chevalley groups, that surjects onto each factor. By a theorem of Hall (Lemma 3.5 in [Kup]), $\Gamma^0/Z(\Gamma^0)$ is itself isomorphic to a direct product of simple Chevalley groups. It follows that $\Gamma^0 = [\Gamma^0, \Gamma^0]Z(\Gamma^0)$. Since Γ^0 is generated by elements of order l and $Z(\Gamma^0)$ is of order prime to l , it follows moreover that Γ^0 is perfect. Therefore Γ^0 is a perfect central extension of a product $\prod H_j$ of simple Chevalley groups H_j , so there exists a surjective homomorphism $\pi : \prod \tilde{H}_j \rightarrow \Gamma^0$ with central kernel, where \tilde{H}_j is the universal perfect central extension of H_j .

As $l > 3$ (to rule out Suzuki and Ree groups) there exist connected simply connected algebraic groups G_j over \mathbb{F}_l such that $H_j \cong G_j(\mathbb{F}_l)/Z(G_j(\mathbb{F}_l))$. (Note that G_j is the restriction of scalars of an absolutely almost simple algebraic group over a finite extension of \mathbb{F}_l .) Since $l > 3$ it is known that $\tilde{H}_j \cong G_j(\mathbb{F}_l)$ (see section 6.1 in [GLS98], particularly table 6.1.3). So we can take $G^0 = \prod G_j$ and $Z_0 = \ker \pi$.

Since $\Gamma^0/Z(\Gamma^0)$ is a product of nonabelian simple groups and since $Z(\Gamma^0)$ and Γ/Γ^0 are of order prime to l , it follows that Γ does not have any composition factor of order l .

Let $G^0 \supset B \supset T$ denote a Borel and maximal torus defined over \mathbb{F}_l .

Step 2. We lift V to a $G_{/\overline{\mathbb{F}}_l}^0$ -module and compare the actions of $T(\mathbb{F}_l)$ and $T(\overline{\mathbb{F}}_l)$ on V . Let U denote the unipotent radical of B and set $N = N_{G^0}(T)$. Let B^{op} denote the opposite Borel subgroup to B containing T and let U^{op} denote its unipotent radical. (See Theorem 14.1 of [Bor91].)

By uniqueness we see it is defined over \mathbb{F}_l .) Let $X = X^*(T_{/\mathbb{F}_l})$ with its subset Φ of roots and Φ^+ (resp. Δ) the set of positive (resp. simple) roots corresponding to B . Let $X^+ \subset X$ be the subset of dominant weights. There is a semisimple algebraic action of $G_{/\mathbb{F}_l}^0$ on V , say $\phi : G_{/\mathbb{F}_l}^0 \rightarrow \mathrm{GL}(V)$, such that:

- (i) the highest weight λ of a simple submodule is restricted (i.e. $0 \leq \langle \lambda, \alpha^\vee \rangle < l$ for all $\alpha \in \Delta$),
- (ii) the action of $G^0(\mathbb{F}_l)$ is the one induced by the map $G^0(\mathbb{F}_l) \rightarrow \Gamma^0$,
- (iii) the subspaces W_i are $G_{/\mathbb{F}_l}^0$ -stable.

(This follows from a result of Steinberg: see Theorem 2.11 in [Hum06]. Note that [Hum06] works with an algebraic group \mathbf{G} that is simple, but the proof given does not depend on that assumption.) By Proposition 3 of [Ser94] we see that if λ in X^+ is a weight of $T_{/\mathbb{F}_l}$ on V then $\sum_{\alpha \in \Phi^+} \langle \lambda, \alpha^\vee \rangle < d$; in particular, $\langle \lambda, \alpha^\vee \rangle < (l-1)/2$ for all $\alpha \in \Phi^+$. (Note that $\dim W_i \leq (l-1)/2$ and that the proof of that proposition does not require that $G_{/\mathbb{F}_l}^0$ be almost simple.) If μ is a weight of $T_{/\mathbb{F}_l}$ on V then we see that there is w in the Weyl group with $w\mu \in X^+$ and $0 \leq \langle w\mu, \alpha^\vee \rangle < (l-1)/2$ for all $\alpha \in \Phi^+$, and we deduce that $|\langle \mu, \alpha^\vee \rangle| < (l-1)/2$ for all $\alpha \in \Phi$. We also deduce that if μ is a weight of $T_{/\mathbb{F}_l}$ on $\mathrm{ad} V$ then $|\langle \mu, \alpha^\vee \rangle| < l-1$ for all $\alpha \in \Delta$.

Step 3. The semisimple group $\bar{I} \subset \mathrm{GL}(V)$ and its simply connected cover $I \subset G_{/\mathbb{F}_l}^0$. Since $|\langle \mu, \alpha^\vee \rangle| < l/2$ for all weights μ of $T_{/\mathbb{F}_l}$ on V and all $\alpha \in \Delta$ we may apply Lemma 4 to $\phi : G_{/\mathbb{F}_l}^0 \rightarrow \mathrm{GL}(V)$. We obtain connected simply connected semisimple algebraic subgroups I, J of $G_{/\mathbb{F}_l}^0$ such that $G_{/\mathbb{F}_l}^0 = I \times J$, $\phi(J) = 1$, and ϕ induces a central isogeny of I onto its image \bar{I} , which is a semisimple algebraic group. Note that $T_{/\mathbb{F}_l} = T_I \times T_J$ and that $B_{/\mathbb{F}_l} = B_I \times B_J$ where (B_I, T_I) (resp. (B_J, T_J)) is a Borel and maximal torus in I (resp. J). Moreover $U_{/\mathbb{F}_l} = U_I \times U_J$. Let $\bar{B}, \bar{T}, \bar{U}, \bar{B}^{\mathrm{op}}, \bar{U}^{\mathrm{op}}$ denote the images of $B_I, T_I, U_I, B_I^{\mathrm{op}}, U_I^{\mathrm{op}}$ in \bar{I} . Then \bar{T} is a maximal torus of \bar{I} , and $\bar{B}, \bar{B}^{\mathrm{op}}$ are opposite Borel subgroups containing it. Also $\bar{U}, \bar{U}^{\mathrm{op}}$ are the unipotent radicals of $\bar{B}, \bar{B}^{\mathrm{op}}$. Since $I \rightarrow \bar{I}$ is a central isogeny, $U_I \rightarrow \bar{U}$ and $U_I^{\mathrm{op}} \rightarrow \bar{U}^{\mathrm{op}}$ are isomorphisms.

Step 4. The maps \log and \exp provide inverse isomorphisms of varieties between $\bar{U} \subset \mathrm{GL}(V)$ and $\mathrm{Lie} \bar{U} \subset \mathrm{ad} V$. This follows from Lemma 5 applied to $\bar{I} \subset \mathrm{GL}(V)$ since $\dim W_i \leq l$ for all i and $|\langle \mu, \alpha^\vee \rangle| < l/2$ for all weights μ of $T_{/\mathbb{F}_l}$ on V and all $\alpha \in \Delta$. (Note that $T_I \rightarrow \bar{T}$

induces a bijection on coroots since $I \rightarrow \bar{I}$ is a central isogeny; thus $T \rightarrow \bar{T}$ induces a surjection on coroots.)

Step 5. The $\bar{\mathbb{F}}_l$ -span of $\log U(\mathbb{F}_l)$ is $\text{Lie } \bar{U}$. Since $d\phi : \text{Lie } U \rightarrow \text{Lie } \bar{U}$ is surjective, it suffices to show that there is an isomorphism $\log : U \rightarrow \text{Lie } U$ defined over \mathbb{F}_l such that $d\phi \circ \log = \log \circ \phi$. Pick an \mathbb{F}_l -structure on V . The map $G^0_{/\mathbb{F}_l} \rightarrow \text{GL}(V)$ can be defined over some \mathbb{F}_{l^s} and so taking restrictions of scalars from \mathbb{F}_{l^s} to \mathbb{F}_l we get an \mathbb{F}_l -vector space V' and a map $\psi : G^0 \rightarrow \text{GL}(V')$. The map $G^0_{/\mathbb{F}_l} \rightarrow \text{GL}(V)$ is obtained from ψ by extending scalars to $\bar{\mathbb{F}}_l$ and projecting to a direct summand V of $V' \otimes \bar{\mathbb{F}}_l$. The dimension of all irreducible factors of $V' \otimes \bar{\mathbb{F}}_l$ is at most l . Moreover for any weight λ of $T_{/\bar{\mathbb{F}}_l}$ on $V' \otimes \bar{\mathbb{F}}_l$ we have $|\langle \lambda, \alpha^\vee \rangle| < (l-1)/2$ for all $\alpha \in \Phi^+$.

By Lemma 4 we see that $\psi : G^0 \rightarrow \text{GL}(V')$ is a central isogeny onto its image. (By construction we have $(\ker \psi)(\mathbb{F}_l) = Z_0$. Suppose that $\ker \psi$ is not finite. Then it has to contain one of the \mathbb{F}_l -almost simple factors of $G^0 = \prod G_j$. But $G_j(\mathbb{F}_l)$ is nonabelian.)

In particular, ψ induces an isomorphism $U \rightarrow \psi(U)$. Then Lemma 5 (applied to the image of $\psi_{/\bar{\mathbb{F}}_l}$) gives the desired map $\log : U \rightarrow \text{Lie } U \subset \text{ad } V'$.

Step 6: Some properties of $G^0(\mathbb{F}_l)$. The pair $(B(\mathbb{F}_l), N(\mathbb{F}_l))$ is a split BN pair in $G^0(\mathbb{F}_l)$ (see section 1.18 of [Car93]). Also $U(\mathbb{F}_l)$ is a Sylow l -subgroup of $G^0(\mathbb{F}_l)$ and $B(\mathbb{F}_l) = N_{G^0(\mathbb{F}_l)}(U(\mathbb{F}_l)) = N_{G^0(\mathbb{F}_l)}(B(\mathbb{F}_l))$ (see Proposition 2.5.1 of [Car93]).

Moreover $T(\mathbb{F}_l)$ is a Sylow l -complement in $B(\mathbb{F}_l)$. Note that $U^{\text{op}}(\mathbb{F}_l)$ is $N(\mathbb{F}_l)$ -conjugate to $U(\mathbb{F}_l)$. (The longest Weyl element w_0 is stable under Frobenius, hence represented by an element $n_0 \in N(\mathbb{F}_l)$. Then use that $U^{\text{op}} = n_0 U n_0^{-1}$.) Moreover the second-last displayed equation on page 74 (section 2.9) of [Car93] shows that $U^{\text{op}}(\mathbb{F}_l)$ is the unique $N(\mathbb{F}_l)$ -conjugate of $U(\mathbb{F}_l)$ with trivial intersection with $U(\mathbb{F}_l)$.

Step 7. We have $N(\mathbb{F}_l) = N_{G^0(\mathbb{F}_l)}(T(\mathbb{F}_l))$ so that $N_{G^0(\mathbb{F}_l)}(T(\mathbb{F}_l)) \cap N_{G^0(\mathbb{F}_l)}(B(\mathbb{F}_l)) = T(\mathbb{F}_l)$ and $Z_0 \subset Z(G^0(\mathbb{F}_l)) \subset T(\mathbb{F}_l)$.

Suppose that g is in $N_{G^0(\mathbb{F}_l)}(T(\mathbb{F}_l))$. One can write g uniquely as unu' where $u \in U(\mathbb{F}_l)$, $n \in N(\mathbb{F}_l)$ maps to w_n in the Weyl group and $u' \in U_{w_n}$ in the notation of Theorem 2.5.14 of [Car93]. Then for any h in $T(\mathbb{F}_l)$ we can find h' and h'' in $T(\mathbb{F}_l)$ such that

$$hunu' = unu'h' \quad \text{and} \quad h''unu' = unu'h,$$

i.e.,

$$(huh^{-1})(hn)u' = u(nh')(h'^{-1}u'h')$$

and

$$(h''uh''^{-1})(h''n)u' = u(nh)(h^{-1}u'h).$$

As $T(\mathbb{F}_l)$ normalizes $U(\mathbb{F}_l)$ and U_{w_n} and as $w_{nh} = w_n = w_{hn}$ the uniqueness assertion of Theorem 2.5.14 of [Car93] tells us that $huh^{-1} = u$ and $u' = h^{-1}u'h$. Thus $u \in Z_{U(\mathbb{F}_l)}(T(\mathbb{F}_l))$ and $u' \in Z_{U_{w_n}}(T(\mathbb{F}_l)) \subset Z_{U(\mathbb{F}_l)}(T(\mathbb{F}_l))$. So it suffices to prove that $Z_{U(\overline{\mathbb{F}}_l)}(T(\mathbb{F}_l)) = 1$. By Proposition 8.2.1 in [Spr09]

it suffices to show that $Z_{U_\alpha(\overline{\mathbb{F}}_l)}(T(\mathbb{F}_l)) = 1$ for all $\alpha \in \Phi^+$. By Proposition 8.1.1(i) in [Spr09]

it suffices that α is non-trivial on $T(\mathbb{F}_l)$ for all $\alpha \in \Phi^+$. As $l \geq 5$, this follows from Lemma 3(i) (applied with Δ_* the set of simple coroots).

Step 8. We find a subgroup H of order prime to l such that $\Gamma = \Gamma^0 H$. Let H denote the subgroup of $h \in \Gamma$ which normalize both the image of $B(\mathbb{F}_l)$ and the image of $T(\mathbb{F}_l)$ in Γ^0 . Then by the previous paragraph we see that $H \cap \Gamma^0$ is $T(\mathbb{F}_l)/Z_0$. Thus H has order prime to l .

Moreover if $\gamma \in \Gamma$ we see that $\gamma(B(\mathbb{F}_l)/Z_0)\gamma^{-1}$ is the normalizer of a Sylow l -subgroup of $G^0(\mathbb{F}_l)/Z_0$ and hence $G^0(\mathbb{F}_l)$ -conjugate to $B(\mathbb{F}_l)/Z_0$, say $\gamma(B(\mathbb{F}_l)/Z_0)\gamma^{-1} = k(B(\mathbb{F}_l)/Z_0)k^{-1}$ with $k \in G^0(\mathbb{F}_l)$. Then $k^{-1}\gamma(T(\mathbb{F}_l)/Z_0)\gamma^{-1}k$ is a Sylow l -complement in $B(\mathbb{F}_l)/Z_0$ and hence (by Hall's theorem) $B(\mathbb{F}_l)/Z_0$ -conjugate to $T(\mathbb{F}_l)/Z_0$, say

$$k^{-1}\gamma(T(\mathbb{F}_l)/Z_0)\gamma^{-1}k = k'(T(\mathbb{F}_l)/Z_0)k'^{-1}$$

for some $k' \in B(\mathbb{F}_l)$. Then $(kk')^{-1}\gamma$ lies in H and we deduce that Γ is generated by H and $G^0(\mathbb{F}_l)/Z_0 = \Gamma^0$.

Step 9. Lifting the conjugation action of H on Γ^0 to G^0 . We first show that $G^0_{/\overline{\mathbb{F}}_l}$ has no simple factor SL_n with $l|n$ by showing that any such factor would act trivially on $V = \bigoplus W_i$, contradicting that $G^0(\mathbb{F}_l)/Z_0$ acts faithfully. So suppose that $\mathrm{SL}_{n/\overline{\mathbb{F}}_l}$ has an irreducible module of dimension less than $l-1$. Then by Proposition 3 in [Ser94] its highest weight λ would satisfy $\sum \langle \lambda, \alpha^\vee \rangle < l-1$, where α runs through the set of positive roots. A calculation shows that the left-hand side is at least $n-1$ if λ is non-zero. So if $n \geq l$, then $\lambda = 0$.

Next we claim that $d\phi : (\mathrm{Lie} G^0)(\overline{\mathbb{F}}_l) \rightarrow \mathrm{ad} V$ is injective on the subspace $(\mathrm{Lie} G^0)(\mathbb{F}_l)$. Note first that it is injective on $(\mathrm{Lie} U)(\mathbb{F}_l)$ as ϕ is injective on $U(\mathbb{F}_l)$. (Consider the isomorphism $\log : U(\mathbb{F}_l) \rightarrow (\mathrm{Lie} U)(\mathbb{F}_l)$ constructed in Step 5.) Similarly $d\phi$ is injective on $(\mathrm{Lie} U^{\mathrm{op}})(\mathbb{F}_l)$. Since ϕ maps U to \overline{U} , T to \overline{T} , U^{op} to $\overline{U}^{\mathrm{op}}$, and since $\mathrm{Lie} G^0 = \mathrm{Lie} U \oplus \mathrm{Lie} T \oplus \mathrm{Lie} U^{\mathrm{op}}$, $\mathrm{Lie} \overline{G} = \mathrm{Lie} \overline{U} \oplus \mathrm{Lie} \overline{T} \oplus \mathrm{Lie} \overline{U}^{\mathrm{op}}$ it follows that the kernel of $d\phi$ on $(\mathrm{Lie} G^0)(\mathbb{F}_l)$ is contained in $(\mathrm{Lie} T)(\mathbb{F}_l)$. But $(\mathrm{Lie} G^0)(\overline{\mathbb{F}}_l)$ contains no non-trivial abelian ideal by Lemma 6. This proves the claim.

Note that H acts by conjugation on $\mathrm{GL}(V)$ and $\mathrm{ad} V$, in particular it preserves the Lie algebra structure of $\mathrm{ad} V$. By definition H stabilises the image of $U(\mathbb{F}_l)$ in $\mathrm{GL}(V)$ and hence by Step 5 it also

stabilises $\log U(\mathbb{F}_l) = d\phi((\text{Lie } U)(\mathbb{F}_l))$. Because $U^{\text{op}}(\mathbb{F}_l)$ is the unique $N_{G^0(\mathbb{F}_l)}(T(\mathbb{F}_l))$ -conjugate of $U(\mathbb{F}_l)$ that has trivial intersection with $U(\mathbb{F}_l)$, it is also stabilised by H . The previous argument then shows that H stabilises $d\phi((\text{Lie } U^{\text{op}})(\mathbb{F}_l))$. Since $[\text{Lie } U, \text{Lie } U^{\text{op}}] = \text{Lie } G^0$ (as we may check over $\overline{\mathbb{F}}_l$), it follows that H stabilises the image of $(\text{Lie } G^0)(\mathbb{F}_l)$ in $\text{ad } V$. By extending scalars, we get a natural action of H on $(\text{Lie } G^0)(\overline{\mathbb{F}}_l)$. This action lifts uniquely to an action on $G_{/\overline{\mathbb{F}}_l}^0$ by Lemma 6.

We claim that with respect to the H -action on $G_{/\overline{\mathbb{F}}_l}^0$ just constructed, $\phi : G_{/\overline{\mathbb{F}}_l}^0 \rightarrow \text{GL}(V)$ is H -equivariant. We first show that the conjugation action of H on $\text{GL}(V)$ stabilises \overline{I} . If $h \in H$ then h sends $U(\mathbb{F}_l)$ to itself and hence $\log U(\mathbb{F}_l)$ to itself and hence $\text{Lie } \overline{U}$ to itself and hence \overline{U} to itself. Similarly h stabilises \overline{U}^{op} . As the root subgroups generate \overline{I} (by Theorem 8.1.5 in [Spr09]), we see that h indeed stabilises \overline{I} . This action of H on \overline{I} lifts uniquely to an action on the simply connected cover I of \overline{I} . (For existence use Theorem 9.6.5 of [Spr09] and the conjugation action of T_l . For uniqueness use the semisimplicity of I .) On the other hand, Lemma 6 shows that the H -action on $G_{/\overline{\mathbb{F}}_l}^0$ respects the decomposition $G_{/\overline{\mathbb{F}}_l}^0 = I \times J$. Since J is killed by ϕ it suffices to show that the two H -actions on I (one coming from \overline{I} and one from $G_{/\overline{\mathbb{F}}_l}^0$) agree. By Lemma 6 we can check this on the Lie algebra. The same lemma shows that $d\phi : \text{Lie } I \rightarrow \text{Lie } \overline{I}$ is an isomorphism, since $\text{Lie } I$ contains no non-trivial abelian ideal. By construction both H -actions on $\text{Lie } I$ are compatible with the H -action on $\text{Lie } \overline{I}$, so the two H -actions on I indeed agree. Therefore ϕ is H -equivariant. A fortiori, it extends to a homomorphism $G_{/\overline{\mathbb{F}}_l}^0 \rtimes H \rightarrow \text{GL}(V)$.

Finally we show that the H -action on $G_{/\overline{\mathbb{F}}_l}^0$ descends to G^0 . Suppose that $h \in H$ and $\sigma \in \text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_l)$. The automorphism $\sigma h \sigma^{-1} h^{-1}$ is trivial on $(\text{Lie } G^0)(\mathbb{F}_l)$, hence trivial on $(\text{Lie } G^0)(\overline{\mathbb{F}}_l)$, hence trivial on $G_{/\overline{\mathbb{F}}_l}^0$ by Lemma 6. Therefore the H -action indeed descends to G^0 .

By construction, the image of $G^0(\mathbb{F}_l) \rtimes H$ is Γ . Let $G = G^0 \rtimes H$ and $r : G_{/\overline{\mathbb{F}}_l} \rightarrow \text{GL}(V)$ the homomorphism we just obtained. It remains to show that r is semisimple. But this follows from Lemma 5(b) in [Ser94] since the restriction of r to $G_{/\overline{\mathbb{F}}_l}^0$ is semisimple and $(G : G^0)$ is prime to l . \square

We remark that for the purpose of proving Theorem 9 we do not need an H -action on G^0 , we only need an H -action on $G_{/\overline{\mathbb{F}}_l}^0$ that is compatible with the H -action on $\text{GL}(V)$. Since $G_{/\overline{\mathbb{F}}_l}^0 = I \times J$, we can

lift the H -action on \bar{I} to I as above and let H act arbitrarily on J ; for this it is not necessary to appeal to Lemma 6.

Lemma 8. *Suppose that G is a linear algebraic group over $\bar{\mathbb{F}}_l$ such that the connected component G^0 is semi-simple and simply connected and such that l does not divide $(G : G^0)$. Let $G^0 \supset B \supset T$ denote a Borel subgroup and a maximal torus and let \mathcal{T} denote the normalizer of the pair (B, T) in G . Then the $G^0(\bar{\mathbb{F}}_l)$ -conjugates of $\mathcal{T}(\bar{\mathbb{F}}_l)$ equal the semisimple elements of $G(\bar{\mathbb{F}}_l)$ and they are Zariski dense in G . In particular, if V is an irreducible representation of G then the $G^0(\bar{\mathbb{F}}_l)$ -conjugates of $\mathcal{T}(\bar{\mathbb{F}}_l)$ span $\text{ad } V$ over $\bar{\mathbb{F}}_l$.*

Proof. By Theorem 7.5 in [Ste68a] every semisimple element of $G(\bar{\mathbb{F}}_l)$ is $G^0(\bar{\mathbb{F}}_l)$ -conjugate to an element of $\mathcal{T}(\bar{\mathbb{F}}_l)$. The converse is clear as $\mathcal{T} \cap G^0 = T$, an element $g \in G(\bar{\mathbb{F}}_l)$ is semisimple iff g is of order prime to l , and l does not divide $(G : G^0)$. Next we have $G = G^0\mathcal{T}$ since Borel subgroups in G^0 are conjugate and maximal tori in B are conjugate. Consider a fixed coset G^0h with $h \in \mathcal{T}(\bar{\mathbb{F}}_l)$. By Lemma 4 of [Spr06] the elements $g(th)g^{-1} = [gt(hgh^{-1})^{-1}]h$ of G^0h , where t runs over $T(\bar{\mathbb{F}}_l)$ and g runs over $G^0(\bar{\mathbb{F}}_l)$, are Zariski dense in G^0h . (Lemma 4 of [Spr06] does not immediately apply to h as h is not a diagram automorphism. However for some $s \in T(\bar{\mathbb{F}}_l)$ the automorphism $g \mapsto shgh^{-1}s^{-1}$ is a diagram automorphism and hence the elements $gt(hgh^{-1})^{-1} = gts^{-1}(shgh^{-1}s^{-1})^{-1}s$ as t runs over $T(\bar{\mathbb{F}}_l)$ and g runs over $G^0(\bar{\mathbb{F}}_l)$ are Zariski dense in G^0 .) Thus the $G^0(\bar{\mathbb{F}}_l)$ -conjugates of $\mathcal{T}(\bar{\mathbb{F}}_l)$ are Zariski dense in $G(\bar{\mathbb{F}}_l)$. For the last claim note that if $\text{tr}(gw) = 0$ for some $w \in \text{ad } V$ and some Zariski dense subset of $g \in G(\bar{\mathbb{F}}_l)$, then $w = 0$. \square

The proof of our main theorem relies on Proposition 7 and thus on the classification of finite simple groups. (It still holds without it for l sufficiently large, depending on d and ineffective, due to the results of Larsen and Pink [LP].)

Theorem 9. *Suppose that V is a finite-dimensional $\bar{\mathbb{F}}_l$ -vector space and that $\Gamma \subset \text{GL}(V)$ is a finite subgroup that acts irreducibly on V . Let $\Gamma^0 \subset \Gamma$ be the subgroup generated by elements of l -power order. Then V is a semisimple Γ^0 -module. Let $d \geq 1$ be the maximal dimension of an irreducible Γ^0 -submodule of V . Suppose that $l \geq 2(d + 1)$. Then:*

- (i) $H^0(\Gamma, \text{ad}^0 V) = H^1(\Gamma, \text{ad}^0 V) = H^1(\Gamma, \bar{\mathbb{F}}_l) = 0$.
- (ii) *The set Γ^{ss} spans $\text{ad } V$ as an $\bar{\mathbb{F}}_l$ -vector space.*

In particular, for any finite subfield k of $\bar{\mathbb{F}}_l$ containing the eigenvalues of all elements of Γ and such that $\Gamma \subset \text{GL}_n(k)$, Γ is adequate.

Proof. Write $V = \bigoplus_i W_i$ as a direct sum of irreducible Γ^0 -modules. Note that Γ/Γ^0 has order prime to l .

We claim that $\dim V$ is prime to l . Let U be an irreducible constituent of V as a Γ^0 -module and let V' be the U -isotypic direct summand of V . Since Γ acts transitively on the set of isotypic components and as $(\Gamma : \Gamma^0)$ is prime to l , it suffices to show that $\dim V'$ is prime to l . Let $\Gamma' \supset \Gamma^0$ be the stabiliser of V' . Then V' is an irreducible Γ' -module. By Theorem 51.7 in [CR62], U extends to a projective representation of Γ' and there is an irreducible projective representation U' of Γ'/Γ^0 such that $V' \cong U \otimes U'$ (as projective Γ' -representation). The claim follows as $\dim U < l$ and Γ'/Γ^0 is of order prime to l .

By Proposition 7 there exists an algebraic group $G = G^0 \rtimes H$ over \mathbb{F}_l and a semisimple representation $r : G_{/\overline{\mathbb{F}}_l} \rightarrow \mathrm{GL}(V)$, where G^0 is connected simply connected semisimple, H is a finite group of order prime to l , and $r(G(\mathbb{F}_l)) = \Gamma$. Moreover Γ has no composition factor of order l , which implies that no quotient of Γ^0 contains a non-trivial normal l -subgroup.

We have

$$H^1(\Gamma, \mathrm{ad} V) = \bigoplus_{i,j} H^1(\Gamma^0, \mathrm{Hom}(W_i, W_j))^\Gamma$$

and

$$H^1(\Gamma^0, \mathrm{Hom}(W_i, W_j)) = \mathrm{Ext}_{\Gamma^0}^1(W_i, W_j),$$

which vanishes by [Gur99], Theorem A, since $\dim W_i + \dim W_j \leq l - 2$. (We apply that theorem to the quotient of Γ^0 that acts faithfully. Note that we saw above that this quotient does not have a non-trivial normal l -subgroup.) Similarly, $2 \leq l - 2$ implies that $H^1(\Gamma, \overline{\mathbb{F}}_l) = 0$. Since $\dim V$ is prime to l it follows that $H^0(\Gamma, \mathrm{ad}^0 V) = 0$ and that $\mathrm{ad}^0 V$ is a direct summand of $\mathrm{ad} V$, so $H^1(\Gamma, \mathrm{ad}^0 V) = 0$. This proves the first part above.

Let $G^0 \supset B \supset T$ denote a Borel and maximal torus defined over \mathbb{F}_l . Proposition 7 also shows that $|\langle \mu, \alpha^\vee \rangle| < (l - 1)/2$ for all weights μ of $T_{/\overline{\mathbb{F}}_l}$ on V and all $\alpha \in \Delta$. In particular, all dominant weights of $T_{/\overline{\mathbb{F}}_l}$ on V and $\mathrm{ad} V$ are restricted. Note that if W is a semisimple $G_{/\overline{\mathbb{F}}_l}^0$ -module such that all dominant weights of $T_{/\overline{\mathbb{F}}_l}$ on W are restricted, then every $G^0(\mathbb{F}_l)$ -submodule of W is also a $G_{/\overline{\mathbb{F}}_l}^0$ -submodule. We apply this first to V (which is semisimple as $G_{/\overline{\mathbb{F}}_l}^0$ -module, since r is semisimple), so the W_i are $G_{/\overline{\mathbb{F}}_l}^0$ -submodules. By Proposition 8 of [Ser94] we see that $\mathrm{ad} V = \bigoplus_{i,j} \mathrm{Hom}(W_i, W_j)$ is a semisimple $G_{/\overline{\mathbb{F}}_l}^0$ -module. (Note

that $\dim W_i + \dim W_j < l + 2$.) Thus every $G^0(\mathbb{F}_l)$ -submodule of $\text{ad } V$ is also a $G^0_{/\mathbb{F}_l}$ -submodule.

By Lemma 3 (applied with Δ_* the set of simple coroots), the $\overline{\mathbb{F}}_l$ -linear span of the image of $T(\mathbb{F}_l)$ in $\text{ad } V$ equals the $\overline{\mathbb{F}}_l$ -linear span of the image of $T(\overline{\mathbb{F}}_l)$. Thus the $G^0(\mathbb{F}_l)$ -submodule of $\text{ad } V$ generated by the $\overline{\mathbb{F}}_l$ -linear span of $r(H)$ equals the $G^0(\overline{\mathbb{F}}_l)$ -submodule generated by $r(T(\overline{\mathbb{F}}_l)H)$. By Lemma 8 (noting that $\mathcal{T}(\overline{\mathbb{F}}_l) = T(\overline{\mathbb{F}}_l)H$) it follows that $r(H)$ spans $\text{ad } V$. As $r(H) \subset \Gamma^{\text{ss}}$, this completes the proof. \square

REFERENCES

- [Bor91] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- [Car93] Roger W. Carter. *Finite groups of Lie type*. Wiley Classics Library. John Wiley & Sons Ltd., Chichester, 1993. Conjugacy classes and complex characters, Reprint of the 1985 original, A Wiley-Interscience Publication.
- [CR62] Charles W. Curtis and Irving Reiner. *Representation theory of finite groups and associative algebras*. Pure and Applied Mathematics, Vol. XI. Interscience Publishers, a division of John Wiley & Sons, New York-London, 1962.
- [GLS98] Daniel Gorenstein, Richard Lyons, and Ronald Solomon. *The classification of the finite simple groups. Number 3. Part I. Chapter A*, volume 40 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1998. Almost simple K -groups.
- [Gur99] Robert M. Guralnick. Small representations are completely reducible. *J. Algebra*, 220(2):531–541, 1999.
- [Hum06] James E. Humphreys. *Modular representations of finite groups of Lie type*, volume 326 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2006.
- [Hur82] James F. Hurley. A note on the centers of Lie algebras of classical type. In *Lie algebras and related topics (New Brunswick, N.J., 1981)*, volume 933 of *Lecture Notes in Math.*, pages 111–116. Springer, Berlin, 1982.
- [Jan03] Jens Carsten Jantzen. *Representations of algebraic groups*, volume 107 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2003.
- [Kup] G. Kuperberg. Denseness and Zariski denseness of Jones braid representations. Preprint.
- [LP] Michael Larsen and Richard Pink. Finite subgroups of algebraic groups. Preprint.
- [Mag54] Wilhelm Magnus. On the exponential solution of differential equations for a linear operator. *Comm. Pure Appl. Math.*, 7:649–673, 1954.
- [Mil] James S. Milne. Algebraic groups, Lie groups, and their arithmetic subgroups. Accessible from www.jmilne.org.
- [Ser92] Jean-Pierre Serre. *Lie algebras and Lie groups*, volume 1500 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, second edition, 1992. 1964 lectures given at Harvard University.

- [Ser94] Jean-Pierre Serre. Sur la semi-simplicité des produits tensoriels de représentations de groupes. *Invent. Math.*, 116(1-3):513–530, 1994.
- [Spr06] T. A. Springer. Twisted conjugacy in simply connected groups. *Transform. Groups*, 11(3):539–545, 2006.
- [Spr09] T. A. Springer. *Linear algebraic groups*. Modern Birkhäuser Classics. Birkhäuser Boston Inc., Boston, MA, second edition, 2009.
- [Ste61] Robert Steinberg. Automorphisms of classical Lie algebras. *Pacific J. Math.*, 11:1119–1129, 1961.
- [Ste68a] Robert Steinberg. *Endomorphisms of linear algebraic groups*. Memoirs of the American Mathematical Society, No. 80. American Mathematical Society, Providence, R.I., 1968.
- [Ste68b] Robert Steinberg. *Lectures on Chevalley groups*. Yale University, New Haven, Conn., 1968. Notes prepared by John Faulkner and Robert Wilson.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA,
LOS ANGELES, CA 90089-2532, USA

E-mail address: guralnic@usc.edu

INSTITUTE FOR ADVANCED STUDY, EINSTEIN DRIVE, PRINCETON, NJ 08540,
USA

E-mail address: herzig@math.ias.edu

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, 1 OXFORD STREET,
CAMBRIDGE, MA 02138, USA

E-mail address: rtaylor@math.harvard.edu

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, 1 OXFORD STREET,
CAMBRIDGE, MA 02138, USA

E-mail address: thorne@math.harvard.edu