

1. Let  $K = \mathbf{Q}(\sqrt{26})$  and let  $\varepsilon = 5 + \sqrt{26}$ . Use Dedekind's theorem to show that the ideal equations

$$\langle 2 \rangle = \langle 2, \varepsilon + 1 \rangle^2, \quad \langle 5 \rangle = \langle 5, \varepsilon + 1 \rangle \langle 5, \varepsilon - 1 \rangle, \quad \langle \varepsilon + 1 \rangle = \langle 2, \varepsilon + 1 \rangle \langle 5, \varepsilon + 1 \rangle$$

hold in  $K$ . Using Minkowski's bound, show that the class number of  $K$  (i.e. the cardinality of the ideal class group  $\text{Cl}(\mathcal{O}_K)$ ) is 2. Verify that  $\varepsilon$  is the fundamental unit. Deduce that all solutions in integers  $x, y$  to the equation  $x^2 - 26y^2 = \pm 10$  are given by  $x + \sqrt{26}y = \pm \varepsilon^n(\varepsilon \pm 1)$  for  $n \in \mathbf{Z}$ .

2. Find the factorisations into prime ideals of  $\langle 2 \rangle$  and  $\langle 3 \rangle$  in  $K = \mathbf{Q}(\sqrt{-23})$ . Verify that  $\langle \omega \rangle = \langle 2, \omega \rangle \langle 3, \omega \rangle$  where  $\omega = \frac{1}{2}(1 + \sqrt{-23})$ . Prove that  $K$  has class number 3.

3. Find the factorisations into prime ideals of  $\langle 2 \rangle$ ,  $\langle 3 \rangle$  and  $\langle 5 \rangle$  in  $K = \mathbf{Q}(\sqrt{-71})$ . Verify that

$$\langle \alpha \rangle = \langle 2, \alpha \rangle \langle 3, \alpha \rangle^2 \quad \text{and} \quad \langle \alpha + 2 \rangle = \langle 2, \alpha \rangle^3 \langle 3, \alpha - 1 \rangle$$

where  $\alpha = \frac{1}{2}(1 + \sqrt{-71})$ . Find an element of  $\mathcal{O}_K$  with norm  $2^a \cdot 3^b \cdot 5$  for some  $a, b \geq 0$ . Hence prove that the class group of  $K$  is cyclic and find its order.

4. Compute the ideal class group of  $\mathbf{Q}(\sqrt{d})$  for  $d = -30, -13, -10, 19$  and  $65$ .

5. (1) Find the fundamental unit in  $\mathbf{Q}(\sqrt{3})$ . Determine all the integer solutions of the equations  $x^2 - 3y^2 = m$  for  $m = -1, 13$  and  $121$ .

(2) Find the fundamental unit in  $\mathbf{Q}(\sqrt{10})$ . Determine all the integer solutions of the equations  $x^2 - 10y^2 = m$  for  $m = -1, 6$  and  $7$ .

6. Find all integer solutions of the equations  $y^2 = x^3 - 13$  and  $y^2 = x^5 - 10$ .

7. Show that  $\mathbf{Q}(\sqrt{-d})$  has class number 1 for  $d = 1, 2, 3, 7, 11, 19, 43, 67, 163$ .

8. Let  $K = \mathbf{Q}(\sqrt{-d})$  where  $d > 3$  is a square-free integer.

(1) Show that if  $\mathcal{O}_K$  is Euclidean, then it contains a principal ideal of norm 2 or 3. [Hint: Suppose that  $\phi : \mathcal{O}_K - \{0\} \rightarrow \mathbf{N}$  is a Euclidean function. Then choose  $x \in \mathcal{O}_K - \{0, \pm 1\}$  with  $\phi(x)$  minimal.]

(2) Use your answer to Question 7 to give an example where  $\mathcal{O}_K$  is a PID, but is not Euclidean.

9. Let  $K = \mathbf{Q}(\alpha)$  where  $\alpha$  is a root of  $f(X) = X^3 - 7X - 1$ . [Note that  $\text{disc}(f) = 5 \times 269$  is square-free.] Compute  $N_{K/\mathbf{Q}}(n + \alpha)$  for  $|n| \leq 3$ . Hence show that  $\langle 5 \rangle = P_1^2 P_2$  and  $\langle 7 \rangle = Q_1 Q_2 Q_3$  where the  $P_i$  and  $Q_j$  are distinct principal prime ideals of  $\mathcal{O}_K$ . Find units generating a subgroup of  $\mathcal{O}_K^\times$  of finite index. [Hint: You can show that the units you have found are independent by considering their images in  $\mathcal{O}_K/7\mathcal{O}_K \cong \mathbf{F}_7 \times \mathbf{F}_7 \times \mathbf{F}_7$ .]

10. Let  $K = \mathbf{Q}(\sqrt{d})$  where  $d \neq 0, 1$  is a square-free integer. Describe the ring  $\mathcal{O}_K/2\mathcal{O}_K$  as explicitly as you can. [The answer depends on  $d \pmod{8}$ .] Show that  $\mathbf{Z}[\sqrt{d}]^\times \subset \mathcal{O}_K^\times$  has index 1 or 3. Give an example where the index is 3.

11. Let  $p$  be an odd prime and let  $\zeta_p = e^{2\pi i/p}$ .

(1) Show that  $\mathbf{Q}(\zeta_p)$  contains a quadratic field with discriminant  $\pm p$ . How does the sign depend on  $p$ ?

(2) Show using the Minkowski bound that  $\mathbf{Z}[\zeta_p]$  is a UFD for  $p = 5$  and  $p = 7$ .

- 12.** Let  $K = \mathbf{Q}(\alpha)$  where  $\alpha$  is a root of  $f(X) = X^3 - 3X + 1$ .
- (1) Show that  $f$  is irreducible over  $\mathbf{Q}$  and compute its discriminant.
  - (2) Show that  $3\mathcal{O}_K = P^3$  where  $P = \langle \alpha + 1 \rangle$  is a prime ideal in  $\mathcal{O}_K$  with residue field  $\mathbf{F}_3$ . Deduce that  $\mathcal{O}_K = \mathbf{Z}[\alpha] + 3\mathcal{O}_K$ .
  - (3) Show that  $\mathcal{O}_K = \mathbf{Z}[\alpha]$ . Compute the class group of  $K$ .
- 13.** Let  $K = \mathbf{Q}(e^{2\pi i/23})$ .
- (1) Show that there are distinct prime ideals  $Q, Q'$  of  $\mathcal{O}_K$  such that  $\langle 2 \rangle = QQ'$  and  $N(Q) = N(Q') = 2^{11}$ . [You may use the fact from Part II Galois Theory that any finite field of order  $p^n$  contains a unique subfield of order  $p^d$  for each  $d|n$ .]
  - (2) Using your answer to Question 2, deduce that the class number of  $K$  is divisible by 3.
- 14.** Let  $B_{r,s}(t) = \{(y_1, \dots, y_r, z_1, \dots, z_s) \in \mathbf{R}^r \times \mathbf{C}^s \mid \sum |y_i| + 2 \sum |z_j| \leq t\}$ . Show that

$$\text{Vol } B_{r+1,s}(t) = \int_{-t}^t \text{Vol } B_{r,s}(t - |y|) dy,$$

and

$$\text{Vol } B_{r,s+1}(t) = \int_{|z| \leq t/2} \text{Vol } B_{r,s}(t - 2|z|) dz.$$

Hence show by induction that

$$\text{Vol } B_{r,s}(t) = 2^r \left(\frac{\pi}{2}\right)^s \frac{t^{r+2s}}{(r+2s)!}.$$

[You should do the second integral by choosing polar coordinates,  $z = \rho e^{i\theta}$ .]