## Computer Science Tripos 2002 Paper 1 Question 8

Thomas Forster

February 4, 2021

Let  $\Omega$  be a set. Write  $\mathcal{P}(\Omega)$  for its power set. Recall the definition of the intersection of  $\mathcal{B} \subseteq \mathcal{P}(\Omega)$ :

$$\bigcap_{B \in \mathcal{B}} \mathcal{B} \ = \ \{x \in \Omega : (\forall B \in \mathcal{B})(x \in B)\}$$

- (a) Let  $\mathcal{B} \subseteq \mathcal{P}(\Omega)$  and  $\mathcal{C} \subseteq \mathcal{P}(\Omega)$ .
  - (i) Prove that


$$(\bigcap_{B \in \mathcal{B}} B) \cup (\bigcap_{C \in \mathcal{C}} C) \subseteq \bigcap_{(B,C) \in \mathcal{B} \times \mathcal{C}} (B \cup C)$$

(ii) Prove that

$$\bigcap_{(B,C)\in\mathcal{B}\times\mathcal{C}}(B\cup C)\subseteq (\bigcap_{B\in\mathcal{B}}B)\cup (\bigcap_{C\in\mathcal{C}}C)$$

(b) This part seems to have caused problems for some. Let's have a look.

We are contemplating relations that hold between elements of  $\Omega$  and subsets of  $\Omega$ . An example of the sort of thing the examiner has in mind is the relation that a point y in the plane bears to a (typically non-convex) region X when y is in the convex hull of X.



The idea is that y is one of the points you have to "add" to obtain something convex. (Check that you know what a convex set is, as i'm going to procede on the assumption that you do, and use it as a—one hopes!—illuminating illustration)

What is  $\mathcal{R}$ ?  $\mathcal{A}$  is an intersection-closed family of subsets of  $\Omega$ . (As it might be, the collection of convex subsets of the plane). We are told that it is the relation that relates y to X whenever anything in A that extends X also contains y. In our illustration—where  $\mathcal{A}$  is the collection of convex subsets of the plane— $\mathcal{R}$  is the relation that hold between X and y whenever y is in the convex hull of X. (If you don't already know the meaning of the expression "convex hull" you can probably guess it from the news that, in the picture above, y is in the convex hull of X.) Certainly in this case any set that is  $\mathcal{R}$ -closed is convex.

Assume C is  $\mathcal{R}$ -closed. That is to say

$$\forall (X,y) \in \mathcal{R}. X \subseteq C \to y \in C \tag{1}$$

(That's in their notation: i'd've written it  $(\forall \langle X, y \rangle \in \mathcal{R})(X \subseteq C \to y \in C)$  which (i think) makes the scoping clearer.)

But  $\mathcal{R} = \{(X, y) \in \mathcal{P}(\Omega) \times \Omega : (\forall A \in \mathcal{A})(X \subseteq A \to y \in A)\}$ . Substituting this for ' $\mathcal{R}$ ' in (1) we obtain

$$\forall (X,y) \in \{(X,y) \in \mathcal{P}(\Omega) \times \Omega : (\forall A \in \mathcal{A})(X \subseteq A \to y \in A)\}.X \subseteq C \to y \in C$$
(2)

which reduces to

$$(\forall X, y)[(\forall A \in \mathcal{A})(X \subseteq A \to y \in A) \land X \subseteq C. \to y \in C] \tag{3}$$

The examiners suggest you should consider the set  $\{A \in \mathcal{A} : C \subseteq A\}$ . I think they want you to look at  $\bigcap \{A \in \mathcal{A} : C \subseteq A\}$ .

If you've followed the action this far you would probably think of this anyway, since this is a set that you know must be in  $\mathcal{A}$  and it seems to stand an outside chance of being equal to C. So let's look again at (3) to see if it does, in fact, tell us that  $\bigcap \{A \in \mathcal{A} : C \subseteq A\}$  is C.

And—of course—it does. First we instantiate 'X' to 'C' in (3) to obtain:

$$\forall y [(\forall A \in \mathcal{A})(C \subseteq A \to y \in A) \to y \in C] \tag{4}$$

Now let y be an arbitrary member of  $\bigcap \{A \in \mathcal{A} : C \subseteq A\}$ . That means that y satisfies the antecedent of 4. So it satisfies the consequent of 4 as well. So we have proved that  $\bigcap \{A \in \mathcal{A} : C \subseteq A\}$  is a subset of C. It was always a superset of C, so it is equal to C. So  $C \in \mathcal{A}$  as desired.